

**CHILDRENS PLUS MULTI SYMPTOM COLD- acetaminophen, chlorpheniramine maleate, dextromethorphan hydrobromide, phenylephrine hydrochloride suspension**  
**Care One (American Sales Company)**

-----

**Drug Facts**

**Active ingredients (in each 5 mL)**

**Acetaminophen 160 mg**

Chlorpheniramine maleate 1 mg

Dextromethorphan HBr 5 mg

Phenylephrine HCl 2.5 mg

**Purposes**

**Pain reliever/fever reducer**

Antihistamine

Cough suppressant

Nasal decongestant

**Uses**

- temporarily relieves the following cold/flu symptoms:
  - minor aches and pains
  - cough
  - headache
  - sore throat
  - sneezing and runny nose
  - nasal congestion
  - stuffy nose
- temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses (10 mL) in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters

- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- to make a child sleepy
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.
- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product

### **Ask a doctor before use if your child has**

- a breathing problem such as chronic bronchitis
- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus)
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- liver disease

### **Ask a doctor or pharmacist before use if your child is taking**

- sedatives or tranquilizers
- the blood thinning drug warfarin

### **When using this product**

- **do not exceed recommended dose (see overdose warning)**
- excitability may occur, especially in children
- marked drowsiness may occur
- sedatives and tranquilizers may increase drowsiness

### **Stop use and ask a doctor if**

- new symptoms occur
- fever gets worse or lasts for more than 3 days
- redness or swelling is present
- pain, nasal congestion or cough gets worse or lasts for more than 5 days
- nervousness, dizziness or sleeplessness occurs
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

### **Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

### **Directions**

- **this product does not contain directions or complete warnings for adult use**
- mL=milliliter
- shake well before using
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- do not give more than 5 doses in any 24-hours period
- if needed, repeat dose every 4 hours while symptoms last
- find the right dose on chart below. If possible, use weight to dose, otherwise, use age.

Weight (lb)	Age (yr)	Dose (mL)
48-95	6-11	10 mL
36-47	4-5	do not use unless directed by a doctor
under 36	under 4	do not use

### **Other Information**

- store between 20-25°C (68-77°F)
- do not refrigerate.

### **Inactive ingredients**

anhydrous citric acid, carboxymethylcellulose sodium, D&C red #33, FD&C blue #1, FD&C red #40, flavor, glycerin, microcrystalline cellulose, purified water, sodium benzoate, sorbitol, sucrose, xanthan gum

### **Questions or comments?**

Call **1-877-753-3935 Monday-Friday 9AM-5PM EST**

### **Principal Display Panel**

For Ages 6-11 Years

### **Children's Multi-Symptom Cold**

Pain Relief / Fever Reducer - ACETAMINOPHEN, 160 mg

Antihistamine - Chlorpheniramine Maleate, 1 mg

Cough Suppressant - Dextromethorphan HBr, 5 mg

Nasal Decongestant - Phenylephrine HCl, 2.5 mg

FOR RELIEF OF:

- Fever & Sore Throat
- Sneezing and Runny Nose
- Cough
- Stuffy Nose

Oral Suspension

Dosage Cup Provided

Grape Flavored

FL OZ (mL)

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

DISTRIBUTED BY:

FOODHOLD U.S.A., LLC

LANDOVER, MD 20785

1-877-846-9949

**Package Label**

**Drug Facts (continued)**

- nervousness, dizziness or sleeplessness occurs
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

**Directions**

- this product does not contain directions or complete warnings for adult use
- mL = milliliter
- shake well before using
- do not give more than 5 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- if needed, repeat dose every 4 hours while symptoms last
- find the right dose on chart below. If possible, use weight to dose; otherwise, use age.

Weight (lb)	Age (yr)	Dose (mL)
48-95	6-11	10 mL
36-47	4-5	do not use unless directed by a doctor
under 36	under 4	do not use

**Other information**

- store between 20-25°C (68-77°F). Do not refrigerate.

**Inactive ingredients** carboxymethylcellulose sodium, citric acid, D&C red #33, FD&C blue #1, FD&C red #40, flavor, glycerin, microcrystalline cellulose, purified water, sodium benzoate, sorbitol, sucrose, xanthan gum

**Questions or comments?**  
Call 1-877-753-3835 Monday-Friday 9AM-5PM EST

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

**PARENTS:**  
Learn about teen medicine abuse  
www.StopMedicineAbuse.org

DISTRIBUTED BY FOODHOLD U.S.A., LLC  
LANDOVER, MD 20785 • 1-877-846-9949  
© 2016 S&S Brands, LLC  
Quality guaranteed or your money back.

NDC 41520-704-04

**CAREone®**

For Ages 6 - 11 Years

**Children's**

**MULTI-SYMPTOM COLD**

Pain Relief/Fever Reducer - ACETAMINOPHEN, 160 mg  
Antihistamine - Chlorpheniramine Maleate, 1 mg  
Cough Suppressant - Dextromethorphan HBr, 5 mg  
Nasal Decongestant - Phenylephrine HCl, 2.5 mg

**FOR RELIEF OF:**

- Fever & Sore Throat • Stuffy Nose
- Sneezing & Runny Nose • Cough

Oral Suspension  
Dosing Cup Provided



**Grape Flavored**

**4 FL OZ (118mL)**

OUR PHARMACISTS RECOMMEND

NDC 41520-704-04

**CAREone®**

For Ages 6 - 11 Years

**Children's**

**MULTI-SYMPTOM COLD**

Pain Relief/Fever Reducer - ACETAMINOPHEN, 160 mg  
Antihistamine - Chlorpheniramine Maleate, 1 mg  
Cough Suppressant - Dextromethorphan HBr, 5 mg  
Nasal Decongestant - Phenylephrine HCl, 2.5 mg

**FOR RELIEF OF:**

- Fever & Sore Throat • Stuffy Nose
- Sneezing & Runny Nose • Cough

Oral Suspension  
Dosing Cup Provided



**Grape Flavored**

**4 FL OZ (118mL)**

OUR PHARMACISTS RECOMMEND

**Drug Facts**

**Active ingredients (in each 5 mL)**

Acetaminophen 160 mg	Pain reliever/fever reducer
Chlorpheniramine maleate 1 mg	Antihistamine
Dextromethorphan HBr 5 mg	Cough suppressant
Phenylephrine HCl 2.5 mg	Nasal decongestant

**Purposes**

- temporarily relieves the following cold/flu symptoms
- minor aches and pains
- sore throat
- sneezing and runny nose
- headache
- nasal congestion
- stuffy nose
- temporarily reduces fever

**Uses**

- temporarily relieves the following cold/flu symptoms
- minor aches and pains
- sore throat
- sneezing and runny nose
- headache
- nasal congestion
- stuffy nose
- temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses (10 mL) in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- to make a child sleep
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product.

**Ask a doctor before use if your child has**

- liver disease
- glaucoma
- thyroid disease
- diabetes
- heart disease
- high blood pressure
- a breathing problem such as chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with asthma

**Ask a doctor or pharmacist before use if your child is taking**

- the blood thinning drug warfarin
- sedatives or tranquilizers

**When using this product**

- do not exceed recommended dose (see Overdose warning)
- excitability may occur, especially in children
- marked drowsiness may occur
- sedatives and tranquilizers may increase drowsiness

**Stop use and ask a doctor if**

- new symptoms occur
- redness or swelling is present
- pain, nasal congestion or cough gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days

S-949 1016



3 41520 31888 7

PLD-1253C FC004119

Lot No.:  
Exp. Date:

### CAREONE Childrens Multi-Symptom Cold Grape Flavor

## CHILDRENS PLUS MULTI SYMPTOM COLD

acetaminophen, chlorpheniramine maleate, dextromethorphan hydrobromide, phenylephrine hydrochloride suspension

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:41520-704
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL
<b>CHLORPHENIRAMINE MALEATE</b> (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE -	CHLORPHENIRAMINE	1 mg

UNII:3U6IO1965U)	MALEATE	in 5 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	5 mg in 5 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	2.5 mg in 5 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>CARBOXYMETHYLCELLULOSE SODIUM</b> (UNII: K679OBS311)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>D&amp;C RED NO. 33</b> (UNII: 9DBA0SBB0L)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCROSE</b> (UNII: C151H8M554)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

### Product Characteristics

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	GRAPE	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41520-704-04	1 in 1 BOX	10/31/2015	
1		118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	10/31/2015	

**Labeler** - Care One (American Sales Company) (809183973)