

ASPIRIN ENTERIC COATED- aspirin tablet, delayed release
CHAIN DRUG MARKETING ASSOCIATION INC

Quality Choice 44-600A

Active ingredient (in each tablet)

Aspirin 81 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

for the temporary relief of minor aches and pains or as recommended by your doctor.
Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take more or for a longer time than directed
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older
- have 3 or more alcoholic drinks every day while using this product

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- you have asthma
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - vomit blood
 - have bloody or black stools
 - feel faint
 - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°C-30°C (59°F-86°F)

- see end flap for expiration date and lot number

Inactive ingredients

corn starch, D&C yellow #10, FD&C yellow #6, hypromellose, methacrylic acid, microcrystalline cellulose, polydextrose, polyethylene glycol, shellac wax, silica, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments?

1-800-426-9391

Principal Display Panel

NDC 63868-363-20

**QC®
QUALITY
CHOICE**

**†Compare to the
Active Ingredient in
BAYER® Low Dose Aspirin**

**Enteric Coated
Aspirin 81 mg**

Low Dose

Pain Reliever (**NSAID**)
Aspirin Regimen**

120 Tablets (81 mg Each)

actual size

**Talk to your doctor or other healthcare provider
before using this product for your heart.

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

†This product is not manufactured or distributed by
Bayer AG, owner of the registered trademark
BAYER® Low Dose Aspirin.
50844 REV0122D60032

**SATISFACTION
GUARANTEED
100%
QC**

Distributed by CDMA, Inc.

43157 W. Nine Mile
 Novi, MI 48375
 www.qualitychoice.com
 Questions:800-935-2362

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Drug Facts
Active ingredient (in each tablet)
 Aspirin 81 mg (NSAID)*
 *nonsteroidal anti-inflammatory drug

Purpose
 Pain reliever

Uses
 for the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide last relief of headaches or other symptoms needing immediate relief.

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:
 ■ hives ■ facial swelling ■ shock ■ asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you
 ■ take more or for a longer time than directed
 ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

NDC 63868-363-20

Compare to the Active Ingredient in BAYER® Low Dose Aspirin

Enteric Coated Aspirin 81 mg Low Dose

Pain Reliever (NSAID) Aspirin Regimen**

120 Tablets (81 mg Each) actual size

Drug Facts (continued)
Questions or comments?
 1-800-426-9391

*This product is not manufactured or distributed by Bayer AG, owner of the registered trademark BAYER® Low Dose Aspirin.
 50844 REV0122D60032

Distributed by CDMA, Inc.
 43157 W Nine Mile
 Novi, MI 48375
 www.qualitychoice.com
 Questions: 800-935-2362

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Drug Facts (continued)
Questions or comments?
 1-800-426-9391

*This product is not manufactured or distributed by Bayer AG, owner of the registered trademark BAYER® Low Dose Aspirin.
 50844 REV0122D60032

Distributed by CDMA, Inc.
 43157 W Nine Mile
 Novi, MI 48375
 www.qualitychoice.com
 Questions: 800-935-2362

Drug Facts (continued)
 If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 ■ do not take more than directed
 ■ drink a full glass of water with each dose
 ■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
 ■ children under 12 years: do not use unless directed by a doctor

Other information
 ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
 ■ see end flap for expiration date and lot number

Inactive ingredients
 corn starch, D&C yellow #10, FD&C yellow #6, hypromellose, methacrylic acid, microcrystalline cellulose, polydextrose, polyethylene glycol, shellac wax, silica, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate

Drug Facts (continued)
 ■ have had stomach ulcers or bleeding problems
 ■ take a blood thinning (anticoagulant) or steroid drug
 ■ are age 60 or older
 ■ have 3 or more alcoholic drinks every day while using this product

Do not use
 ■ if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if
 ■ you have ever had an allergic reaction to this product or any of its ingredients
 ■ you have asthma
 ■ stomach bleeding warning applies to you
 ■ you have a history of stomach problems, such as heartburn
 ■ you are taking a diuretic
 ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug for
 ■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if
 ■ an allergic reaction occurs. Seek medical help right away.
 ■ you experience any of the following signs of stomach bleeding:
 ■ vomit blood
 ■ have bloody or black stools
 ■ feel faint
 ■ have stomach pain that does not get better
 ■ ringing in the ears or a loss of hearing occurs
 ■ pain gets worse or lasts more than 10 days
 ■ fever gets worse or lasts more than 3 days
 ■ redness or swelling is present
 ■ new symptoms occur

These could be signs of a serious condition.

Drug Facts (continued)
 ■ have had stomach ulcers or bleeding problems
 ■ take a blood thinning (anticoagulant) or steroid drug
 ■ are age 60 or older
 ■ have 3 or more alcoholic drinks every day while using this product

Do not use
 ■ if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if
 ■ you have ever had an allergic reaction to this product or any of its ingredients
 ■ you have asthma
 ■ stomach bleeding warning applies to you
 ■ you have a history of stomach problems, such as heartburn
 ■ you are taking a diuretic
 ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug for
 ■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if
 ■ an allergic reaction occurs. Seek medical help right away.
 ■ you experience any of the following signs of stomach bleeding:
 ■ vomit blood
 ■ have bloody or black stools
 ■ feel faint
 ■ have stomach pain that does not get better
 ■ ringing in the ears or a loss of hearing occurs
 ■ pain gets worse or lasts more than 10 days
 ■ fever gets worse or lasts more than 3 days
 ■ redness or swelling is present
 ■ new symptoms occur

These could be signs of a serious condition.

Drug Facts (continued)
 ■ have had stomach ulcers or bleeding problems
 ■ take a blood thinning (anticoagulant) or steroid drug
 ■ are age 60 or older
 ■ have 3 or more alcoholic drinks every day while using this product

Do not use
 ■ if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if
 ■ you have ever had an allergic reaction to this product or any of its ingredients
 ■ you have asthma
 ■ stomach bleeding warning applies to you
 ■ you have a history of stomach problems, such as heartburn
 ■ you are taking a diuretic
 ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug for
 ■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if
 ■ an allergic reaction occurs. Seek medical help right away.
 ■ you experience any of the following signs of stomach bleeding:
 ■ vomit blood
 ■ have bloody or black stools
 ■ feel faint
 ■ have stomach pain that does not get better
 ■ ringing in the ears or a loss of hearing occurs
 ■ pain gets worse or lasts more than 10 days
 ■ fever gets worse or lasts more than 3 days
 ■ redness or swelling is present
 ■ new symptoms occur

These could be signs of a serious condition.

Drug Facts (continued)
 ■ have had stomach ulcers or bleeding problems
 ■ take a blood thinning (anticoagulant) or steroid drug
 ■ are age 60 or older
 ■ have 3 or more alcoholic drinks every day while using this product

Do not use
 ■ if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if
 ■ you have ever had an allergic reaction to this product or any of its ingredients
 ■ you have asthma
 ■ stomach bleeding warning applies to you
 ■ you have a history of stomach problems, such as heartburn
 ■ you are taking a diuretic
 ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug for
 ■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if
 ■ an allergic reaction occurs. Seek medical help right away.
 ■ you experience any of the following signs of stomach bleeding:
 ■ vomit blood
 ■ have bloody or black stools
 ■ feel faint
 ■ have stomach pain that does not get better
 ■ ringing in the ears or a loss of hearing occurs
 ■ pain gets worse or lasts more than 10 days
 ■ fever gets worse or lasts more than 3 days
 ■ redness or swelling is present
 ■ new symptoms occur

These could be signs of a serious condition.

Quality Choice 44-600A

ASPIRIN ENTERIC COATED
 aspirin tablet, delayed release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63868-363
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

Inactive Ingredients	
Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
METHACRYLIC ACID (UNII: 1CS02G8656)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SHELLAC (UNII: 46N107B71O)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

Product Characteristics			
Color	yellow	Score	no score
Shape	ROUND	Size	6mm
Flavor		Imprint Code	L
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63868-363-20	1 in 1 CARTON	05/01/2011	
1		120 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:63868-363-36	365 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	05/01/2011	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	05/01/2011	

Labeler - CHAIN DRUG MARKETING ASSOCIATION INC (011920774)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(63868-363)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(63868-363)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(63868-363) , pack(63868-363)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(63868-363)

Revised: 4/2024

CHAIN DRUG MARKETING ASSOCIATION INC