

CHILDRENS MOTRIN- ibuprofen suspension
Kenvue Brands LLC

Children's MOTRIN

Drug Facts

Active ingredient (in each 5 mL)

Ibuprofen 100 mg (NSAID) ¹

¹ nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

Warnings

Allergy alert

Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin,

ibuprofen, naproxen, or others)

- takes more or for a longer time than directed

Heart attack and stroke warning

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Sore throat warning

Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding:
 - feels faint
 - vomits blood
 - has bloody or black stools
 - has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling

- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed**
- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.
- if needed, repeat dose every **6-8 hours**
- do not use more than **4 times a day**
- replace original bottle cap to maintain child resistance

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL) *
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

* or as directed by a doctor

Other information

- each 5 mL contains: **sodium 2 mg**
- store between 20-25°C (68-77°F)
- **do not use if carton tape or bottle wrap imprinted with "Sealed For Your Safety" is broken or missing**

Inactive ingredients

acesulfame potassium, anhydrous citric acid, D&C yellow no. 10, FD&C red no. 40, flavors, glycerin, polysorbate 80, pregelatinized starch, purified water, sodium benzoate, sucrose, xanthan gum

Questions or comments?

Call **1-877-895-3665** (toll-free) or **215-273-8755** (collect)

PRINCIPAL DISPLAY PANEL

NDC 50580-601-22

For Ages 2 to 11 Years

***Children's
Motrin[®]***

contains Ibuprofen

Oral Suspension (NSAID)

100 mg per 5 mL

Pain Reliever / Fever Reducer

***Lasts up to
8 hours***

Alcohol Free

***Original
Berry Flavor***

1 fl oz (30 mL)

Children's Motrin
contains
Ibuprofen Oral Suspension (NSAID)

Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA
©J&JCI 2021 www.motrin.com 30050814

Original Berry Flavor

NDC 50580-601-22

For Ages 2 to 11 Years

Children's Motrin
contains **Ibuprofen Oral Suspension (NSAID)**
100 mg per 5 mL
Pain Reliever / Fever Reducer



Important: Read all product information before using. Keep this box for important information. This product is intended for use in children.

Drug Facts

Active ingredient (in each 5 mL) Purpose
Ibuprofen 100 mg (NSAID)*.....Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses temporarily: ■ relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
■ reduces fever

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
■ hives ■ facial swelling ■ asthma (wheezing)
■ shock ■ skin reddening ■ rash ■ blisters
If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child: ■ has had stomach ulcers or bleeding problems
■ takes a blood thinning (anticoagulant) or steroid drug
■ takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ takes more or for a longer time than directed
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Drug Facts (continued)

Do not use ■ if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
■ right before or after heart surgery

Ask a doctor before use if ■ stomach bleeding warning applies to your child ■ child has a history of stomach problems, such as heartburn ■ child has problems or serious side effects from taking pain relievers or fever reducers
■ child has not been drinking fluids ■ child has lost a lot of fluid due to vomiting or diarrhea ■ child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke ■ child has asthma ■ child is taking a diuretic

Ask a doctor or pharmacist before use if the child is
■ under a doctor's care for any serious condition
■ taking any other drug

When using this product
■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if ■ child experiences any of the following signs of stomach bleeding: ■ feels faint
■ vomits blood ■ has bloody or black stools
■ has stomach pain that does not get better ■ child has symptoms of heart problems or stroke: ■ chest pain
■ trouble breathing ■ weakness in one part or side of body
■ slurred speech ■ leg swelling ■ the child does not get any relief within first day (24 hours) of treatment ■ fever or pain gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions ■ this product does not contain directions or complete warnings for adult use ■ do not give more than directed ■ shake well before using ■ mL = milliliter
■ find right dose on chart. If possible, use weight to dose; →

Drug Facts (continued)

otherwise use age. ■ use only enclosed dosing cup. Do not use any other dosing device. ■ if needed, repeat dose every 6-8 hours ■ do not use more than 4 times a day
■ replace original bottle cap to maintain child resistance

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

*or as directed by a doctor

Other information ■ each 5 mL contains: sodium 2 mg
■ store between 20-25°C (68-77°F) ■ do not use if carton tape or bottle wrap imprinted with "Sealed For Your Safety" is broken or missing

Inactive ingredients acesulfame potassium, anhydrous citric acid, D&C yellow no. 10, FD&C red no. 40, flavors, glycerin, polysorbate 80, pregelatinized starch, purified water, sodium benzoate, sucrose, xanthan gum

Questions or comments?
Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)



USA
746452
904

LOT: _____
EXP: _____

CHILDRENS MOTRIN
ibuprofen suspension

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50580-601
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	100 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SUCROSE (UNII: C151H8M554)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	orange	Score	
Shape		Size	
Flavor	BERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50580-601-08	2 in 1 PACKAGE	01/30/2002	
1	NDC:50580-601-04	1 in 1 CARTON		
1		120 mL in 1 BOTTLE, PLASTIC; Type 1: Convenience Kit of Co-Package		
2	NDC:50580-601-21	1 in 1 CARTON	03/01/2012	
2		120 mL in 1 BOTTLE, PLASTIC; Type 1: Convenience Kit of Co-Package		
3	NDC:50580-601-22	1 in 1 CARTON	03/01/2012	
3		30 mL in 1 BOTTLE, PLASTIC; Type 1: Convenience Kit of Co-Package		
4	NDC:50580-601-20	1 in 1 CARTON	03/01/2012	
4		120 mL in 1 BOTTLE, PLASTIC; Type 1: Convenience Kit of Co-Package		
5	NDC:50580-601-25	2 in 1 PACKAGE	07/10/2012	
5		1 in 1 CARTON		
5		120 mL in 1 BOTTLE, PLASTIC; Type 1: Convenience Kit of Co-Package		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020516	09/01/1995	

Labeler - Kenvue Brands LLC (118772437)

Revised: 11/2024

Kenvue Brands LLC