

**COLD AND FLU DAYTIME- acetaminophen, dextromethorphan hbr,
phenylephrine hcl solution
Bi-Mart**

Bi-Mart 44-012

Active ingredients (in each 15 mL)

Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer
Cough suppressant
Nasal decongestant

Uses

- temporarily relieves common cold and flu symptoms:
 - sore throat
 - nasal congestion
 - fever
 - headache
 - minor aches and pains
 - cough due to minor throat and bronchial irritation

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a

doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if the user has

- difficulty in urination due to enlargement of the prostate gland
- cough that occurs with too much phlegm (mucus)
- liver disease
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- heart disease
- diabetes
- thyroid disease
- high blood pressure

Ask a doctor or pharmacist before use if the user is

taking the blood thinning drug warfarin.

When using this product

do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed**
- mL = milliliter; FL OZ = fluid ounce
- only use the dose cup provided
- do not take more than 5 doses per 24 hours

adults and children 12 years and over	30 mL every 4 hours
children 6 to under 12 years	15 mL every 4 hours
children under 6 years	do not use

Other information

- **each 15 mL contains:** sodium 13 mg
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients

anhydrous citric acid, FD&C yellow #6, flavors, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate dihydrate, sodium metabisulfite, sodium saccharin, sorbitol, sucralose

Questions or comments?

1-800-935-6737

Principal display panel**BI-MART**

Compare to the active ingredients
in Vicks® DayQuil® Cold & Flu*

**DAYTIME
COLD & FLU**

Acetaminophen (Pain reliever/fever reducer)
Dextromethorphan HBr (Cough suppressant)
Phenylephrine HCl (Nasal decongestant)

Headache, Fever, Sore Throat,
Minor Aches & Pains,
Nasal Congestion,
Cough

Alcohol Free

Menthol Flavored

12 FL OZ (355 mL)

TAMPER EVIDENT: DO NOT USE IF PRINTED

NECK WRAP IS BROKEN OR MISSING

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

BI-MART UNCONDITIONAL MONEY BACK GUARANTEE

If you are not completely satisfied with your BI-MART QUALITY PRODUCT, regardless of the reason, return the unused portion and your entire purchase price will be cheerfully refunded.

*This product is not manufactured or distributed by The Procter & Gamble Company, distributor of Vicks® DayQuil® Cold & Flu.

50844 REV0722A01202

Distributed by:

BI-MART

Eugene, OR 97402

TAMPER EVIDENT: DO NOT USE IF PRINTED
NECK WRAP IS BROKEN OR MISSING



PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

BI-MART UNCONDITIONAL MONEY BACK GUARANTEE
If you are not completely satisfied with your BI-MART
QUALITY PRODUCT, regardless of the reason, return the
unused portion and your entire purchase price will be
cheerfully refunded.

*This product is not manufactured or
distributed by The Procter & Gamble
Company, distributor of Vicks®
DayQuil® Cold & Flu.

50844 REV0722A01202

Distributed by:
BI-MART
Eugene, OR 97402

3001000549

69526-012-11-1

No Print / No Varnish Area
Lot # and Exp. Info

BI-MART

Compare to the active ingredients
in Vicks® DayQuil® Cold & Flu*

DAYTIME

COLD & FLU

Acetaminophen (Pain reliever/fever reducer)
Dextromethorphan HBr (Cough suppressant)
Phenylephrine HCl (Nasal decongestant)

Headache, Fever, Sore Throat,
Minor Aches & Pains,
Nasal Congestion,
Cough



Menthol Flavored

Alcohol Free

12 FL OZ (355 mL)

Drug Facts

Active ingredients (in each 15 mL) Purpose

Acetaminophen 325 mg Pain reliever/fever reducer
Dextromethorphan HBr 10 mg Cough suppressant
Phenylephrine HCl 5 mg Nasal decongestant

Uses

temporarily relieves common cold and flu symptoms:
sore throat nasal congestion fever
headache minor aches and pains
cough due to minor throat and bronchial irritation

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

adult takes more than 4,000 mg of acetaminophen in 24 hours
child takes more than 5 doses in 24 hours
taken with other drugs containing acetaminophen

adult has 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening blisters rash

If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if the user has

difficulty in urination due to enlargement of the prostate gland
cough that occurs with too much phlegm (mucus) liver disease
persistent or chronic cough such as occurs with smoking, asthma, or emphysema heart disease diabetes thyroid disease
high blood pressure

PEEL HERE FOR MORE DRUG FACTS

Questions or comments? 1-800-935-6737

Inactive ingredients anhydrous citric acid, FD&C yellow #6, sodium saccharin, sorbitol, sucralose, sodium benzoate, sodium citrate dihydrate, sodium metabisulfite, flavors, glycerin, polyethylene glycol, propylene glycol, purified water.

Other information each 15 mL contains: sodium 13 mg
store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) use by expiration date on package

adults and children 12 years and over	30 mL every 4 hours
children 6 to under 12 years	15 mL every 4 hours
children under 6 years	do not use

Directions
do not take more than directed
mL = milliliter; FL OZ = fluid ounce
only use the dose cup provided
do not take more than 5 doses per 24 hours

could be signs of a serious condition.
cough comes back or occurs with rash or headache that lasts. These (children) or 7 days (adults)
pain, nasal congestion, or cough gets worse or lasts more than 5 days
redness or swelling is present new symptoms occur
fever gets worse or lasts more than 3 days
nervousness, dizziness, or sleeplessness occur

Stop use and ask a doctor if
When using this product do not exceed recommended dosage.
thinning drug warfarin.

Drug Facts (continued)
Ask a doctor or pharmacist before use if the user is taking the blood

Bi-Mart 44-012

COLD AND FLU DAYTIME

acetaminophen, dextromethorphan hbr, phenylephrine hcl solution

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:37835-526

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
SODIUM METABISULFITE (UNII: 4VON5FNS3C)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color	orange	Score	
Shape		Size	
Flavor	MENTHOL	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37835-526-30	354 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/18/2018	01/06/2023
2	NDC:37835-526-02	355 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/06/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/18/2018	

Labeler - Bi-Mart (027630078)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	manufacture(37835-526) , pack(37835-526)

Revised: 1/2024

Bi-Mart