

**DAYTIME NIGHTTIME COLD FLU RELIEF- acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl
EQUATE (Wal-Mart Stores, Inc.) (see also WAL-MART INC)**

DRUG FACTS

Active ingredients for Daytime (in each softgel)

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

Active ingredients for Nighttime (in each softgel)

Acetaminophen 325 mg

Dextromethorphan HBr 15 mg

Doxylamine succinate 6.25 mg

Purpose for Daytime

Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

Purpose for Nighttime

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Uses

DAYTIME

- temporarily relieves common cold and flu symptoms
 - cough due to minor throat and bronchial irritation
 - nasal congestion
 - headache
 - minor aches and pains
 - fever
 - sore throat

NIGHTTIME

- temporarily relieves common cold and flu symptoms
 - cough due to minor throat and bronchial irritation
 - sore throat
 - headache
 - minor aches and pains
 - fever
 - runny nose and sneezing

Warnings

DAYTIME

Liver warning: These products contain acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using these products

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

NIGHTTIME

Liver warning: This product contain acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or following by fever, headache, rash, nausea, vomiting, consult a doctor promptly.

Do not use

DAYTIME

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

NIGHTTIME

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease, or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

DAYTIME

- liver disease
- heart disease
- diabetes
- thyroid disease
- high blood pressure
- cough that occurs with too much phlegm (mucus)
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

NIGHTTIME

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

DAYTIME

taking the blood thinning drug warfarin

NIGHTTIME

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product,

DAYTIME

do not exceed recommended dosage

NIGHTTIME

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

DAYTIME

- nervousness, dizziness or sleeplessness occur
- pain, cough, and nasal congestion gets worse or lasts more than 7 days
- redness or swelling is present
- new symptoms occur
- fever gets worse or lasts more than 3 days
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

NIGHTTIME

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash headache that lasts.

These could be a signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

DAYTIME

- **do not take more than directed (see Overdose warning)**
- do not take more than 4 doses in 24 hours
- swallow whole; do not crush, chew, or dissolve
- adults and children 12 years and over; take 2 softgels with water every 4 hours.
- children under 12 years: do not use

NIGHTTIME

- **do not take more than directed (see Overdose warning)**

- do not take more than 4 doses in 24 hours
- swallow whole; do not crush, chew, or dissolve
- adults and children 12 years and over: take 2 softgels with water every 6 hours
- children under 12 years: do not use

Other information

- store between 15-30°C (59-86°F)
- avoid excessive heat

Inactive ingredients

DAYTIME

butylated hydroxyanisole, butylated hydroxytoluene, FD&C yellow #6, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

NIGHTTIME

D&C yellow #10, FD&C blue #1, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

Questions or comments?

Call **1-888-287-1915**

Principal Display Panel

Compare to the active ingredients in Vicks® DayQuil® and NyQuil® Cold & Flu LiquiCaps®

DAYTIME

Multi-Symptom

Cold & Flu Relief

Acetaminophen 325 mg- Pain reliever / Fever reducer

Dextromethorphan HBr 10 mg - Cough suppressant

Phenylephrine HCl 5 mg - Nasal decongestant

Relieves

- Aches, fever, sore throat
- Cough
- Nasal congestion

Non-drowsy

Alcohol-Free

Antihistamine-Free

Softgels**

(** Liquid-filled capsules)

NIGHTTIME

Multi-Symptom

Cold & Flu Relief

Acetaminophen 325 mg - Pain reliever / Fever reducer

Dextromethorphan HBr 15 mg - Cough suppressant

Doxylamine succinate 6.25 mg - Antihistamine

Relieves

- Aches, fever, sore throat
- Cough
- Runny nose, sneezing

Alcohol-free

Softgels**

(**Liquid-filled capsules)

When using Daytime and Nighttime products, carefully read the labeling to ensure correct dosing.

†This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, DayQuil, NyQuil®, and LiquiCaps® are registered

trademarks of the Procter and Gamble Company.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOW SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Distributed by:

Walmart Inc.,

Bentonville, AR 72716

Product Label



FREE AREA
COATING

<p>Drug Facts (continued)</p> <p>Do not use</p> <ul style="list-style-type: none"> ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ liver disease ■ heart disease ■ high blood pressure ■ diabetes ■ thyroid disease ■ kidney disease ■ trouble urinating due to an enlarged prostate gland ■ asthma, chronic bronchitis, or emphysema ■ trouble breathing or wheezing such as occurs with smoking, or <p>Ask a doctor or pharmacist before use if you are:</p> <ul style="list-style-type: none"> ■ taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers <p>When using this product, do not exceed recommended dosage.</p> <p>Stop use and ask a doctor if:</p> <ul style="list-style-type: none"> ■ drowsiness, dizziness, or sleepiness occur ■ pain, cough, or nasal congestion gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. <p>Keep out of reach of children. Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p>	<p>Drug Facts (continued)</p> <p>Do not use</p> <ul style="list-style-type: none"> ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ liver disease ■ glaucoma ■ trouble breathing or wheezing such as occurs with smoking, or ■ trouble urinating due to an enlarged prostate gland <p>Ask a doctor or pharmacist before use if you are:</p> <ul style="list-style-type: none"> ■ taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers <p>When using this product, do not exceed recommended dosage.</p> <p>Stop use and ask a doctor if:</p> <ul style="list-style-type: none"> ■ drowsiness, dizziness, or sleepiness occur ■ pain or cough gets worse or lasts more than 3 days ■ fever gets worse or lasts more than 3 days ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. <p>Keep out of reach of children. Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p>
---	--

<p>Drug Facts (continued)</p> <p>Do not use</p> <ul style="list-style-type: none"> ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ liver disease ■ heart disease ■ diabetes ■ thyroid disease ■ kidney disease ■ trouble urinating due to an enlarged prostate gland ■ asthma, chronic bronchitis, or emphysema ■ trouble breathing or wheezing such as occurs with smoking, or <p>Ask a doctor or pharmacist before use if you are:</p> <ul style="list-style-type: none"> ■ taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers <p>When using this product, do not exceed recommended dosage.</p> <p>Stop use and ask a doctor if:</p> <ul style="list-style-type: none"> ■ drowsiness, dizziness, or sleepiness occur ■ pain, cough, or nasal congestion gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. <p>Keep out of reach of children. Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p>	<p>Drug Facts (continued)</p> <p>Do not use</p> <ul style="list-style-type: none"> ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ liver disease ■ heart disease ■ diabetes ■ thyroid disease ■ kidney disease ■ trouble urinating due to an enlarged prostate gland ■ asthma, chronic bronchitis, or emphysema ■ trouble breathing or wheezing such as occurs with smoking, or <p>Ask a doctor or pharmacist before use if you are:</p> <ul style="list-style-type: none"> ■ taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers <p>When using this product, do not exceed recommended dosage.</p> <p>Stop use and ask a doctor if:</p> <ul style="list-style-type: none"> ■ drowsiness, dizziness, or sleepiness occur ■ pain, cough, or nasal congestion gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. <p>Keep out of reach of children. Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p>
---	---

COMBO PACK

<p>equate™</p> <p>DAYTIME NON-DROWSY</p> <p>Cold & Flu Multi-Symptom Relief</p> <p>Acetaminophen 325 mg - Pain reliever/fever reducer Dextromethorphan HBr 10 mg - Cough suppressant Phenylephrine HCl 5 mg - Nasal decongestant</p> <p>48 Total Softgels</p> <p>Relieves • Aches, Fever, Sore Throat • Nasal Congestion, Cough Alcohol-Free Antihistamine-Free</p> <p>32** Softgels (**Liquid-filled capsules)</p> <p>Actual size</p>	<p>NIGHTTIME</p> <p>Cold & Flu Multi-Symptom Relief</p> <p>Acetaminophen 325 mg - Pain reliever/fever reducer Dextromethorphan HBr 15 mg - Cough suppressant Doxylamine succinate 6.25 mg - Antihistamine</p> <p>16** Softgels (**Liquid-filled capsules)</p> <p>Actual size</p>
--	--

48 Total Softgels

Relieves
• Aches, Fever, Sore Throat
• Nasal Congestion, Cough
Alcohol-Free
Antihistamine-Free

32** Softgels
(**Liquid-filled capsules)

Actual size

Relieves
• Aches, Fever, Sore Throat
• Runny Nose, Sneezing
• Cough
Alcohol-Free

16** Softgels
(**Liquid-filled capsules)

Actual size

<p>Drug Facts (continued)</p> <p>Directions</p> <ul style="list-style-type: none"> ■ do not take more than directed (see Overdose warning) ■ do not take more than 4 doses in 24 hours ■ do not take more than 2 softgels with water every 6 hours ■ swallow whole, do not crush, chew, or dissolve ■ adults and children 12 years and over: take 2 softgels with water every 6 hours ■ children under 12 years: do not use <p>Other information</p> <ul style="list-style-type: none"> ■ store between 15-30°C (59-86°F) ■ avoid excessive heat <p>Inactive ingredients D&C yellow #10, FD&C blue #1, gelatin, glycerin, lactan, light mineral oil, mannitol, polyethylene glycol, polyethylene glycol, purified water, sorbitan, sorbitol, white ink</p> <p>Questions or comments? 1-888-287-1915</p>	<p>Drug Facts (continued)</p> <p>Directions</p> <ul style="list-style-type: none"> ■ do not take more than directed (see Overdose warning) ■ do not take more than 4 doses in 24 hours ■ do not take more than 2 softgels with water every 6 hours ■ swallow whole, do not crush, chew, or dissolve ■ adults and children 12 years and over: take 2 softgels with water every 6 hours ■ children under 12 years: do not use <p>Other information</p> <ul style="list-style-type: none"> ■ store between 15-30°C (59-86°F) ■ avoid excessive heat <p>Inactive ingredients D&C yellow #10, FD&C blue #1, gelatin, glycerin, lactan, light mineral oil, mannitol, polyethylene glycol, polyethylene glycol, purified water, sorbitan, sorbitol, white ink</p> <p>Questions or comments? 1-888-287-1915</p>
---	---

EQUATE Daytime Nighttime Multi-Symptom Cold & Flu Relief

48374

8113136224

6 5

PARENTS:
Child-resistant container required under the Poison Prevention Packaging Act (PPA). This is to help protect children from accidental ingestion. For more information on child-resistant containers, call 1-888-237-1915.

DISTRIBUTED BY: Walmart Inc.
Bentonville, AR 72716
PROJUL27 OF CHINA

This product is not manufactured or distributed by The Procter & Gamble Company, Norwalk, CT. DayQuil®, NyQuil®, and Lulicaps® are registered trademarks of The Procter and Gamble Company. FC008099

TAMPER EVIDENCE DO NOT USE IF CARTON IS OPENED OR IF FOLSTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.
 KEEP OUT OF REACH OF CHILDREN. READ COMPLETE WARNINGS AND PRODUCT INFORMATION.

Lot No.:
Exp. Date:

DAYTIME NIGHTTIME COLD FLU RELIEF

acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49035-192
---------------------	----------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49035-192-48	1 in 1 KIT; Type 0: Not a Combination Product	01/30/2020	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	16 BLISTER PACK	16
Part 2	32 BLISTER PACK	32

Part 1 of 2

NIGHTTIME COLD FLU RELIEF

acetaminophen, dextromethorphan hbr, doxylamine succinate capsule

Product Information

Route of Administration	ORAL
--------------------------------	------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
DOXYLAMINE SUCCINATE (UNII: V9BI9B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

Inactive Ingredients

Ingredient Name	Strength
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	

POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITAN (UNII: 6O92ICV9RU)	
SORBITOL (UNII: 506T60A25R)	
MANNITOL (UNII: 3OWL53L36A)	

Product Characteristics

Color	green	Score	no score
Shape	CAPSULE	Size	21mm
Flavor		Imprint Code	PC10
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		16 in 1 CARTON		
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	01/30/2020	

Part 2 of 2

DAYTIME COLD FLU RELIEF

acetaminophen, dextromethorphan hbr, phenylephrine hcl capsule

Product Information

Route of Administration	ORAL
--------------------------------	------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITAN (UNII: 6O921CV9RU)	
SORBITOL (UNII: 506T60A25R)	
MANNITOL (UNII: 3OWL53L36A)	
BUTYLATED HYDROXYANISOLE (UNII: REK4960K2U)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	

Product Characteristics

Color	orange	Score	no score
Shape	CAPSULE	Size	20mm
Flavor		Imprint Code	PC9
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		32 in 1 CARTON		
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	01/30/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	01/30/2020	

Labeler - EQUATE (Wal-Mart Stores, Inc.) (see also WAL-MART INC) (051957769)