


MIOSTAT - carbachol solution
Alcon Laboratories, Inc.

MIOSTAT*
(CARBACHOL INTRAOCULAR SOLUTION, USP) 0.01%

DESCRIPTION:

MIOSTAT* (carbachol intraocular solution, USP) 0.01% is a sterile balanced salt solution of carbachol for intraocular injection. The active ingredient is represented by the chemical structure:



Established Name: Carbachol

Chemical Name: Ethanaminium, 2-[(aminocarbonyl)oxy]-N,N,Ntrimethyl-, chloride.

Molecular Formula: C₆H₁₅ClN₂O₂

Molecular Weight: 182.65

Each mL of MIOSTAT* (carbachol intraocular solution, USP) 0.01% contains: Active: carbachol 0.01%.

Inactives: sodium chloride 0.64%, potassium chloride 0.075%, calcium chloride dehydrate 0.048%, magnesium chloride hexahydrate 0.03%, sodium acetate trihydrate 0.39%, sodium citrate dihydrate 0.17%, sodium hydroxide and/or hydrochloric acid (to adjust pH) and Water for Injection. pH range is 6.5-7.5.

CLINICAL PHARMACOLOGY:

Carbachol is a potent cholinergic (parasympathomimetic) agent which produces constriction of the iris and ciliary body resulting in reduction in intraocular pressure (IOP). The exact mechanism by which carbachol lowers IOP is not precisely known.

INDICATIONS AND USAGE:

Intraocular use for obtaining miosis during surgery. In addition, MIOSTAT* (carbachol intraocular solution, USP) 0.01% reduces the intensity of IOP elevation in the first 24 hours after cataract surgery.

CONTRAINDICATIONS:

Should not be used in those persons showing hypersensitivity to any of the components of this preparation.

WARNINGS:

For single-dose intraocular use only. Discard unused portion. Intraocular carbachol 0.01% should be used with caution in patients with acute cardiac failure, bronchial asthma, peptic ulcer, hyperthyroidism, G.I. spasm, urinary tract obstruction and Parkinson's disease. The vial stopper contains natural rubber (latex) which may cause severe allergic reactions.

PRECAUTIONS:

Use only if the container is undamaged.

Carcinogenesis:

Studies in animals to evaluate the carcinogenic potential have not been conducted.

Pregnancy:

There are no adequate and well-controlled studies in pregnant women. MIOSTAT* (carbachol intraocular solution, USP) 0.01% should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Nursing Mothers:

It is not known if this medication is excreted in breast milk. Exercise caution when administering to a nursing woman.

Pediatric Use:

Safety and efficacy in pediatric patients have not been established.

Geriatric Use:

No overall differences in safety or effectiveness have been observed between elderly and younger patients.

ADVERSE REACTIONS:

Ocular: Corneal clouding, persistent bullous keratopathy, retinal detachment and postoperative iritis following cataract extraction have been reported.

Systemic: Side effects such as flushing, sweating, epigastric distress, abdominal cramps, tightness in urinary bladder, and headache have been reported with topical or systemic application of carbachol.

The following additional reactions have been identified during post-approval use of MIOSTAT (carbachol intraocular solution, USP) 0.01% in clinical practice. Because they are reported voluntarily from a population of unknown size, estimates of frequency cannot be made. The reactions, which have been chosen for inclusion due to either their seriousness, frequency of reporting, possible causal connection to MIOSTAT, or a combination of these factors, include: corneal edema, drug effect

prolonged (miosis), eye inflammation, eye pain, intraocular pressure increased, ocular hyperemia, vision blurred, visual impairment, and vomiting.

DOSAGE AND ADMINISTRATION:

Aseptically remove the sterile vial from the blister package by peeling the backing paper and dropping the vial onto a sterile tray. Withdraw the contents into a dry sterile syringe, and replace the needle with an atraumatic cannula prior to intraocular instillation. No more than one-half milliliter should be gently instilled into the anterior chamber for the production of satisfactory miosis. It may be instilled before or after securing sutures. Miosis is usually maximal within two to five minutes after application.

HOW SUPPLIED:

In a 2.0 mL glass vial with a 1.5 mL fill, grey butyl stopper and aluminum seal packaged twelve to a carton.....NDC 0065-0023-15

STORAGE: Store at 15° - 30°C (59° - 86°F).

© 2002, 2003, 2015,2018 Novartis

Distributed by:
Alcon Laboratories, Inc.
Fort Worth, Texas 76134

***A trademark of Novartis**

Revised: June 2018
T2018-XX

PRINCIPAL DISPLAY PANEL

NDC 0065-0023-15

**MIOSTAT®
(CARBACHOL INTRAOCULAR SOLUTION, USP) 0.01%**

TWELVE 1.5 mL Sterile Glass Vials

Alcon®
Alcon Laboratories, Inc., Fort Worth, Texas 76134 USA

USUAL DOSAGE: 0.5 mL

STORAGE: Store at 15 - 30C (59 - 86F).


Rx Only

INGREDIENTS: A sterile balanced salt solution containing (per mL) **Active:** carbachol 0.01%.
Inactives: sodium chloride 0.64%, potassium chloride 0.075%, calcium chloride dihydrate 0.048%, magnesium chloride hexahydrate 0.03%, sodium acetate trihydrate 0.39%, sodium citrate dihydrate 0.17%, sodium hydroxide and/or hydrochloric acid (to adjust pH) and water for injection.

LOT:

EXP.:

390121-0513



MIOSTAT®
(carbachol intraocular solution, USP) 0.01%

Alcon®

© 2001-2005 Alcon, Inc.

320121-0305

LOT: EXP.:



LOT: EXP: **305190-1104**

NDC 0065-0023-15

MIOSTAT®
(CARBACHOL INTRAOCULAR SOLUTION, USP) 0.01% 1.5mL

Rx Only. Sterile Unless Package Open or Damaged Read enclosed insert. **INGREDIENTS: Active:** carbachol 0.01%. **Inactives:** sodium chloride 0.64%, potassium chloride 0.075%, calcium chloride dihydrate 0.048%, magnesium chloride hexahydrate 0.03%, sodium acetate trihydrate 0.39%, sodium citrate dihydrate 0.17%, sodium hydroxide and/or hydrochloric acid (to adjust pH) and water for injection. **USUAL DOSAGE:** 0.5mL **STORAGE:** Store at 15° - 30°C (59° - 86°F).

©2004 Alcon, Inc.

Alcon Labs., Inc. Fort Worth, TX. 76134

LOT:	2	EXP.:	2	305190-1104
NDC 0065-0023-15		MIOSTAT[®]		
(CARBACHOL INTRAOCULAR SOLUTION, USP) 0.01% 1.5 mL				
Rx Only. Sterile Unless Package Open or Damaged Read enclosed insert. INGREDIENTS:				
Active: carbachol 0.01%. Inactives: sodium chloride 0.64%, potassium chloride 0.075%, calcium chloride dihydrate 0.048%, magnesium chloride hexahydrate 0.03%, sodium acetate trihydrate 0.39%, sodium citrate dihydrate 0.17%, sodium hydroxide and/or hydrochloric acid (to adjust pH) and water for injection. USUAL DOSAGE: 0.5 mL STORAGE: Store at 15° - 30°C (59° - 86°F).				
©2004 Alcon, Inc. Alcon Labs., Inc. Fort Worth, TX. 76134				
				
3 0065002315 3				

MIOSTAT

carbachol solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0065-0023
Route of Administration	OPHTHALMIC		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CARBACHOL (UNII: 8 Y164V895Y) (Carbamoylcholine - UNII:54Z8M50D6Q)	CARBACHOL	0.1 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
POTASSIUM CHLORIDE (UNII: 660YQ98I10)	

CALCIUM CHLORIDE (UNII: M4I0D6VV5M)	
MAGNESIUM CHLORIDE (UNII: 02F3473H9O)	
SODIUM ACETATE (UNII: 4550K0SC9B)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	
WATER (UNII: 059QF0KO0R)	

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0065-0023-15	12 in 1 CARTON	04/15/1974	
1		1.5 mL in 1 VIAL, GLASS; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA016968	04/15/1974	

Labeler - Alcon Laboratories, Inc. (008018525)

Registrant - Alcon Laboratories, Inc. (008018525)

Establishment			
Name	Address	ID/FEI	Business Operations
Alcon Research Ltd		007672236	manufacture(0065-0023)