

**ALLERGY RELIEF- diphenhydramine hcl tablet
BETTER LIVING BRANDS LLC.**

160R_Albertsons_21130-596_Diphenhydramine HCl 25 mg Tablets

Drug Facts

Active ingredients (in each tablet)

Diphenhydramine HCl 25 mg

Purpose

Antihistamine

Uses

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

temporarily relieves these symptoms due to the common cold:

- runny nose
- sneezing

Warnings

Do not use

- to make a child sleepy
- with any other product containing diphenhydramine, even one used on skin

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.

When using this product

- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- take every 4 to 6 hours, or as directed by a doctor
- do not take more than 6 times in 24 hours

adults and children 12 years and over	1 to 2 tablets
children 6 to under 12 years	1 tablet
children under 6 years	do not use

Other information

- **each tablet contains:** calcium 25 mg
- protect from light
- store at 25°C (77° F) excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients carnauba wax, colloidal silicon dioxide, croscarmellose sodium, D&C red #27 aluminum lake, dibasic calcium phosphate dihydrate, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol (PEG) 400, polysorbate 80, titanium dioxide

Questions or comments?

Call **1-877-290-4008**



Compare to Benadryl®
Allergy Ultratab®
active ingredient
NDC 2130-596-35

Quality
Guaranteed

Allergy Relief

Diphenhydramine HCl 25 mg
Antihistamine Minitabs

- Easy to swallow
- Relief of: Sneezing, runny nose, itchy throat, itchy & watery eyes



Actual Size

365 MINITABS

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS GOOT, TORN, BROKEN OR MISSING.

Drug Facts

Active ingredient (in each tablet) Purpose
Diphenhydramine HCl 25 mg.....Antihistamine

Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
 - sneezing
 - itchy, watery eyes
 - itching of the nose or throat
- temporarily relieves these symptoms due to the common cold:
- runny nose
 - sneezing

Warnings

Do not use if you are taking sedatives or tranquilizers, or if you have been drinking alcohol within 14 days before use.

Do not use if you are taking a blood thinner, such as warfarin, or if you are taking a medicine for high blood pressure.

Do not use if you are taking a medicine for asthma, such as albuterol, or if you are taking a medicine for diabetes, such as insulin.

Do not use if you are taking a medicine for heart disease, such as digoxin, or if you are taking a medicine for thyroid disease, such as levothyroxine.

Do not use if you are taking a medicine for depression, such as amitriptyline, or if you are taking a medicine for anxiety, such as diazepam.

Do not use if you are taking a medicine for Parkinson's disease, such as levodopa, or if you are taking a medicine for multiple sclerosis, such as interferon beta.

Do not use if you are taking a medicine for HIV/AIDS, such as zidovudine, or if you are taking a medicine for hepatitis, such as interferon alpha.

Do not use if you are taking a medicine for malaria, such as chloroquine, or if you are taking a medicine for tuberculosis, such as isoniazid.

Do not use if you are taking a medicine for cancer, such as cyclophosphamide, or if you are taking a medicine for leukemia, such as cytarabine.

Do not use if you are taking a medicine for AIDS, such as zidovudine, or if you are taking a medicine for HIV, such as zidovudine.

Do not use if you are taking a medicine for hepatitis, such as interferon alpha, or if you are taking a medicine for multiple sclerosis, such as interferon beta.

Do not use if you are taking a medicine for Parkinson's disease, such as levodopa, or if you are taking a medicine for multiple sclerosis, such as interferon beta.

Do not use if you are taking a medicine for HIV/AIDS, such as zidovudine, or if you are taking a medicine for hepatitis, such as interferon alpha.

Do not use if you are taking a medicine for malaria, such as chloroquine, or if you are taking a medicine for tuberculosis, such as isoniazid.

Do not use if you are taking a medicine for cancer, such as cyclophosphamide, or if you are taking a medicine for leukemia, such as cytarabine.

Do not use if you are taking a medicine for AIDS, such as zidovudine, or if you are taking a medicine for HIV, such as zidovudine.

Do not use if you are taking a medicine for hepatitis, such as interferon alpha, or if you are taking a medicine for multiple sclerosis, such as interferon beta.

Do not use if you are taking a medicine for Parkinson's disease, such as levodopa, or if you are taking a medicine for multiple sclerosis, such as interferon beta.

Do not use if you are taking a medicine for HIV/AIDS, such as zidovudine, or if you are taking a medicine for hepatitis, such as interferon alpha.

Do not use if you are taking a medicine for malaria, such as chloroquine, or if you are taking a medicine for tuberculosis, such as isoniazid.

Do not use if you are taking a medicine for cancer, such as cyclophosphamide, or if you are taking a medicine for leukemia, such as cytarabine.

Do not use if you are taking a medicine for AIDS, such as zidovudine, or if you are taking a medicine for HIV, such as zidovudine.

Do not use if you are taking a medicine for hepatitis, such as interferon alpha, or if you are taking a medicine for multiple sclerosis, such as interferon beta.

Do not use if you are taking a medicine for Parkinson's disease, such as levodopa, or if you are taking a medicine for multiple sclerosis, such as interferon beta.

Do not use if you are taking a medicine for HIV/AIDS, such as zidovudine, or if you are taking a medicine for hepatitis, such as interferon alpha.

Do not use if you are taking a medicine for malaria, such as chloroquine, or if you are taking a medicine for tuberculosis, such as isoniazid.

Do not use if you are taking a medicine for cancer, such as cyclophosphamide, or if you are taking a medicine for leukemia, such as cytarabine.

Do not use if you are taking a medicine for AIDS, such as zidovudine, or if you are taking a medicine for HIV, such as zidovudine.

Do not use if you are taking a medicine for hepatitis, such as interferon alpha, or if you are taking a medicine for multiple sclerosis, such as interferon beta.

Do not use if you are taking a medicine for Parkinson's disease, such as levodopa, or if you are taking a medicine for multiple sclerosis, such as interferon beta.

Do not use if you are taking a medicine for HIV/AIDS, such as zidovudine, or if you are taking a medicine for hepatitis, such as interferon alpha.

Do not use if you are taking a medicine for malaria, such as chloroquine, or if you are taking a medicine for tuberculosis, such as isoniazid.

Do not use if you are taking a medicine for cancer, such as cyclophosphamide, or if you are taking a medicine for leukemia, such as cytarabine.

Do not use if you are taking a medicine for AIDS, such as zidovudine, or if you are taking a medicine for HIV, such as zidovudine.

Do not use if you are taking a medicine for hepatitis, such as interferon alpha, or if you are taking a medicine for multiple sclerosis, such as interferon beta.

Do not use if you are taking a medicine for Parkinson's disease, such as levodopa, or if you are taking a medicine for multiple sclerosis, such as interferon beta.

Do not use if you are taking a medicine for HIV/AIDS, such as zidovudine, or if you are taking a medicine for hepatitis, such as interferon alpha.

Do not use if you are taking a medicine for malaria, such as chloroquine, or if you are taking a medicine for tuberculosis, such as isoniazid.

Drug Facts (continued under label)


This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., McNeil Consumer Healthcare Division, owner of the registered trademark Benadryl®.

DISTRIBUTED BY: BETTER LIVING BRANDS LLC
P.O. BOX 99, PLEASANTON, CA 94566-0099
1-888-723-3929
180R 0223

LOVE IT
OR IT'S ON US

RD 23487

SS954



Scan here for more information



Lot No.:

Exp. Date:

PEE HERE

Varnish Omit Area

Drug Facts (continued)

When using this product:

- marked drowsiness may occur
- avoid alcoholic drinks, alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- take every 4 to 6 hours, or as directed by a doctor
- do not take more than 6 times in 24 hours

adults and children 12 years and over 1 to 2 tablets

children 6 to under 12 years 1 tablet

children under 6 years do not use

Other information

- each tablet contains: calcium 25 mg
- protected from light
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients

cantharal wax, colloidal silicon dioxide, croscarmellose sodium, D&C red #27 aluminum lake, dibasic calcium phosphate dihydrate, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol (PEG), 400, polysorbate 80, titanium dioxide

Questions or comments? 1-877-290-4008



Adhesive Area



Quality
Guaranteed

NDC 21130-596-20

Allergy Relief

Diphenhydramine HCl 25 mg
Antihistamine Minitabs

- Easy to swallow
- Relief of: Sneezing, runny nose, itchy throat, itchy & watery eyes

200 MINITABS

READ AND KEEP CAUTION FOR COMPLETE PRODUCT WARNINGS AND PRODUCT INFORMATION

TEMPERATURE SENSITIVE: DO NOT USE THIS PRODUCT IF THE PRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts

Active ingredient (in each tablet)

Diphenhydramine HCl 25 mg

Purpose

Antihistamine

Temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat
- watery eyes

Temporarily relieves these symptoms due to the common cold:

- runny nose
- sneezing

Warnings

Do not use

- to make a child sleepy
- with any other product containing diphenhydramine, even one used on skin

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Directions

- Take 1 to 6 hours, or as directed by doctor
- Do not take more than 6 times in 24 hours
- adults and children 12 years and over: 1 to 2 tablets
- children 6 to under 12 years: 1 tablet
- children under 6 years: do not use

Other information

- each tablet contains: calcium 25 mg
- protect from light
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Questions or comments?

1-877-296-4008

how2recycle.info
650926
PLASTIC BOTTLE

DISTRIBUTED BY: BETTER LIVING BRANDS, LLC
P.O. BOX 590 PLEASANTON, CA 94566-0059
PH: 887-723-3839
0000 0223
S554

Lot No:

Exp. Date:

Vamish Omit Area

Coating/Print Omit Area



READ AND KEEP CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION
TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts

Active Ingredient (In each tablet) Purpose
Diphenhydramine HCl 25 mg Antihistamine

Uses
temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
■ runny nose
■ sneezing
■ itchy, watery eyes
■ itching of the nose or throat
temporarily relieves these symptoms due to the common cold:
■ runny nose
■ sneezing

Warnings
Do not use
■ to make a child sleepy
■ with any other product containing diphenhydramine, even one used on skin

Ask a doctor before use if you have
■ a breathing problem such as emphysema or chronic bronchitis
■ glaucoma
■ trouble urinating due to enlarged prostate gland

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.

When using this product
■ marked drowsiness may occur
■ avoid alcoholic drinks
■ alcohol, sedatives, and tranquilizers may increase drowsiness
■ be careful when driving a motor vehicle or operating machinery
■ excitability may occur, especially in children

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions
■ take every 4 to 6 hours, or as directed by a doctor
■ do not take more than 6 times in 24 hours
adults and children 12 years and over 1 to 2 tablets
children 6 to under 12 years 1 tablet
children under 6 years do not use

Other information
■ each tablet contains: calcium 25 mg
■ protect from light
■ store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)
■ use by expiration date on package

Drug Facts (continued)

Inactive Ingredients
carnauba wax, colloidal silicon dioxide, croscarmellose sodium, D&C red #27 aluminum lake, dibasic calcium phosphate dihydrate, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol (PEG) 400, polysorbate 80, titanium dioxide

Questions or comments?
1-877-260-4808

*This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., McNeil Consumer Healthcare Division, owner of the registered trademark Benadryl®

DISTRIBUTED BY:
BETTER LIVING BRANDS LLC
P.O. BOX 99
PLEASANTON, CA 94566-0000
#1-888-723-3929
160R 0223

Coating Omit

Compare to Benadryl® Allergy UltraTab®
200 MINTABS
Allergies
Allergy Relief
Diphenhydramine HCl 25 mg
Antihistamine Minitabs
• Easy to swallow
• Relief of: Sneezing, runny nose, itchy throat, itchy & watery eyes



Coating/Print Omit Area

Coating Omit



Adhesive Area

Drug Facts (continued)

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- take every 4 to 6 hours, or as directed by a doctor
- do not take more than 6 times in 24 hours
- adults and children 12 years and over 1 to 2 tablets
- children 6 to under 12 years 1 tablet
- children under 6 years do not use

Other information

- each tablet contains calcium 25 mg
- protect from light
- store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Questions or comments? 1-877-230-4008



NDC 21130-596-01

Allergy Relief

Diphenhydramine HCl 25 mg
Antihistamine Minitabs

- Easy to swallow
- Relief of: Sneezing, runny nose, itchy throat, itchy & watery eyes

100 MINITABS

READ AND KEEP CARTRON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts

Active ingredient (in each tablet) Purpose
Diphenhydramine HCl 25 mg.....Antihistamine

Uses

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

temporarily relieves these symptoms due to the common cold:

- runny nose
- sneezing

Warnings

- Do not use
- to make a child sleepy
- with any other product containing diphenhydramine, even one used on skin

how2recycle.info

LOVE IT OR IT'S ON US +

DISTRIBUTED BY:
BETTER LIVING BRANDS LLC
P.O. BOX 99
PLEASANTON, CA 94566-0099
1-888-723-3929

160R 0223

RD 23059
55854

Drug Facts (continued under label)

Varnish Omit Area

Lot#: _____ Exp. Date: _____



READ AND KEEP CARTRON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts

Active ingredient (in each tablet) Purpose
Diphenhydramine HCl 25 mg.....Antihistamine

Uses

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

temporarily relieves these symptoms due to the common cold:

- runny nose
- sneezing

Warnings

- Do not use
- to make a child sleepy
- with any other product containing diphenhydramine, even one used on skin

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma

Drug Facts (continued)

- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.

When using this product

- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- take every 4 to 6 hours, or as directed by a doctor
- do not take more than 6 times in 24 hours
- adults and children 12 years and over 1 to 2 tablets
- children 6 to under 12 years 1 tablet
- children under 6 years do not use

Other information

- each tablet contains calcium 25 mg
- protect from light

Drug Facts (continued)

- store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients carnauba wax, colloidal silicon dioxide, croscarmellose sodium, D&G red #27 aluminum lake, calcium phosphate dihydrate, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol (PEG) 400, polyisobutyl 80, titanium dioxide

Questions or comments? 1-877-230-4008

This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., McNeil Consumer Healthcare Division, owner of the registered trademark Benadryl.

DISTRIBUTED BY:
BETTER LIVING BRANDS LLC
P.O. BOX 99
PLEASANTON, CA 94566-0099
1-888-723-3929

160R 0223

Scan here for more information

Compare to Signature Care UltraTab[®] now registered

NDC 21130-596-01

Allergy Relief

Diphenhydramine HCl 25 mg
Antihistamine Minitabs

100 MINITABS

- Easy to swallow
- Relief of: Sneezing, runny nose, itchy throat, itchy & watery eyes

Varnish/Print Omit Area

E3144R



ALLERGY RELIEF

diphenhydramine hcl tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:21130-596
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	25 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
D&C RED NO. 27 ALUMINUM LAKE (UNII: ZK64F7XSTX)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	

Product Characteristics

Color	pink	Score	no score
Shape	CAPSULE	Size	11mm
Flavor		Imprint Code	T;061
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21130-596-20	1 in 1 CARTON	05/31/2023	
1		200 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:21130-596-01	1 in 1 CARTON	05/31/2023	
2		100 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:21130-596-35	365 in 1 BOTTLE; Type 0: Not a Combination Product	07/27/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/31/2023	

Labeler - BETTER LIVING BRANDS LLC. (009137209)

Registrant - TIME CAP LABORATORIES, INC. (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
TIME CAP LABORATORIES, INC.		037052099	manufacture(21130-596)

Revised: 10/2023

BETTER LIVING BRANDS LLC.