

**NAPROXEN SODIUM- naproxen sodium tablet  
BETTER LIVING BRANDS LLC.**

-----  
**609R\_21130-690\_Albertsons Naproxen Sodium 220 mg Tablets**

***Drug Facts***

***Active ingredient (in each tablet)***

Naproxen sodium 220 mg

(naproxen 200 mg) (NSAID)\*

\*nonsteroidal anti-inflammatory drug

***Purposes***

Pain reliever/fever reducer

***Uses***

- temporarily relieves minor aches and pains due to:
  1. minor pain of arthritis
  2. muscular aches
  3. backache
  4. menstrual cramps
  5. headache
  6. toothache
  7. the common cold
- temporarily reduces fever

***Warnings***

**Allergy alert:** Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems

- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

### **Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

### **Ask a doctor before use if**

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

### **Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

### **When using this product**

- take with food or milk if stomach upset occurs

### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  1. feel faint
  2. vomit blood
  3. have bloody or black stools
  4. have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  1. chest pain
  2. trouble breathing
  3. weakness in one part or side of body
  4. slurred speech
  5. leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days

- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

### ***Directions***

- **do not take more than directed**
- **the smallest effective dose should be used**
- drink a full glass of water with each dose

---

Adults and children 12 years and older	<ul style="list-style-type: none"> <li>• take 1 tablet every 8 to 12 hours while symptoms last</li> <li>• for the first dose you may take 2 tablets within the first hour</li> <li>• do not exceed 2 tablets in any 8- to 12-hour period</li> <li>• do not exceed 3 tablets in a 24-hour period</li> </ul>
Children under 12 years	<ul style="list-style-type: none"> <li>• ask a doctor</li> </ul>

---

### ***Other information***

- **each tablet contains:** sodium 20 mg
- store at 20-25°C (68-77°F).
- avoid high humidity and excessive heat above 40°C (104°F).

### ***Inactive ingredients***

colloidal silicon dioxide, croscarmellose sodium, FD&C blue #2 Lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, titanium dioxide

**Questions or comments?1-877-290-4008**



NDC 21130-690-05

# Naproxen Sodium TABLETS USP, 220 mg Pain Reliever/Fever Reducer (NSAID)

Strength to last 12 hours

50 TABLETS

**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION**  
**TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING**

### Drug Facts

#### Active ingredient (in each tablet)

Naproxen sodium 220 mg.....Pain reliever/fever reducer (naproxen 220 mg) (NSAID)\*

\*Nonsteroidal anti-inflammatory drug.

**Uses** ■ temporarily relieves minor aches and pains due to:  
● minor pain of arthritis ● muscular aches ● backache  
● menstrual cramps ● headache ● toothache ● the common cold ■ temporarily reduces fever

#### Warnings

**Allergy alert:** Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

■ hives ■ facial swelling ■ asthma (wheezing)

■ shock ■ skin redness ■ rash ■ blisters

If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if

you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug

■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more

alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal.

The risk is higher if you use more than directed or for longer than directed. If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.**

**Directions** ■ do not take more than directed ■ the smallest effective dose should be used ■ drink a full glass of water with each

dose. Adults and children 12 years and older: ● take 1 tablet every 8 to 12 hours while symptoms last ● for the first dose you may take 2

tablets within the first hour ● do not exceed 2 tablets in any 8- to 12-hour period ● do not exceed 3 tablets in a 24-hour period.

Children under 12 years: ● ask a doctor.

**Questions or comments? 1-877-298-4008**

### LOVE IT

OR IT'S ON US\*

DISTRIBUTED BY:

BETTER LIVING BRANDS LLC

P.O. BOX 99

PLEASANTON, CA 94566-0099

†1-888-723-3829

PO 23053 S5934

Made in India \*08P 0223

how2recycle.info



Lot No.:

Exp Date:

Varnish Omit Area



# NAPROXEN SODIUM

naproxen sodium tablet

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:21130-690
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>NAPROXEN SODIUM</b> (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)	NAPROXEN SODIUM	220 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>FD&amp;C BLUE NO. 2</b> (UNII: L06K8R7DQK)	
<b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

<b>CROSCARMELLOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	

### Product Characteristics

<b>Color</b>	blue	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	141
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21130-690-05	1 in 1 CARTON	06/13/2023	
1		50 in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090545	06/13/2023	

**Labeler** - BETTER LIVING BRANDS LLC. (009137209)

**Registrant** - TIME CAP LABORATORIES, INC. (037052099)

### Establishment

Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LIMITED		925822975	manufacture(21130-690)

Revised: 6/2023

BETTER LIVING BRANDS LLC.