

ASPIRIN- aspirin tablet
Advance Pharmaceutical Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

ASPIRIN 325mg

Active Ingredient

(in each tablet)

Aspirin 325 mg (NSAID) *

*nonsteroidal anti-inflammatory drug

Purpose

Pain Reliever

Uses

temporarily relieves

- headache
- menstrual pain
- minor pain or arthritis
- muscle pain
- toothache
- pain and fever and cold

Warnings

Reye's Syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early signs of Reye's syndrome, a rare but serious illness.

Allergy Alert: Aspirin may cause a severe allergic reaction which may include

- hives
- asthma (wheezing)
- shock
- facial swelling

Stomach Bleeding Warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood-thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (ibuprofen, naproxen, or others)
- takes more or for a longer time than directed
- have 3 or more alcoholic drinks every day while using this product.

Do not use

- if you have ever had an allergic reaction to any other pain reliever / fever reducer

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout or arthritis
- taking any other drug
- under a doctor's care for any serious condition

Stop use and ask a doctor if

- you experiences any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear
- ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

DRINK A FULL GLASS OF WATER WITH EACH DOSE

- **adults and children 12 years and older** : 1-2 tablets every 4 hours while symptoms last, but not more than 12 tablets in 24 hours.
- **Children under 12 years**: ask a doctor

Other Information

- avoid excessive heat or humidity
- store at 15-30 °C (59-86 °F)

Inactive Ingredients

corn starch, PVP

Questions or Comments

call 631-981-4600, 8.30 am-4.30 pm ET, Monday - Friday

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

For temporary relief of minor aches and pains

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL



Advance
Pharmaceutical Inc.

PAIN RELIEVER/FEVER REDUCER
100 TABLETS

**ASPIRIN TABLETS
(NSAID)**

NDC 17714-010-01
*Compare to active
ingredient in Bayer®
Aspirin Regimen 325 mg



Drug Facts

Active ingredient (in each tablet) Purpose

Aspirin 325 mg (NSAID)*Pain reliever/Fever reducer
*nonsteroidal anti-inflammatory drug

Uses temporarily relieves ■ headache ■ menstrual pain
■ minor pain of arthritis ■ muscle pain ■ toothache
■ pain and fever of colds

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include ■ hives ■ facial swelling
■ asthma (wheezing) ■ shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- take more or for a longer time than directed
- have 3 or more alcoholic drinks every day while using this product

Drug Facts continued on back of label.

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

*Advance Pharmaceutical Inc. is not affiliated with the owner of the trademark BAYER® Aspirin Regimen 325 mg.

Manufactured by: Advance Pharmaceutical Inc., Holtsville, NY 11742, USA



LA0413



Lot No.:

Exp. Date:

Drug Facts (continued)

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic ■ you have asthma

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout or arthritis
- under a doctor's care for any serious condition
- taking any other drug

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear
- ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions DRINK A FULL GLASS OF WATER WITH EACH DOSE ■ **adults and children 12 years and over:** 1-2 tablets every 4 hours while symptoms last, not more than 12 tablets in 24 hours
■ **children under 12 years:** ask a doctor

Other information ■ avoid excessive heat or humidity
■ store at room temperature 15°-30°C (59°-86°F)

Inactive ingredients corn starch, PVP

Questions or comments?

call 631-981-4600, 8:30 am-4:30 pm ET, Monday - Friday



**Advance
Pharmaceutical Inc.**

**PAIN RELIEVER/FEVER REDUCER
300 TABLETS**

325 mg (5gr)

**ASPIRIN TABLETS
(NSAID)**

NDC 17714-010-03
*Compare to active
ingredient in Bayer®
Aspirin Regimen 325 mg



Drug Facts

Active ingredient (in each tablet)	Purpose
Aspirin 325 mg (NSAID)*	Pain reliever/Fever reducer

*nonsteroidal anti-inflammatory drug

Uses temporarily relieves

- headache
- menstrual pain
- minor pain of arthritis
- muscle pain
- toothache
- pain and fever of colds

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- take more or for a longer time than directed
- have 3 or more alcoholic drinks every day while using this product

Drug Facts continued on back of label.

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

*Advance Pharmaceutical Inc. is not affiliated with the owner of the trademark BAYER® Aspirin Regimen 325 mg.

Manufactured by: Advance Pharmaceutical Inc., Holtsville, NY 11742, USA



LA0413



Lot No.:

Exp. Date:

Drug Facts (continued)

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic ■ you have asthma

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout or arthritis
- under a doctor's care for any serious condition
- taking any other drug

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear
- ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions DRINK A FULL GLASS OF WATER WITH EACH DOSE ■ **adults and children 12 years and over:** 1- 2 tablets every 4 hours while symptoms last, not more than 12 tablets in 24 hours
■ **children under 12 years:** ask a doctor

Other information ■ avoid excessive heat or humidity
■ store at room temperature 15°-30°C (59°-86°F)

Inactive ingredients corn starch, PVP

Questions or comments?

call 631-981-4600, 8:30 am-4:30 pm ET, Monday - Friday

NDC: 17714-010-01 – 100 Tablets

NDC: 17714-010-03 - 300 Tablets

ASPIRIN

aspirin tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:17714-010
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
POVIDONE (UNII: FZ989GH94E)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	AP;010
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:17714-010-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	12/01/2015	
2	NDC:17714-010-03	300 in 1 BOTTLE; Type 0: Not a Combination Product	12/01/2015	
3	NDC:17714-010-10	1000 in 1 BOTTLE; Type 0: Not a Combination Product	01/09/2002	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part343	01/09/2002	

Labeler - Advance Pharmaceutical Inc. (078301063)**Registrant** - Advance Pharmaceutical Inc. (078301063)**Establishment**

Name	Address	ID/FEI	Business Operations
Advance Pharmaceutical Inc.		078301063	manufacture(17714-010)