

**LOW DOSE ASPIRIN 81MG- aspirin tablet  
BETTER LIVING BRANDS LLC.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**481R\_Albertson\_21130-481\_Aspirin 81mg EC Yellow**

**Drug Facts**

**Active ingredient (in each tablet)**

Aspirin 81 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever

**Uses**

- for the temporary relief of minor aches and pains or as recommended by your doctor. **Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.**
- ask your doctor about other uses for enteric coated 81 mg Aspirin.

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer

- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

### **Ask a doctor or pharmacist before use if you are** taking a prescription drug for

- gout
- diabetes
- arthritis

### **Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  1. feel faint
  2. vomit blood
  3. have bloody or black stools
  4. have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or a loss of hearing occurs

**If pregnant or breast-feeding**, ask a health professional before use . **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

### **Directions**

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

### **Other information**

- store at room temperature. Avoid excessive heat above 40°C (104°F)

**Inactive ingredients** anhydrous lactose, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10 aluminum lake, iron oxide ochre, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate

Questions or comments? 1-877-290-4008

Adhesive Area

**Drug Facts (continued)**

Take more or for a longer time than directed

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer. If you have ever had an allergic reaction to this product or any of its ingredients.

**Ask a doctor before use if:**

- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for:

- gout
- diabetes
- arthritis

**Stop using and ask a doctor if:**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - faint or vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- symptoms worsen or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or a loss of hearing occurs

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

- Drink a full glass of water with each dose
- Adults and children 12 years and over: Take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- Children under 12 years: consult a doctor

**Other information**

■ Store at room temperature. Avoid excessive heat above 40°C (104°F).

**Inactive ingredients** include: hydroxypropyl methylcellulose, croscarmellose sodium, DCP yellow #10 aluminum lake, iron oxide ochre, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polyethylene glycol 400, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate

Questions or comments? 1-877-290-4008




Compare to Bayer® Low Dose Aspirin active ingredient\* NDC 2130-481-90

Quality Guaranteed

Low Dose Aspirin 81 mg Pain reliever (NSAID)

- Enteric coated
- Aspirin regimen\*\*



\*\*Talk to your doctor or other health care professional for your heart. TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CRACKED, BROKEN OR MISSING

**Drug Facts**

**Active ingredient (in each tablet)** Aspirin 81 mg (NSAID) - nonsteroidal anti-inflammatory drug

**Purpose** Pain reliever

**Uses** for the temporary relief of minor aches and pains or as recommended by your doctor. Because of the delayed action, this product will not provide fast relief of headaches or other symptoms a need immediate relief. ask your doctor about other uses for enteric coated 81 mg Aspirin.

**Warnings** **Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. **Allergy alert:** Aspirin may cause a severe allergic reaction, which may include: hives, facial swelling, asthma (wheezing), shock **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroidal drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product

**Drug Facts (continued under label)**

This product is not meant to be used if distributed by Bayer AG. [how2recycle.info](http://how2recycle.info)

DISTRIBUTED BY BETTER LIVING BRANDS, LLC  
 P.O. BOX 96, PLEASANTON, CA 94566-0096  
 1-888-729-3829

Scan for info

Lot No.: Exp. Date:

48FR 0823 3 21130176029 0

Varnish Omit Area



NDC 21190-481-20

Low Dose  
**Aspirin 81 mg**  
Pain reliever (NSAID)

- Enteric coated
- Aspirin regimen\*\*

200 TABLETS

**DO NOT TAKE ANY OTHER ASPIRIN PRODUCTS.** **DO NOT TAKE ANY OTHER PAIN RELIEVERS.** **DO NOT TAKE ANY OTHER NSAID PRODUCTS.** **DO NOT TAKE ANY OTHER BLOOD THINNING DRUGS.** **DO NOT TAKE ANY OTHER DRUGS.** **DO NOT TAKE ANY OTHER MEDICATIONS.**

**Drug Facts**

**Purpose**  
Aspirin 81 mg (NSAID) is a pain reliever.

**Uses** ■ For the temporary relief of minor aches and pains or as a blood thinner when used as directed. It is used for the relief of inflammation and to prevent blood clots. **DO NOT TAKE ANY OTHER ASPIRIN PRODUCTS.** **DO NOT TAKE ANY OTHER PAIN RELIEVERS.** **DO NOT TAKE ANY OTHER NSAID PRODUCTS.** **DO NOT TAKE ANY OTHER BLOOD THINNING DRUGS.** **DO NOT TAKE ANY OTHER DRUGS.** **DO NOT TAKE ANY OTHER MEDICATIONS.**

**Warnings** ■ Ask your doctor about other uses for enteric-coated 81 mg Aspirin. **DO NOT TAKE ANY OTHER ASPIRIN PRODUCTS.** **DO NOT TAKE ANY OTHER PAIN RELIEVERS.** **DO NOT TAKE ANY OTHER NSAID PRODUCTS.** **DO NOT TAKE ANY OTHER BLOOD THINNING DRUGS.** **DO NOT TAKE ANY OTHER DRUGS.** **DO NOT TAKE ANY OTHER MEDICATIONS.**

**Directions** **Directions:** See label for full directions. Take 1 tablet (81 mg) with water 2 to 3 times daily as needed every 4 to 6 hours. Do not take more than 4 tablets in 24 hours. **DO NOT TAKE ANY OTHER ASPIRIN PRODUCTS.** **DO NOT TAKE ANY OTHER PAIN RELIEVERS.** **DO NOT TAKE ANY OTHER NSAID PRODUCTS.** **DO NOT TAKE ANY OTHER BLOOD THINNING DRUGS.** **DO NOT TAKE ANY OTHER DRUGS.** **DO NOT TAKE ANY OTHER MEDICATIONS.**

**Other information** ■ Aspirin may cause severe allergic reactions in some people. ■ Allergic reactions include: hives, rash, itching, swelling, dizziness, trouble breathing. ■ If you experience these symptoms, stop taking this product and seek medical attention immediately. **DO NOT TAKE ANY OTHER ASPIRIN PRODUCTS.** **DO NOT TAKE ANY OTHER PAIN RELIEVERS.** **DO NOT TAKE ANY OTHER NSAID PRODUCTS.** **DO NOT TAKE ANY OTHER BLOOD THINNING DRUGS.** **DO NOT TAKE ANY OTHER DRUGS.** **DO NOT TAKE ANY OTHER MEDICATIONS.**

**Contains aspirin** ■ Aspirin may cause severe allergic reactions in some people. ■ Allergic reactions include: hives, rash, itching, swelling, dizziness, trouble breathing. ■ If you experience these symptoms, stop taking this product and seek medical attention immediately. **DO NOT TAKE ANY OTHER ASPIRIN PRODUCTS.** **DO NOT TAKE ANY OTHER PAIN RELIEVERS.** **DO NOT TAKE ANY OTHER NSAID PRODUCTS.** **DO NOT TAKE ANY OTHER BLOOD THINNING DRUGS.** **DO NOT TAKE ANY OTHER DRUGS.** **DO NOT TAKE ANY OTHER MEDICATIONS.**

**Questions or comments?** Visit [www.lovetheplanet.com](http://www.lovetheplanet.com)



**How2Recycle.info**  
PLASTIC BOTTLE

**LOT'S ON US!**  
LOVE IT  
GALIT'S ON US!

**Lot No.:**  
**Exp. Date:**

**Signature Care Quality Guaranteed**

Coating/Print Omit Area

Let No. Exp Date:



RD 23199 52954

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

### Drug Facts

Active ingredient (in each tablet) Purpose  
Aspirin 81 mg (NSAID) Pain reliever  
hydroxyacetic acid/anticoagulant drug

**Uses**  
For the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will provide fast relief of headaches or other symptoms needing immediate relief. Ask your doctor about other aspirin coated 81 mg Aspirin.

**Warnings**  
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, it changes in behavior with nausea and vomiting occur, contact a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include: • Hives • Facial swelling • Asthma (wheezing) • Shock  
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: • are age 65 or older • have had stomach ulcers or bleeding problems • take other drugs (particularly) or stomach drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin)

### Drug Facts (continued)

• Pregnant, pregnant, or others • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed

Do not use if you are allergic to aspirin or any other pain reliever or reducer • If you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if • stomach bleeding warning applies to you • you have a history of stomach problems, such as heartburn • you have high blood pressure, heart disease, liver disease, or kidney disease • you are taking a diuretic • you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for • gout • diabetes • arthritis

Stop use and ask a doctor if • an allergic reaction occurs. Seek medical help right away • you experience any of the following signs of stomach bleeding: • fast heart • vomit blood • have bloody or black stools • have stomach pain that does not get better • pain gets worse or lasts more than 3 days • lower gas worse or lasts more than 3 days • nausea or swelling is present • new symptoms occur • ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

### Drug Facts (continued)

**Directions**  
• Drink a full glass of water with each dose • Adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor • Children under 12 years: consult a doctor

**Other information** • Store at room temperature. Avoid excessive heat above 40°C (104°F)

**Inactive ingredients** antyoxen lactate, carmellose, croscarmellose sodium, D&G yellow #10 aluminum lake, iron oxide ochre, methylacrylate acid and ethyl acrylate copolymer, microcrystalline cellulose, polyvinyl alcohol, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate

Questions or comments? 1-877-266-4068

This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Bayer.

DISTRIBUTED BY:  
BETTER LIVING BRANDS LLC  
P.O. BOX 50  
PLEASANTON, CA 94566-0050  
1-888-723-3320  
481R 0623



Coating Omit

Company to Bayer Low Dose Aspirin active ingredient

NDC 23199-81-30



200 TABLETS

# Aspirin 81 mg

Pain reliever (NSAID)

- Enteric coated
- Aspirin regimen\*\*



Low Dose Aspirin 81 mg Pain reliever (NSAID) 200 TABLETS



NDC 23199-81-30

Coating Omit

Coating/Print Omit Area



Quality  
Guaranteed

NDC 21130-481-12

# Low Dose Aspirin 81 mg Pain reliever (NSAID)

- Enteric coated
- Aspirin regimen\*\*

120 TABLETS

\*\*Talk to your doctor or other healthcare provider before using this product, your health.

**READ AND KEEP CHILDREN FROM EATING COMPLETE PRODUCT. WARNING SAND**

**TAMPER EVIDENT DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL**

**OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING**

### Drug Facts

#### Active ingredient (in each tablet)

Aspirin (in mg (NSAID)) ..... Pain reliever

Aspirin (in mg (NSAID)) ..... Pain reliever

Use as directed. Do not use for more than 10 days unless directed by a doctor.

Do not use if you are allergic to aspirin or any of the ingredients. This

product will not provide fast relief of headaches or other symptoms

unless you take it as directed. Ask your doctor about other uses for

enteric coated 81 mg Aspirin.

#### Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering

from chicken pox or flu-like symptoms should not use this product. When

using this product, if changes in behavior with nausea and vomiting

occur, consult a doctor because these symptoms could be an early sign

of Reye's syndrome, a rare but serious illness. **Allergy alert:** Aspirin may

cause a severe allergic reaction which may include: ■ hives ■ facial

swelling ■ asthma (wheezing) ■ shock. **Stomach bleeding warning:**

This product contains an NSAID, which may cause severe stomach

bleeding, which may be fatal. This risk increases with longer use. **Use**

**cautiously if you have:** ■ a history of ulcers or bleeding problems ■ take a blood thinner

(anticoagulant) or steroid drug ■ take other drugs containing

aspirin or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or

others) ■ have 3 or more alcoholic drinks every day while using this

product ■ take more or for a longer time than directed. **If pregnant or**

**breast-feeding,** ask a health professional before use. **Keep out of reach**

**of children.**

**Directions:** ■ drink a full glass of water with each dose ■ adults and

children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed

42 tablets in 24 hours unless directed by a doctor ■ children under 12

years: consult a doctor

**Questions or comments? 1-877-290-4800**

481 R 0823

©2014 Varnish Omit Area, LLC

P.O. BOX 481, PLEASANTON, CA 94566-0481

1-888-726-8829

Lot No.:

Exp. Date:

481 R 0823

©2014 VARNISH OMIT AREA, LLC

P.O. BOX 481, PLEASANTON, CA 94566-0481

1-888-726-8829

Lot No.:

Exp. Date:

481 R 0823

©2014 VARNISH OMIT AREA, LLC

P.O. BOX 481, PLEASANTON, CA 94566-0481

1-888-726-8829

Lot No.:

Exp. Date:

481 R 0823

©2014 VARNISH OMIT AREA, LLC

P.O. BOX 481, PLEASANTON, CA 94566-0481

1-888-726-8829

Lot No.:

Exp. Date:

Varnish Omit Area

Side of Package

Coating/Print Omit Area



READ AND KEEP CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION  
TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

**Drug Facts**

**Active ingredient (in each tablet)**  
Aspirin 81 mg (NSAID)\*  
**Purpose**  
Pain reliever  
\*nonsteroidal anti-inflammatory drug

**Uses** ■ for the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide best relief of headaches or other symptoms needing immediate relief. ■ ask your doctor about other uses for enteric coated 81 mg Aspirin.

**Warnings**  
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock  
Stomach bleeding warning: This product contains

**Drug Facts (continued)**

an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or stomach drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Do not use ■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for ■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if ■ an allergic reaction occurs. Seek medical help right away. ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ pain gets worse or lasts more than 10 days ■ never gets worse or lasts more

**Drug Facts (continued)**

than 5 days ■ redness or swelling is present ■ new symptoms occur ■ ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center. (1-800-222-1222) right away.

**Directions** ■ drink a full glass of water with each dose ■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor ■ children under 12 years: consult a doctor

**Other information** ■ Store at room temperature. Avoid excessive heat above 40°C (104°F)

**Inactive ingredients** anhydrous lactose, croscarmellose sodium, DSC yellow #10 aluminum lake, iron oxide ochre, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polyacrylate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate

**Questions or comments?** 1-877-298-4008

Coating Omit

Compare to Bayer Low Dose Aspirin active ingredient  
NDC 2110-4812  
Actual Size  
120 TABLETS



Quality Guaranteed

Low Dose Aspirin 81 mg  
Pain reliever (NSAID)  
• Enteric coated  
• Aspirin regimen\*\*

Coating/Print Omit Area

Coating Omit

\*This product is not manufactured or distributed by Bayer AG, owner of the registered trademark "Aspirin".



100% Recycled Content







## LOW DOSE ASPIRIN 81MG

aspirin tablet

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:21130-481
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>CROSCARMELOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	

<b>SODIUM HYDROXIDE</b> (UNII: 55X04QC32I)
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)
<b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)
<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O)
<b>ANHYDROUS LACTOSE</b> (UNII: 3SY5LH9PMK)
<b>METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER</b> (UNII: NX76LV5T8J)
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)
<b>TALC</b> (UNII: 7SEV7J4R1U)
<b>TRIETHYL CITRATE</b> (UNII: 8Z96QXD6UM)
<b>FERRIC OXIDE YELLOW</b> (UNII: EX438O2MRT)

### Product Characteristics

<b>Color</b>	yellow	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	7mm
<b>Flavor</b>		<b>Imprint Code</b>	HEART
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21130-481-32	1 in 1 CARTON	07/14/2023	
1		32 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:21130-481-12	1 in 1 CARTON	07/14/2023	
2		120 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:21130-481-20	1 in 1 CARTON	07/14/2023	
3		200 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:21130-481-50	500 in 1 BOTTLE; Type 0: Not a Combination Product	07/14/2023	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	07/14/2023	

**Labeler** - BETTER LIVING BRANDS LLC. (009137209)

**Registrant** - TIME CAP LABORATORIES, INC. (037052099)

## Establishment

Name	Address	ID/FEI	Business Operations
TIME CAP LABORATORIES, INC.		037052099	manufacture(21130-481)

Revised: 8/2023

BETTER LIVING BRANDS LLC.