

**ASPIRIN 81MG- aspirin tablet, delayed release
BETTER LIVING BRANDS LLC.**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

482R_Albertson_21130-428_Aspirin Delayed -Release Tablets USP, 81mg EC

Drug Facts

Active ingredient (in each tablet)

Aspirin 81 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

- temporarily relieves minor aches and pain
- ask your doctor about other uses for enteric coated 81 mg Aspirin.

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: • hives • facial swelling • asthma (wheezing) • shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problem
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

- you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for gout, diabetes or arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - ●feel faint ● vomit blood ● have bloody or black stools ● have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- ringing in the ears or a loss of hearing occurs
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- drink a full glass of water with each dose
- adults and children 12 years of age and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years of age: do not use unless directed by a doctor

Other information

- store at 20-25°C (68-77°F)
- close cap tightly after use

Inactive ingredients anhydrous lactose, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, iron oxide yellow, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, triethyl citrate

Questions or comments?1-877-290-4008

Side of Package

Coating/Print Omit Area

Lot No.:
Exp. Date:

RD 23159
692
52964

4828 0828
1-800-898-7222
P.O. BOX 990
LETTERMAN BRANCH, LLC
600 99669 000
P.O. BOX 990
LETTERMAN BRANCH, LLC
more information
go to www.aspirin.com

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts

Active ingredient (in each tablet) **Purpose**

Aspirin 81 mg (NSAID)* Pain reliever
*nonsteroidal anti-inflammatory drug

Uses ■ temporarily relieves minor aches and pain
■ ask your doctor about other uses for enteric coated 81 mg Aspirin.

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems

Drug Facts (continued)

■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product
■ take more or for a longer time than directed

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for gout, diabetes or arthritis

Stop use and ask a doctor if ■ an allergic reaction occurs. Seek medical help right away. ■ you experience any of the following signs of stomach bleeding: ● feel faint ● vomit blood ● have bloody or black stools ● have stomach pain that does not get better

■ pain gets worse or lasts more than 10 days
■ ringing in the ears or a loss of hearing occurs
■ redness or swelling is present
■ new symptoms occur
These could be signs of a serious condition

Drug Facts (continued)

if pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions ■ drink a full glass of water with each dose ■ adults and children 12 years of age and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor ■ children under 12 years of age: do not use unless directed by a doctor

Other information ■ store at 20-25°C (68-77°F) ■ close cap tightly after use

Inactive ingredients anhydrous lactose, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, iron oxide yellow, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, triethyl citrate

Questions or comments? 1-877-290-4008

*This product is not manufactured or distributed by Foundation Consumer Healthcare LLC, owner of the registered trademark St. Joseph® Low Dose 81 mg Aspirin.

Compare to St. Joseph® Low Dose Safety Coated 81 mg Aspirin active ingredient!

120 TABLETS

Actual Size

Low Dose Aspirin 81 mg

Pain reliever (NSAID)

• Enteric coated
• Aspirin regimen**



Coating/Print Omit Area

**Talk to your doctor or other healthcare provider before using this product for your heart.



Coating Omit

Coating/Print
Omit Area



READ AND KEEP CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION
TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.

Drug Facts

Active ingredient (in each tablet) Aspirin 81 mg (NSAID)
Purpose Pain reliever, nonsteroidal anti-inflammatory drug

Uses temporarily relieves minor aches and pain
Use as your doctor orders other uses for enteric coated 81 mg Aspirin

Warnings
People's symptoms: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Always read: Aspirin may cause a severe allergic reaction which may include:
• hives
• facial swelling
• asthma (wheezing)
• shock
Stomach bleeding warnings: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
• are age 60 or older
• have had stomach ulcers or bleeding problems
• take a blood thinning (anticoagulant) or steroid drug
• take other drugs containing prescription or over-the-counter aspirin, ibuprofen, naproxen, or others
• have 3 or more alcoholic drinks every day while using this product
• take more or for a longer time than directed

Drug Facts (continued)

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer
Ask a doctor before use if
• stomach bleeding warning applies to you
• you have a history of stomach problems, such as heartburn
• you have high blood pressure, heart disease, liver disease, or kidney disease
• you are taking a diuretic or you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for gout, diabetes or arthritis

Stop use and ask a doctor if
• an allergic reaction occurs. Seek medical help right away, if you experience any of the following signs of stomach bleeding:
• feel faint
• vomit blood
• have bloody or black stools
• pain gets worse or lasts more than 10 days
• ringing in the ears or a loss of hearing occurs
• indigestion or swelling in breast
• new symptoms occur
These could be signs of a serious condition
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away

Directions
• drink a full glass of water with each dose
• adults and children 12 years of age and over, take 4 to

Drug Facts (continued)

8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
• children under 12 years of age do not use unless directed by a doctor

Other information
• store at 20-25°C (68-77°F)
• close cap tightly after use

Inactive ingredients anhydrous lactose, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, iron oxide yellow, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polyethylene glycol, sodium hydroxide, sodium lauryl sulfate, starch, talc, triethyl citrate

Questions or comments? 1-877-298-4008

This product is not manufactured or distributed by Foundation Consumer-Healthcare LLC, owner of the registered trademark St. Joseph® Low Dose 81 mg Aspirin.
DISTRIBUTED BY:
BETTER LIVING BRANDS LLC
P.O. BOX 99
PLEASANTON, CA 94566-0099
41-888-723-2929
482R 0623

Coating
Omit

Compare to St. Joseph® Low Dose Safety Coated 81 mg Aspirin active ingredient
NDC 700-429-20
Actual Size
200 TABLETS
Low Dose Aspirin 81 mg
Pain reliever (NSAID)
• Enteric coated
• Aspirin regimen**



Coating
Omit

**Talk to your doctor or other healthcare provider before using the product for your heart.



NDC 21130-428-20



Low Dose Aspirin 81 mg
Pain reliever (NSAID)

- Enteric coated
- Aspirin regimen**

200 TABLETS

*Ask your doctor or other healthcare provider before using this product for your best. **READ AND KEEP CAREFULLY COMPLETE PRODUCT WARNINGS AND DIRECTIONS. THIS PRODUCT IS A MEDICINE AND SHOULD BE USED ONLY AS DIRECTED ON THE LABEL OR IN THE INFORMATION STATEMENT.**

Drug Facts

Active ingredient (in each tablet)
Aspirin 81 mg (NSAID)*

Purpose
Nonsteroidal anti-inflammatory drug

Uses ■ Temporarily relieves minor aches and pain ■ Ask your doctor for other uses for enteric coated 81 mg Aspirin.

Warnings
Heart's symptoms: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms may be signs of Reye's syndrome, a rare but serious illness. Allergies about: Aspirin may cause a severe allergic reaction which may include: ■ Hives ■ Facial swelling ■ Asthma (wheezing) ■ Shock

Side effects ■ Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you take aspirin for a long time, take higher doses, or take other drugs that may increase your risk of bleeding. ■ Take a blood thinner (anticoagulant) or steroid drug ■ Take other drugs combining prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ Have 3 or more alcoholic drinks every day while using this product ■ Take more or less than directed ■ Are pregnant or breast-feeding, ask a health professional before use. ■ Keep out of reach of children.

Directions ■ Drink a full glass of water with each dose ■ Adults and children 17 years of age and over: Take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours. Pains directed by a doctor. ■ Children 12 years of age or not use unless directed by a doctor.

Questions or comments?
1-877-234-4088
© 2012 BETTER LIVING BRANDS, LLC
10000 W. 16th Avenue, Suite 100, Golden, CO 80601
NADA 141-224-001
4420.0625.5264

how2recycle.info
51
0122



PLASTIC BOTTLE

LOVE IT OR IT'S ON US.

Lot No: _____
Exp. Date: _____

Varnish Omit Area

ASPIRIN 81MG

aspirin tablet, delayed release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:21130-428
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	

SODIUM HYDROXIDE (UNII: 55X04QC32I)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
TALC (UNII: 7SEV7J4R1U)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER (UNII: NX76LV5T8J)	
STARCH, CORN (UNII: O8232NY3SJ)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

Product Characteristics

Color	pink (Peach Colored tablets)	Score	no score
Shape	ROUND	Size	7mm
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21130-428-12	1 in 1 CARTON	08/08/2023	
1		120 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:21130-428-20	1 in 1 CARTON	08/08/2023	
2		200 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	08/08/2023	

Labeler - BETTER LIVING BRANDS LLC. (009137209)

Registrant - TIME CAP LABORATORIES, INC. (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
TIME CAP LABORATORIES, INC.		037052099	manufacture(21130-428)

