

NICORETTE- nicotine polacrilex lozenge
Haleon US Holdings LLC

Drug Facts

Active ingredient (in each lozenge) (2 mg)

Nicotine polacrilex, 2 mg

Active ingredient (in each lozenge) (4 mg)

Nicotine polacrilex, 4 mg

Purpose

Stop smoking aid

Use

- reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings

If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider.

Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- stomach ulcer or diabetes
- history of seizures

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

Stop use and ask a doctor if

- mouth problems occur
- persistent indigestion or severe sore throat occurs

- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets.

Nicotine lozenges may have enough nicotine to make children and pets sick. If you need to remove the lozenge, wrap it in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions (2 mg)

- **if you are under 18 years of age, ask a doctor before use. No studies have been done to show if this product will work for you.**
- before using this product, read the enclosed User’s Guide for complete directions and other important information
- begin using the lozenge on your quit day
- **if you smoke your first cigarette within 30 minutes of waking up,** use 4 mg nicotine lozenge
- **if you smoke your first cigarette more than 30 minutes after waking up,** use 2 mg nicotine lozenge according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 lozenge every 1 to 2 hours	1 lozenge every 2 to 4 hours	1 lozenge every 4 to 8 hours

- **nicotine lozenge is a medicine and must be used a certain way to get the best results**
- place the lozenge in your mouth and allow the lozenge to slowly dissolve. Minimize swallowing. **Do not chew or swallow lozenge.**
- you may feel a warm or tingling sensation
- occasionally move the lozenge from one side of your mouth to the other until completely dissolved
- do not eat or drink 15 minutes before using or while the lozenge is in your mouth
- to improve your chances of quitting, use at least 9 lozenges per day for the first 6 weeks
- do not use more than one lozenge at a time or continuously use one lozenge after another since this may cause you hiccups, heartburn, nausea or other side effects
- **do not use more than 5 lozenges in 6 hours. Do not use more than 20 lozenges per day.**
- it is important to complete treatment. If you feel you need to use the lozenge for a longer period to keep from smoking, talk to your health care provider.

Directions (4 mg)

- **if you are under 18 years of age, ask a doctor before use. No studies have been done to show if this product will work for you.**
- before using this product, read the enclosed User’s Guide for complete directions and other important information

- begin using the lozenge on your quit day
- **if you smoke your first cigarette more than 30 minutes after waking up**, use 2 mg nicotine lozenge
- **if you smoke your first cigarette within 30 minutes of waking up**, use 4 mg nicotine lozenge according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 lozenge every 1 to 2 hours	1 lozenge every 2 to 4 hours	1 lozenge every 4 to 8 hours

- **nicotine lozenge is a medicine and must be used a certain way to get the best results**
- place the lozenge in your mouth and allow the lozenge to slowly dissolve. Minimize swallowing. **Do not chew or swallow lozenge.**
- you may feel a warm or tingling sensation
- occasionally move the lozenge from one side of your mouth to the other until completely dissolved
- do not eat or drink 15 minutes before using or while the lozenge is in your mouth
- to improve your chances of quitting, use at least 9 lozenges per day for the first 6 weeks
- do not use more than one lozenge at a time or continuously use one lozenge after another since this may cause you hiccups, heartburn, nausea or other side effects
- **do not use more than 5 lozenges in 6 hours. Do not use more than 20 lozenges per day.**
- it is important to complete treatment. If you feel you need to use the lozenge for a longer period to keep from smoking, talk to your health care provider.

Other information (Ice Mint)

- store at 20 - 25°C (68 - 77°F)
- Store in the original container

Inactive ingredients (Ice Mint)

acesulfame potassium, flavors, hypromellose, magnesium stearate, mannitol, microcrystalline cellulose, polysorbate 80, potassium aluminum silicate, sodium carbonate, xanthan gum

Questions or comments?

call toll-free **1-888-569-1743**

Principal Display Panel

NDC 0135-0625-02

NEW

Nicorette

nicotine polacrilex lozenge, 2 mg •stop smoking aid

Lozenge

Coated

Ice Mint

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.

If you smoke your first cigarette **WITHIN 30 MINUTES** of waking up, use Nicorette 4mg Lozenge.

2 mg

80 LOZENGES, 2mg EACH

(4 Packs of 20)

Includes User's Guide

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. **Use Enough** -Use **at least 9 Nicorette Lozenges** per day during the first six weeks.
3. **Use Long Enough** -Use Nicorette for the full 12 weeks.
4. **Use With a Support Program** as directed in the enclosed User's Guide.

For more information and for a FREE individualized stop smoking program, please visit www.Nicorette.com or see inside for more details.

TO OPEN

PUSH

LIFT

TO CLOSE

Push Cap Closed until you hear a CLICK to ensure child resistance

CLICK

Distributed By:

GlaxoSmithKline Consumer Healthcare

Warren, NJ 07059

Made in the Switzerland

Trademarks are owned by or licensed to the GSK group of companies.

©2018 GSK group of companies or its licensor.

106600XA

NDC 0135-0625-02

gsk

Nicorette

nicotine polacrilex lozenge, 2 mg • stop smoking aid

Lozenge



NEW

Coated *Ice Mint*

FOR THOSE WHO
SMOKE THEIR
FIRST CIGARETTE
MORE THAN 30
MINUTES AFTER
WAKING UP.

If you smoke your
first cigarette
WITHIN 30
MINUTES of waking
up, use Nicorette
4 mg Lozenge.

2 mg

80 LOZENGES, 2 mg EACH
(4 Packs of 20)

**Includes
User's Guide**

Principal Display Panel

NDC 0135-0626-02

NEW

Nicorette

nicotine polacrilex lozenge, 4 mg • stop smoking aid

Lozenge

Coated Ice Mint

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.

If you smoke your first cigarette **MORE THAN 30 MINUTES** after waking up, use Nicorette 2 mg Lozenge.

4 mg

80 LOZENGES, 2mg EACH

(4 Packs of 20)

Includes User's Guide

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. **Use Enough** -Use **at least 9 Nicorette Lozenges** per day during the first six weeks.
3. **Use Long Enough** -Use Nicorette for the full 12 weeks.
4. **Use With a Support Program** as directed in the enclosed User's Guide.

For more information and for a FREE individualized stop smoking program, please visit www.Nicorette.com or see inside for more details.

TO OPEN

PUSH

LIFT

TO CLOSE

Push Cap Closed until you hear a CLICK to ensure child resistance

CLICK

Distributed By:

GlaxoSmithKline Consumer Healthcare

Warren, NJ 07059

Made in the Switzerland

Trademarks are owned by or licensed to the GSK group of companies.

©2018 GSK group of companies or its licensor.

106608XA

NDC 0135-0626-02



Nicorette

nicotine polacrilex lozenge, 4 mg • stop smoking aid



NEW

Coated Ice Mint

FOR THOSE WHO
SMOKE THEIR
FIRST CIGARETTE
WITHIN 30
MINUTES OF
WAKING UP.

If you smoke your
first cigarette
MORE THAN 30
MINUTES after
waking up, use
Nicorette 2 mg
Lozenge.

4 mg

80 LOZENGES, 4 mg EACH
(4 Packs of 20)

Includes
User's Guide

NICORETTE

nicotine polacrilex lozenge

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0135-0625
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)	NICOTINE	2 mg

Inactive Ingredients

Ingredient Name	Strength
POLACRILIN (UNII: RCZ785HI7S)	
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MANNITOL (UNII: 3OWL53L36A)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
POTASSIUM (UNII: RWP5GA015D)	
ALUMINUM SILICATE (UNII: T1FAD4SS2M)	
SODIUM CARBONATE (UNII: 45P3261C7T)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	white (white to off-white)	Score	no score
Shape	OVAL	Size	15mm
Flavor	MINT (Ice Mint)	Imprint Code	n;2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0135-0625-01	1 in 1 CARTON	04/11/2019	
1		20 in 1 CONTAINER; Type 0: Not a Combination Product		
2	NDC:0135-0625-02	4 in 1 CARTON	04/11/2019	
2		20 in 1 CONTAINER; Type 0: Not a Combination Product		
3	NDC:0135-0625-03	6 in 1 CARTON	10/23/2019	
3		20 in 1 CONTAINER; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA021330	04/11/2019	

NICORETTE

nicotine polacrilex lozenge

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0135-0626
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)	NICOTINE	4 mg

Inactive Ingredients

Ingredient Name	Strength
POTASSIUM (UNII: RWP5GA015D)	
ALUMINUM SILICATE (UNII: T1FAD4SS2M)	
SODIUM CARBONATE (UNII: 45P3261C7T)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
XANTHAN GUM (UNII: TTV12P4NEE)	
POLACRILIN (UNII: RCZ785HI7S)	
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MANNITOL (UNII: 3OWL53L36A)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	

Product Characteristics

Color	white (white to off-white)	Score	no score
Shape	OVAL	Size	15mm
Flavor	MINT (Ice Mint)	Imprint Code	n;4
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0135-0626-01	1 in 1 CARTON	04/11/2019	
1		20 in 1 CONTAINER; Type 0: Not a Combination Product		
2	NDC:0135-0626-02	4 in 1 CARTON	04/11/2019	

2		20 in 1 CONTAINER; Type 0: Not a Combination Product		
3	NDC:0135-0626-03	6 in 1 CARTON	10/23/2019	
3		20 in 1 CONTAINER; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA021330	04/11/2019	

Labeler - Haleon US Holdings LLC (079944263)

Revised: 7/2024

Haleon US Holdings LLC