

**ACID REDUCER- ranitidine tablet**  
**Fred's Inc.**

-----

***Drug Facts***

***ACTIVE INGREDIENT (IN EACH TABLET)***

Ranitidine 150 mg (as ranitidine hydrochloride USP, 168 mg)

***PURPOSE***

Acid reducer

***USES***

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by certain foods and beverages

***WARNINGS***

**Allergy alert:** Do not use if you are allergic to ranitidine or other acid reducers

**Do not use**

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers
- if you have kidney disease, except under the advice and supervision of a doctor

**Ask a doctor before use if you have**

- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

**Stop use and ask a doctor if**

- your heartburn continues or worsens
- you need to take this product for more than 14 days

**If pregnant or breast-feeding,**

Ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

## ***DIRECTIONS***

- adults and children 12 years and over:
  - to **relieve** symptoms, swallow 1 tablet with a glass of water
  - to **prevent** symptoms, swallow 1 tablet with a glass of water **30 to 60 minutes before** eating food or drinking beverages that cause heartburn
  - can be used up to twice daily (do not take more than 2 tablets in 24 hours)
- children under 12 years: ask a doctor

## ***OTHER INFORMATION***

- **TAMPER EVIDENT: DO NOT USE IF THE CARTON OR PRINTED FOIL UNDER CAP IS OPEN OR TORN.**
- store at 20° - 25° C (68° - 77° F)
- avoid excessive heat or humidity
- this product is sodium and sugar free

## ***INACTIVE INGREDIENTS***

Colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, talc, titanium dioxide

## ***QUESTIONS?***

Call **1-800-406-7984**

## **PRINCIPAL DISPLAY PANEL**

**fred's<sup>®</sup>**

**Maximum Strength**

**Ranitidine Tablets, USP**

**150 mg**

**Acid Reducer**

**Prevents & Relieves Heartburn Associated with Acid Indigestion & Sour Stomach**

**50 TABLETS**

**†Compare To the active ingredient of Zantac 150<sup>®</sup>**

**DISTRIBUTED BY: fred's, Inc.**

**5092927/0112**

**Drug Facts**

**Active ingredient (in each tablet)**  
Ranitidine 150 mg (as ranitidine hydrochloride USP, 168 mg).....Acid reducer

**Uses**  
■ relieves heartburn associated with acid indigestion and sour stomach  
■ prevents heartburn associated with acid indigestion and sour stomach brought on by certain foods and beverages

**Warnings**  
■ Do not use if you are allergic to ranitidine or other acid reducers  
■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor  
■ if you have kidney disease, except under the advice and supervision of a doctor

**fred's**<sup>®</sup>

**Maximum Strength**



Shown at Actual Size

**Ranitidine Tablets, USP  
150 mg**

**Acid Reducer**

Prevents & Relieves Heartburn  
Associated with Acid Indigestion  
& Sour Stomach



**50 TABLETS**

**Compare To**  
the active ingredient of  
**Zantac 150**<sup>®</sup>

Read the directions, consumer information leaflet and warnings before use. Keep the carton. It contains important information.

Expiration Date: **NON VARNISH**

Batch No.

**fred's**<sup>®</sup>

**Maximum Strength**

**Ranitidine Tablets, USP 150 mg**

**Acid Reducer**

**Directions**  
■ adults and children 12 years and over:  
to relieve symptoms, swallow 1 tablet with a glass of water  
to prevent symptoms, swallow 1 tablet with a glass of water 30 to 60 minutes before eating  
■ food or drinking beverages that cause heartburn  
can be used up to twice daily (do not take more than 2 tablets in 24 hours)  
■ children under 12 years: ask a doctor

**Other Information**  
■ TAMPER EVIDENT: DO NOT USE IF THE CARTON OR PRINTED FOIL UNDER CAP IS OPEN OR TORN.  
■ store at 20° - 25° C (68° - 77° F)  
■ avoid excessive heat or humidity  
■ this product is sodium and sugar free

**Inactive ingredients** colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, talc, titanium dioxide

**Questions?** call 1-800-406-7984

This product is not manufactured or distributed by Boehringer Ingelheim Pharmaceuticals Inc., owner of the registered trademark Zantac 150<sup>®</sup>.

0112

MADE IN INDIA



8 4579 112567 8

DISTRIBUTED BY: fred's, Inc.  
4300 NEW GEMWELL RD, MEMPHIS, TN 38118  
www.fredsinc.com

**fred's Promise**  
Not Happy With This Product?  
Just Bring It Back For a Refund!

**Stop use and ask a doctor if** ■ your heartburn continues or worsens  
shoulders, or lightheadedness  
■ heartburn with lightheadedness, sweating or dizziness  
■ chest pain or shoulder pain with shortness of breath, sweating; pain spreading to arms, neck or back


**If pregnant or breast-feeding,** ask a health professional before use.  
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**  
■ adults and children 12 years and over:  
to relieve symptoms, swallow 1 tablet with a glass of water  
to prevent symptoms, swallow 1 tablet with a glass of water 30 to 60 minutes before eating  
■ food or drinking beverages that cause heartburn  
can be used up to twice daily (do not take more than 2 tablets in 24 hours)  
■ children under 12 years: ask a doctor

**Other Information**  
■ TAMPER EVIDENT: DO NOT USE IF THE CARTON OR PRINTED FOIL UNDER CAP IS OPEN OR TORN.  
■ store at 20° - 25° C (68° - 77° F)  
■ avoid excessive heat or humidity  
■ this product is sodium and sugar free

**Inactive ingredients** colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, talc, titanium dioxide

**Questions?** call 1-800-406-7984




1262605

Expiration Date: **NON VARNISH**

Batch No.



5092927



## ACID REDUCER

ranitidine tablet

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:55315-351
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
RANITIDINE HYDROCHLORIDE (UNII: BK76465IHM) (RANITIDINE - UNII:884KT10YB7)	RANITIDINE	150 mg

### Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	pink	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	12mm
<b>Flavor</b>		<b>Imprint Code</b>	9R
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55315-351-50	1 in 1 CARTON		
1		50 in 1 BOTTLE		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA200536	03/30/2012	

**Labeler** - Fred's Inc. (005866116)

## Establishment

Name	Address	ID/FEI	Business Operations
Shasun Pharmaceuticals Limited		915786829	manufacture(55315-351)

Revised: 1/2013

Fred's Inc.