

**COLD AND FLU HBP- acetaminophen, chlorpheniramine maleate,  
dextromethorphan hbr tablet, film coated  
Rite Aid Corporation**

-----  
**Rite Aid 44-738 Cold and Flu**

***Active ingredients (in each caplet)***

Acetaminophen 325 mg  
Chlorpheniramine maleate 2 mg  
Dextromethorphan HBr 10 mg

***Purpose***

Pain reliever/fever reducer  
Antihistamine  
Cough suppressant

***Uses***

- temporarily relieves these common cold and flu symptoms:
  - minor aches and pains
  - sore throat
  - sneezing
  - headache
  - runny nose
  - cough
- temporarily reduces fever

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- blisters
- rash
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a

doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if you have**

- difficulty in urination due to enlargement of the prostate gland
- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema

### **Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

### **When using this product**

- avoid alcoholic beverages
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- excitability may occur, especially in children
- use caution when driving a motor vehicle or operating machinery

### **Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### ***Directions***

- **do not take more than directed**
- adults and children 12 years and over: take 2 caplets every 4 hours while symptoms persist. Do not take more than 10 caplets in 24 hours.
- children under 12 years: ask a doctor

### ***Other information***

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- protect from excessive moisture
- see end flap for expiration date and lot number

### ***Inactive ingredients***

corn starch, crospovidone, FD&C red #40 aluminum lake, lecithin, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

### ***Questions or comments?***

**1-800-426-9391**

### ***Principal display panel***

NDC 11822-7380-9

Compare to the active ingredients in  
**Coricidin® HBP Maximum Strength Flu\***

**MAXIMUM STRENGTH  
COLD &  
FLU HBP**

**ACETAMINOPHEN  
CHLORPHENIRAMINE MALEATE  
DEXTROMETHORPHAN HBr**

PAIN RELIEVER/FEVER REDUCER  
ANTIHISTAMINE/COUGH SUPPRESSANT

Relieves body aches, pains & headache,  
fever, cough, runny nose, sore throat, sneezing

For people with high blood  
pressure

Decongestant-free

ACTUAL SIZE

**20**

CAPLETS

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING**

**PARENTS:**

Learn about teen medicine abuse  
www.StopMedicineAbuse.org

\*This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Coricidin®  
HBP Maximum Strength Flu.

50844 ORG101973809

**SATISFACTION GUARANTEE**

If you're not satisfied, we'll happily refund your money.

**DISTRIBUTED BY:**

RITE AID, 30 HUNTER LANE,  
CAMP HILL, PA 17011  
www.riteaid.com

\*This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Coricidin® HBP Maximum Strength Flu. 50844 ORG101973809

**PARENTS:**  
Learn about teen medicine abuse  
www.StopMedicineAbuse.org

**SATISFACTION GUARANTEE**  
If you're not satisfied, we'll happily refund your money.

**DISTRIBUTED BY:**  
RITE AID, 30 HUNTER LANE,  
CAMP HILL, PA 17011  
www.riteaid.com

NDC 11822-7380-9      Compare to the active ingredients in  
Coricidin® HBP Maximum Strength Flu\*

# MAXIMUM STRENGTH COLD & FLU HBP

**ACETAMINOPHEN  
CHLORPHENIRAMINE MALEATE  
DEXTROMETHORPHAN HBr**

PAIN RELIEVER/FEVER REDUCER  
ANTIHISTAMINE/COUGH SUPPRESSANT

Relieves body aches, pains & headache,  
fever, cough, runny nose, sore throat, sneezing

For people with high blood  
pressure  
Decongestant-free

**Drug Facts (continued)**

protect from excessive moisture  
see end flap for expiration date and lot number

**Inactive ingredients**  
FD&C red #40 aluminum lake, lecithin, magnesium stearate, corn starch, croscopolone.

**Questions or comments?** 1-800-426-9391

**Drug Facts (continued)**

temporarily relieves these common cold and flu symptoms:  
minor aches and pains    sore throat  
runny nose    cough  
sneezing    headache  
temporarily reduces fever

**Drug Facts (continued)**

**Warnings**  
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take  
more than 4,000 mg of acetaminophen in 24 hours  
with other drugs containing acetaminophen  
3 or more alcoholic drinks every day while using this product  
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:  
blisters    rash    skin reddening  
If a skin reaction occurs, stop use and seek medical help right away.  
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.  
Do not use    with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.  
if you have ever had an allergic reaction to this product or any of its ingredients  
Ask a doctor before use if you have  
difficulty in urination due to enlargement of the prostate gland    liver disease    glaucoma  
cough that occurs with too much phlegm (mucus)  
a breathing problem or persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema  
Ask a doctor or pharmacist before use if you are  
taking sedatives or tranquilizers

**Drug Facts (continued)**

**When using this product**  
avoid alcoholic beverages  
marked drowsiness may occur  
alcohol, sedatives, and tranquilizers may increase drowsiness  
excitability may occur, especially in children  
use caution when driving a motor vehicle or operating machinery  
**Stop use and ask a doctor if**  
pain or cough gets worse or lasts more than 7 days    new symptoms occur  
fever gets worse or lasts more than 3 days  
redness or swelling is present  
cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.  
**If pregnant or breast-feeding**, ask a health professional before use.  
**Keep out of reach of children.** In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

ACTUAL SIZE

## 20 CAPLETS

**Drug Facts (continued)**

temporarily relieves these common cold and flu symptoms:  
minor aches and pains    sore throat  
runny nose    cough  
sneezing    headache  
temporarily reduces fever

**Drug Facts (continued)**

**Warnings**  
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take  
more than 4,000 mg of acetaminophen in 24 hours  
with other drugs containing acetaminophen  
3 or more alcoholic drinks every day while using this product  
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:  
blisters    rash    skin reddening  
If a skin reaction occurs, stop use and seek medical help right away.  
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.  
Do not use    with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.  
if you have ever had an allergic reaction to this product or any of its ingredients  
Ask a doctor before use if you have  
difficulty in urination due to enlargement of the prostate gland    liver disease    glaucoma  
cough that occurs with too much phlegm (mucus)  
a breathing problem or persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema  
Ask a doctor or pharmacist before use if you are  
taking sedatives or tranquilizers

**Drug Facts (continued)**

**When using this product**  
avoid alcoholic beverages  
marked drowsiness may occur  
alcohol, sedatives, and tranquilizers may increase drowsiness  
excitability may occur, especially in children  
use caution when driving a motor vehicle or operating machinery  
**Stop use and ask a doctor if**  
pain or cough gets worse or lasts more than 7 days    new symptoms occur  
fever gets worse or lasts more than 3 days  
redness or swelling is present  
cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.  
**If pregnant or breast-feeding**, ask a health professional before use.  
**Keep out of reach of children.** In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Drug Facts (continued)**

**Directions**  
do not take more than directed  
adults and children 12 years and over: take 2 caplets every 4 hours while symptoms persist. Do not take more than 10 caplets in 24 hours.  
children under 12 years: ask a doctor

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

0 11822 20214 5

No print/No varnish  
Lot & Exp date

B-1702-738-09-R2  
ORG101973809

Rite Aid 44-738

**COLD AND FLU HBP**

acetaminophen, chlorpheniramine maleate, dextromethorphan hbr tablet, film coated

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:11822-7380
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>CHLORPHENIRAMINE MALEATE</b> (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg

**Inactive Ingredients**

Ingredient Name	Strength
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>CROSPVIDONE, UNSPECIFIED</b> (UNII: 2S7830E561)	
<b>FD&amp;C RED NO. 40 ALUMINUM LAKE</b> (UNII: 6T47AS764T)	
<b>LECITHIN, SOYBEAN</b> (UNII: 1DI56QDM62)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

**Product Characteristics**

<b>Color</b>	red	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	16mm
<b>Flavor</b>		<b>Imprint Code</b>	44;738
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	NDC:11822			

1	NDC:11822-7380-9	2 in 1 CARTON	06/25/2021	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>		<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M012		06/25/2021	

**Labeler** - Rite Aid Corporation (014578892)

<b>Establishment</b>			
<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
LNK International, Inc.		832867837	manufacture(11822-7380) , pack(11822-7380)

<b>Establishment</b>			
<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
LNK International, Inc.		832867894	manufacture(11822-7380)

<b>Establishment</b>			
<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
LNK International, Inc.		117025878	manufacture(11822-7380)

Revised: 6/2024

Rite Aid Corporation