

VICKS DAYQUIL AND NYQUIL VAPOCOOL SEVERE COLD AND FLU PLUS CONGESTION- acetaminophen, phenylephrine hydrochloride, and doxylamine succinate
The Procter & Gamble Manufacturing Company

Vicks ® DayQuil™ & Nyquil™ VapoCOOL SEVERE COLD & FLU + CONGESTION, Coated Caplets

DayQuil™ VapoCOOL™ SEVERE COLD & FLU + CONGESTION

Active ingredients (in each caplet)

Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Guaifenesin 200 mg
Phenylephrine HCl 5 mg

Purpose

Pain reliever/Fever reducer
Cough suppressant
Expectorant
Nasal decongestant

Uses

- temporarily relieves common cold/flu symptoms:
- nasal congestion
- sinus congestion & pressure
- cough due to minor throat & bronchial irritation
- minor aches & pains
- headache
- fever
- sore throat
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning:

This product contains acetaminophen.
Severe liver damage may occur if you take

- more than 8 caplets in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy Alert:

Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening
- Blisters
- Rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning:

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

When using this product,

do not use more than directed.

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- do not exceed 8 caplets per 24 hrs

adults & children 12 yrs & over	2 caplets with water every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

each caplet contains:

- sodium 4 mg
- store at no greater than 25°C

Inactive ingredients

croscarmellose sodium, crospovidone, FD&C Yellow No. 6 Aluminum Lake, flavor, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, silicon dioxide, stearic acid, sucralose, talc, titanium dioxide

Questions?

1-800-362-1683

NyQuil™ VapoCOOL™ SEVERE COLD & FLU + CONGESTION

Active ingredients (in each caplet)

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Doxylamine succinate 6.25 mg

Phenylephrine HCl 5 mg

Purpose

Pain reliever/Fever reducer

Cough suppressant

Antihistamine

Nasal decongestant

Uses

temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion & pressure
- cough due to minor throat & bronchial irritation
- cough to help you sleep
- minor aches & pains
- headache
- fever
- sore throat
- runny nose & sneezing •reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage

Warnings

Liver warning:

This product contains acetaminophen.

Severe liver damage may occur if you take

- more than 8 caplets in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy Alert:

acetaminophen may cause severe skin reactions.

Symptoms may include:

- Skin reddening
- Blisters

- Rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning:

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- **do not use more than directed**
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives & tranquilizers may increase drowsiness

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days

- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- do not exceed 8 caplets per 24 hrs

adults & children 12 yrs & over	2 caplets with water every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- store at no greater than 25°C

Inactive ingredients

crospovidone, D&C Yellow No. 10 Aluminum Lake, FD&C Blue No. 1 Aluminum Lake, FD&C Blue No. 2 Aluminum Lake, FD&C Yellow No. 6 Aluminum Lake, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, silicon dioxide, stearic acid, sucralose, talc, titanium dioxide

Questions?

1-800-362-1683

TAMPER EVIDENT:This package is safety-sealed & child resistant. Use only if blisters are intact. If difficult to open, use scissors.

**DIST. BY PROCTER & GAMBLE,
CINCINNATI, OH 45202.**

PRINCIPAL DISPLAY PANEL - Kit Carton

Vicks®

DayQuil™ VapoCOOL™ SEVERE COLD & FLU + CONGESTION

Acetaminophen, Guaifenesin, Phenylephrine HCl, Dextromethorphan HBr

Headache, Fever, Sore Throat, Minor Aches & Pains

Chest Congestion, Thins & Loosens Mucus

Nasal Congestion, Sinus Pressure

Cough

32 DAYQUIL COATED CAPLETS

NyQuil™ VapoCOOL™ SEVERE COLD & FLU + CONGESTION

Acetaminophen, Phenylephrine HCl, Doxylamine Succinate, Dextromethorphan HBr

Headache, Fever, Sore Throat, Minor Aches & Pains

Nasal Congestion, Sinus Pressure

Sneezing, Runny Nose

Cough

16 NYQUIL COATED CAPLETS

32 DAYQUIL COATED CAPLETS, 16 NYQUIL COATED CAPLETS: 48 TOTAL COATED CAPLETS

<p>Drug Facts (continued)</p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p>Directions • take only as directed • do not exceed 8 caplets per 24 hrs adults & children 12 yrs & over 2 caplets with water every 4 hrs children 4 to under 12 yrs ask a doctor children under 4 yrs do not use</p> <p>Other information • store at no greater than 25°C</p> <p>Inactive ingredients crospravonide, D&C Yellow No. 10 Aluminum Lake, FD&C Blue No. 1 Aluminum Lake, FD&C Blue No. 2 Aluminum Lake, FD&C Yellow No. 6 Aluminum Lake, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, silicon dioxide, stearic acid, sucralose, talc, titanium dioxide</p> <p>Questions? 1-800-362-1683</p>	<p>Drug Facts (continued)</p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p>Directions • take only as directed • do not exceed 8 caplets per 24 hrs adults & children 12 yrs & over 2 caplets with water every 4 hrs children 4 to under 12 yrs ask a doctor children under 4 yrs do not use</p> <p>Other information • each caplet contains: sodium 4 mg • store at no greater than 25°C</p> <p>Inactive ingredients crospravonide sodium, crospravonide, FD&C Yellow No. 6 Aluminum Lake, flavor, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, silicon dioxide, stearic acid, sucralose, talc, titanium dioxide</p> <p>Questions? 1-800-362-1683</p>
--	--

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

TAMPER EVIDENT: This package is safety-sealed & child resistant. Use only if blisters are intact. If difficult to open, use scissors.



ACTUAL SIZE



DayQuil
VapoCOOL[®]
SEVERE COLD & FLU
+ CONGESTION
Acetaminophen, Guaifenesin,
Phenylephrine HCl, Dextromethorphan HBr

NyQuil
VapoCOOL[®]
SEVERE COLD & FLU
+ CONGESTION
Acetaminophen, Phenylephrine HCl,
Doxylamine Succinate, Dextromethorphan HBr



3 23900 03846 2



ACTUAL SIZE

Headache, Fever, Sore Throat,
Minor Aches & Pains
Chest Congestion, Thins & Loosens Mucus
Nasal Congestion, Sinus Pressure
Cough

32 DAYQUIL COATED CAPLETS

Headache, Fever, Sore Throat,
Minor Aches & Pains
Nasal Congestion, Sinus Pressure
Sneezing, Runny Nose
Cough

16 NYQUIL COATED CAPLETS: COATED CAPLETS

DIST. BY PROCTER & GAMBLE,
CINCINNATI OH 45202

P&G www.pg.com
www.vicks.com
Patents: www.pg.com/patents
90884399

<p>Drug Facts NyQuil + VapoCool Cold & Flu</p> <p>Active ingredients (in each caplet)</p> <p>Acetaminophen 325 mg Pain reliever/fever reducer Dextromethorphan HBr 10 mg Cough suppressant Doxylamine succinate 6.25 mg Antihistamine Phenylephrine HCl 5 mg Nasal decongestant</p> <p>Uses temporarily relieves common cold/flu symptoms: • nasal congestion • sinus congestion & pressure • cough due to minor throat & bronchial irritation • cough to help you sleep • minor aches & pains • headache • fever • sore throat • runny nose & sneezing • reduces swelling of nasal passages • temporarily restores freer breathing through the nose • promotes nasal and/or sinus drainage</p> <p>Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take • more than 8 caplets in 24 hours, which is the maximum daily amount for this product • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product</p>	<p>Drug Facts DayQuil + VapoCool Cold & Flu</p> <p>Active ingredients (in each caplet)</p> <p>Acetaminophen 325 mg Pain reliever/fever reducer Dextromethorphan HBr 10 mg Cough suppressant Guaifenesin 200 mg Expectorant Phenylephrine HCl 5 mg Nasal decongestant</p> <p>Uses temporarily relieves common cold/flu symptoms: • nasal congestion • sinus congestion & pressure • cough due to minor throat & bronchial irritation • minor aches & pains • headache • fever • sore throat • reduces swelling of nasal passages • temporarily restores freer breathing through the nose • promotes nasal and/or sinus drainage • helps loosen phlegm (mucus) and thin bronchial secretions to aid the bronchial passageways of bothersome mucus and make coughs more productive.</p> <p>Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take</p>
--	---

<p>Drug Facts (continued)</p> <p>Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash If a skin reaction occurs, stop use and seek medical help right away Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p> <p>Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p> <p>Ask a doctor before use if you have • liver disease • heart disease • high blood pressure • thyroid disease • diabetes • glaucoma • cough that occurs with too much phlegm (mucus) • a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema • trouble urinating due to enlarged prostate gland</p> <p>Ask a doctor or pharmacist before use if you are • taking sedatives or tranquilizers • taking the blood thinning drug warfarin</p> <p>When using this product • do not use more than directed • excitability may occur, especially in children • marked drowsiness may occur • avoid alcoholic drinks • be careful when driving a motor vehicle or operating machinery • alcohol, sedatives, and tranquilizers may increase drowsiness</p> <p>Stop use and ask a doctor if • you get nervous, dizzy or sleepless • pain, nasal congestion, or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.</p>	<p>Drug Facts (continued)</p> <p>• more than 8 caplets in 24 hours, which is the maximum daily amount for this product • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product</p> <p>Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash If a skin reaction occurs, stop use and seek medical help right away Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p> <p>Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p> <p>Ask a doctor before use if you have • liver disease • heart disease • high blood pressure • thyroid disease • diabetes • trouble urinating due to enlarged prostate gland • cough that occurs with too much phlegm (mucus) • persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema</p> <p>Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.</p> <p>When using this product, do not use more than directed.</p> <p>Stop use and ask a doctor if • you get nervous, dizzy or sleepless • pain, nasal congestion, or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.</p>
---	---

VICKS DAYQUIL AND NYQUIL VAPOCOOL SEVERE COLD AND FLU PLUS CONGESTION

acetaminophen, phenylephrine hydrochloride, and doxylamine succinate kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69423-988
--------------	----------------	--------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69423-988-24	1 in 1 PACKAGE; Type 1: Convenience Kit of Co-Package	03/22/2021	
2	NDC:69423-988-48	1 in 1 PACKAGE; Type 1: Convenience Kit of Co-Package	03/22/2021	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	16 BLISTER PACK	16
Part 2	8 BLISTER PACK	8

Part 1 of 2

VICKS DAYQUIL VAPOCOOL SEVERE COLD AND FLU PLUS CONGESTION

acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride tablet

Product Information

Item Code (Source)	NDC:69423-987
--------------------	---------------

Route of Administration	ORAL
-------------------------	------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CROSPVIDONE (UNII: 2S7830E561)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
TALC (UNII: 7SEV7J4R1U)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
STARCH, CORN (UNII: O8232NY3SJ)	

Product Characteristics

Color	orange	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	DQ
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	08/01/2018	

Part 2 of 2

VICKS NYQUIL VAPOCOOL SEVERE COLD AND FLU PLUS CONGESTION

acetaminophen, phenylephrine hydrochloride, doxylamine succinate, and dextromethorphan hydrobromide tablet, coated

Product Information

Item Code (Source)		NDC:69423-986		
Route of Administration		ORAL		
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		ACETAMINOPHEN	325 mg	
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)		DOXYLAMINE SUCCINATE	6.25 mg	
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)		PHENYLEPHRINE HYDROCHLORIDE	5 mg	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)		DEXTROMETHORPHAN HYDROBROMIDE	10 mg	
Inactive Ingredients				
Ingredient Name			Strength	
FD&C BLUE NO. 1 ALUMINUM LAKE (UNII: J9EQA3S2JM)				
FD&C BLUE NO. 2 ALUMINUM LAKE (UNII: 4AQJ3LG584)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
POVIDONE (UNII: FZ989GH94E)				
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)				
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)				
ALUMINUM OXIDE (UNII: LMI26O6933)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)				
TALC (UNII: 7SEV7J4R1U)				
CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561)				
STEARIC ACID (UNII: 4ELV7Z65AP)				
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)				
SUCRALOSE (UNII: 96K6UQ3ZD4)				
STARCH, CORN (UNII: O8232NY3SJ)				
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
Product Characteristics				
Color	green	Score	no score	
Shape	OVAL	Size	19mm	
Flavor		Imprint Code	NQ	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	08/01/2018	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/22/2021	

Labeler - The Procter & Gamble Manufacturing Company (004238200)

Revised: 10/2023

The Procter & Gamble Manufacturing Company