

**COLD MULTI-SYMPTOM- acetaminophen, dextromethorphan hbr,
phenylephrine hcl tablet, film coated
Kroger Company**

Kroger 44-470C

Active ingredients (in each caplet)

Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer
Cough suppressant
Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - minor aches and pains
 - nasal congestion
 - headache
 - cough
 - sinus congestion and pressure
 - sore throat
- helps clear nasal passages
- promotes nasal and sinus drainage
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- blisters
- rash
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- diabetes
- heart disease
- thyroid disease
- high blood pressure
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

When using this product

do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed**
- adults and children 12 years and over
 - take 2 caplets every 4 hours
 - swallow whole - do not crush, chew, or dissolve
 - do not take more than 10 caplets in 24 hours
- children under 12 years: ask a doctor

Other information

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°C-30°C (59°F-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

corn starch, croscarmellose sodium, crospovidone, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, stearic acid, sucralose, talc, titanium dioxide

Questions or comments?

1-800-632-6900

Principal Display Panel

COMPARE TO the active ingredients of TYLENOL® COLD MAX *See bottom panel

NDC 30142-470-08

Kroger®

DayTime

Cold Multi-Symptom

Acetaminophen

Pain Reliever/Fever Reducer

Dextromethorphan HBr

Cough Suppressant

Phenylephrine HCl

Nasal Decongestant

NON-DROWSY

Headache

Fever, Sore Throat

Nasal Congestion

Cough

OUR PHARMACIST RECOMMENDED

actual size

24 CAPLETS

TAMPER EVIDENT: DO NOT USE IF PACKAGE IF OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

*Tylenol® Cold Max is a registered trademark of Johnson & Johnson Corporation. New Brunswick, New Jersey 08933, Johnson & Johnson Corporation is not affiliated with The Kroger Co. or this product.

50844 ORG071847008

**DISTRIBUTED BY THE KROGER CO.
CINCINNATI, OHIO 45202**

For **More Product Information, Scan UPC**
Using Your **Kroger App** or Call 800-632-6900

Our Brands. Our Guarantee. Love it or
Your Money Back. www.kroger.com

PARENTS:

Learn about teen medicine abuse
www.StopMedicineAbuse.org

80% SIZE
12-CAPLET-A
(non-FPO Issued)
For Position Only
0 41260 01356 0

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

No Print/No Varnish
Lot and Expiration No.

Drug Facts (continued)
Inactive ingredients (continued)
corn starch, croscarmellose sodium, crospovidone, glycol, polyvinyl alcohol, povidone, silicon dioxide, stearic acid, sucralose, talc, titanium dioxide

Drug Facts (continued)
Questions or comments? 1-800-632-6900
Tylenol® Cold Max is a registered trademark of Johnson & Johnson Corporation, New Brunswick, New Jersey 08933. Johnson & Johnson Corporation is not affiliated with The Kroger Co. or this product. 50844 ORG071847008

Drug Facts (continued)
Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ blisters ■ rash ■ skin redness
If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
Do not use
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
■ if you have ever had an allergic reaction to this product or any of its ingredients
Ask a doctor before use if you have
■ liver disease ■ diabetes ■ heart disease ■ thyroid disease ■ high blood pressure ■ difficulty in urination due to enlargement of the prostate gland ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema
■ cough that occurs with too much phlegm (mucus)



Drug Facts (continued)
Directions
■ do not take more than directed
■ adults and children 12 years and over
■ take 2 caplets every 4 hours
■ swallow whole - do not crush, chew, or dissolve
■ do not take more than 10 caplets in 24 hours
■ children under 12 years: ask a doctor
Other information
■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
■ see end flap for expiration date and lot number

Drug Facts
Active ingredients (in each caplet)
Acetaminophen 325 mg
Dextromethorphan Hbr 10 mg
Phenylephrine HCl 5 mg
Nasal decongestant
Purpose
Pain reliever/fever reducer
Cough suppressant
Nasal decongestant

Drug Facts (continued)
Uses
temporarily relieves these common cold and flu symptoms:
■ minor aches and pains ■ headache ■ nasal congestion ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ nervousness, dizziness, or sleeplessness occur ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
Keep out of reach of children. In case of accidental overdose, get medical help or contact Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
Directions
■ do not take more than directed
■ adults and children 12 years and over
■ take 2 caplets every 4 hours
■ swallow whole - do not crush, chew, or dissolve
■ do not take more than 10 caplets in 24 hours
■ children under 12 years: ask a doctor

COMPARE TO the active ingredients of **TYLENOL® COLD MAX*** See bottom panel

NDC 30142-470-08

DayTime Cold Multi-Symptom


Acetaminophen
Pain Reliever/Fever Reducer

Dextromethorphan HBr
Cough Suppressant

Phenylephrine HCl
Nasal Decongestant

NON-DROWSY

Headache
Fever, Sore Throat
Nasal Congestion
Cough



actual size **24 CAPLETS**

DISTRIBUTED BY THE KROGER CO.
CINCINNATI, OHIO 45202
For more Product Information, Scan UPC
Using Your Kroger App or Call 800-632-6900
Our Brands. Our Guarantee. Love It or
Your Money Back. www.kroger.com

PLEASE RECYCLE

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

8-1050-470C-08-CM
ORG071847008

Kroger 44-470C

COLD MULTI-SYMPOTOM
acetaminophen, dextromethorphan hbr, phenylephrine hcl tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:30142-470
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	white	Score	no score
Shape	OVAL	Size	17mm
Flavor	MENTHOL	Imprint Code	44;470
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:30142-470-08	2 in 1 CARTON	05/12/2021	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/12/2021	

Labeler - Kroger Company (006999528)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(30142-470) , pack(30142-470)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(30142-470)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(30142-470)

Revised: 5/2024

Kroger Company