BASIC CARE NIGHTTIME COLD AND FLU- acetaminophen, dextromethorphan hbr, doxylamine succinate capsule, liquid filled
Amazon.com Services LLC

Amazon Nighttime Cold & Flu Drug Facts

Active ingredients (in each softgel)

Acetaminophen 325 mg

Dextromethorphan HBr 15 mg

Doxylamine succinate 6.25 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Uses

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever
- runny nose and sneezing

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough as occurs with smoking, asthma, or emphysema
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed – see overdose warning
- do not exceed 4 doses per 24 hours

adults & children 12 yrs & over	2 softgels with water every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- store at 20-25˚C (68-77˚F)

Inactive ingredients

D&C yellow no. 10, edible ink*, FD&C blue no. 1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution *may contain this ingredient

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

Multi-Symptom Relief

Compare to Vicks® NyQuil® Cold & Flu active ingredients

Nighttime Cold & Flu

Acetaminophen, Dextromethorphan HBr

Doxylamine Succinate

Pain Reliever/Fever Reducer, Antihistamine

Cough Suppressant

Alcohol Free

actual size

48 SOFTGELS

amazon
basic care
Multi-Symptom Relief
Compare to Vicks® NyQuil® Cold & Flu active ingredients**
Nighttime Cold & Flu
amazon
basic care
Multi-Symptom Relief
NDC 72288-056-67
Compare to Vicks® NyQuil® Cold & Flu active ingredients**
Nighttime Cold & Flu
Acetaminophen, Dextromethorphan HBr
Doxylamine Succinate
Pain Reliever/Fever Reducer, Antihistamine
Cough Suppressant
Alcohol Free
actual size
48 SOFTGELS

CONVENIENT RECLOSING TAB

Drug Facts

DO NOT USE IF BLISTER UNIT IS BROKEN OR TORN

Active ingredients (in each softgel)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 15 mg	Cough suppressant
Doxylamine succinate 6.25 mg	Antihistamine

Uses temporarily relieves common cold/flu symptoms:
- cough due to minor throat and bronchial irritation
- minor aches and pains
- runny nose and sneezing
- sore throat
- headache
- fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have
- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough as occurs with smoking, asthma, or emphysema
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are
- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product
- excitability may occur, especially in children
- be careful when driving a motor vehicle or operating machinery
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- marked drowsiness may occur

Stop use and ask a doctor if
- pain or cough gets worse or lasts more than 7 days
- redness or swelling is present
- fever gets worse or lasts more than 3 days
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. ▶

OPEN OTHER END

3 70030 14519 5

Drug Facts (continued)

Directions

- take only as directed – see overdose warning
- do not exceed 4 doses per 24 hours

adults & children 12 yrs & over	2 softgels with water every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information
- store at 20-25°C (68-77°F)

Inactive ingredients D&C yellow no. 10, edible ink*, FD&C blue no. 1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution
*may contain this ingredient

Questions or comments? 1-800-719-9260

MADE IN CANADA

DISTRIBUTED BY:
AMAZON.COM SERVICES LLC
410 TERRY AVENUE N.
SEATTLE, WA 98109

© 2020 AMAZON.COM, INC. OR ITS AFFILIATES. ALL RIGHTS RESERVED.
BASIC CARE AND ALL RELATED LOGOS ARE TRADEMARKS OF AMAZON.COM, INC. OR ITS AFFILIATES.

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

**This product is not manufactured or distributed by Procter & Gamble, distributor of Vicks® NyQuil® Cold & Flu.

Gluten Free

BASIC CARE NIGHTTIME COLD AND FLU

acetaminophen, dextromethorphan hbr, doxylamine succinate capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	**Item Code (Source)**	NDC:72288-056
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

Inactive Ingredients

Ingredient Name	Strength
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WQ0SDW1A)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	
SORBITAN (UNII: 6O92ICV9RU)	

Product Characteristics

Color	GREEN (clear)	**Score**	no score
Shape	OVAL	**Size**	20mm
Flavor		**Imprint Code**	056
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72288-056-67	24 in 1 CARTON	12/28/2020	

1		2 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:72288-056-62	12 in 1 CARTON	09/07/2021	
2		2 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	12/28/2020	

Labeler - Amazon.com Services LLC (128990418)

Revised: 10/2024 Amazon.com Services LLC