

MUCUS RELIEF SINUS PRESSURE PAIN AND COUGH- acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride tablet, coated
WINCO FOODS, LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

1199-WNC-2023-0411

Drug Facts

Active ingredients (in each caplet)	Purpose
Acetaminophen 325 mg	Pain reliever
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves:
 - headache
 - nasal congestion
 - minor aches and pains
 - sinus congestion and pressure
 - cough
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning

- This product contains acetaminophen. Severe liver damage may occur if you take
- more than 12 caplets in 24 hours, which is the maximum daily amount
 - with other drugs containing acetaminophen
 - 3 or more alcoholic drinks daily while using this product

Allergy alert

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- diabetes
- liver disease
- heart disease
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see Overdose warning)**

- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years of age and older: take 2 caplets every 4 hours
- children under 12 years of age: do not use

Other information

- store between 20-25°C (68-77°F) in a dry place

Inactive ingredients

colloidal silicon dioxide, corn starch, croscarmellose sodium, crospovidone, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, stearic acid, talc, titanium dioxide

Questions?

Call 800-824-1706 Monday - Friday 9am - 4pm MST

PRINCIPAL DISPLAY PANEL

WinCo® Foods

Maximum Strength

Mucus Relief

SINUS PRESSURE, PAIN & COUGH

ACETAMINOPHEN,

Dextromethorphan HBr, Guaifenesin, Phenylephrine HCl

Pain Reliever, Cough Suppressant, Expectorant, Nasal Decongestant

* Compare to the active ingredients in MUCINEX® Sinus-Max® Pressure, Pain & Cough

Relieves Sinus Pressure & Congestion, Headache, and Cough

Thins & Loosens Mucus

For Ages 12+

20 Caplets

NC

Maximum Strength
Mucus Relief
 SINUS PRESSURE,
 PAIN & COUGH

ACETAMINOPHEN,
 Dextromethorphan HBr, Guaifenesin, Phenylephrine HCl
 Pain Reliever, Cough Suppressant, Expectorant, Nasal Decongestant

20
 Caplets

*Compare to the active ingredients in MUCINEX®
 Sinus-Max® Pressure, Pain & Cough

Relieves Sinus Pressure & Congestion,
 Headache, and Cough

Thins & Loosens Mucus

For Ages 12+

NC



Maximum Strength

Mucus Relief

SINUS PRESSURE, PAIN & COUGH

Drug Facts

Active ingredients (in each caplet): Purpose
 Acetaminophen 325 mg.....Pain reliever
 Dextromethorphan HBr 10 mg.....Cough suppressant
 Guaifenesin 200 mg.....Expectorant
 Phenylephrine HCl 5 mg.....Nasal decongestant

Uses ■ temporarily relieves: ■ headache
 ■ nasal congestion ■ minor aches and pains
 ■ sinus congestion and pressure ■ cough
 ■ promotes nasal and/or sinus drainage
 ■ helps loosen phlegm (mucus) and thin bronchial
 secretions to rid the bronchial passageways of
 bothersome mucus and make coughs
 more productive

Warnings
 Liver warning: This product contains acetaminophen.
 Severe liver damage may occur if you take
 ■ more than 12 caplets in 24 hours, which is the
 maximum daily amount
 ■ with other drugs containing acetaminophen
 ■ 3 or more alcoholic drinks daily while using
 this product
Allergy alert: Acetaminophen may cause severe
 skin reactions. Symptoms may include:
 ■ skin reddening ■ blisters ■ rash
 If a skin reaction occurs, stop use and seek medical
 help right away.

Do not use
 ■ with any other drug containing acetaminophen
 (prescription or nonprescription). If you are not
 sure whether a drug contains acetaminophen,
 ask a doctor or pharmacist.
 ■ if you are now taking a prescription monoamine
 oxidase inhibitor (MAOI) (certain drugs for
 depression, psychiatric, or emotional conditions,
 or Parkinson's disease), or for 2 weeks after stopping
 the MAOI drug, if you do not know if your
 prescription drug contains an MAOI, ask a doctor
 or pharmacist before taking this product.

Ask a doctor before use if you have ■ diabetes
 ■ liver disease ■ heart disease
 ■ high blood pressure ■ thyroid disease
 ■ trouble urinating due to an enlarged prostate gland
 ■ cough that occurs with too much phlegm (mucus) ▶

Drug Facts (continued)

■ persistent or chronic cough such as occurs with
 smoking, asthma, chronic bronchitis, or emphysema
Ask a doctor or pharmacist before use if you
 are taking the blood thinning drug warfarin
When using this product do not use more than directed

Stop use and ask a doctor if
 ■ nervousness, dizziness, or sleeplessness occur
 ■ pain, nasal congestion, or cough gets worse or
 lasts more than 7 days
 ■ fever gets worse or lasts more than 3 days
 ■ redness or swelling is present
 ■ new symptoms occur
 ■ cough comes back or occurs with rash or
 headache that lasts

These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health
professional before use.
Keep out of reach of children.
Overdose warning: Taking more than the
 recommended dose (overdose) may cause liver
 damage. In case of overdose, get medical help or
 contact a Poison Control Center right away
 (1-800-222-1222). Quick medical attention is
 critical for adults as well as for children even if you
 do not notice any signs or symptoms.

Directions ■ do not take more than
 directed (see Overdose warning)
 ■ do not take more than 12 caplets in any
 24-hour period
 ■ adults and children 12 years of age and over:
 take 2 caplets every 4 hours
 ■ children under 12 years of age: do not use

Other information
 ■ store between 20-25°C (68-77°F) in a dry place
 ■ retain outer package for complete product
 information and warnings

Inactive ingredients colloidal silicon
 dioxide, croscarmellose sodium, crospovidone,
 FD&C red #40 aluminum lake, FD&C yellow #6
 aluminum lake, magnesium stearate, maltodextrin,
 microcrystalline cellulose, polyethylene glycol,
 polyvinyl alcohol, povidone, pregelatinized starch,
 stearic acid, talc, titanium dioxide ▶



DISTRIBUTED BY:
 WINCO FOODS, LLC, BOISE, ID 83704
100% SATISFACTION GUARANTEE
 Comments & Questions? www.wincofoods.com

**DO NOT USE IF BLISTER UNITS
 ARE TORN OR BROKEN**

Drug Facts (continued)
 Call 800-824-1706
 Monday - Friday 9am - 4pm MST
PER 4-HOUR DOSE
MAXIMUM STRENGTH

WF2209 F119902WNC_RO

0 70552 23070 5

WinCo FOODS
Maximum Strength
Mucus Relief
 SINUS PRESSURE, PAIN & COUGH

MUCUS RELIEF SINUS PRESSURE PAIN AND COUGH

acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:67091-299
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	red	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	AAA;1165
Contains			

Packaging

#	Item Code	Package Description	Marketing Start	Marketing End
---	-----------	---------------------	-----------------	---------------

#	Item Code	Package Description	Date	Date
1	NDC:67091-299-02	2 in 1 CARTON	04/11/2023	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	04/11/2023	

Labeler - WINCO FOODS, LLC (056098817)

Revised: 4/2023

WINCO FOODS, LLC