

CHILDRENS ACETAMINOPHEN- acetaminophen tablet, chewable
United Natural Foods, Inc. dba UNFI

Equaline 44-449-children

Active ingredient (in each chewable tablet)

Acetaminophen 160 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - sore throat
 - flu
 - toothache
 - the common cold
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this

product

Ask a doctor before use if your child has
liver disease.

Ask a doctor or pharmacist before use if your child is
taking the blood thinning drug warfarin.

When using this product
do not exceed recommended dosage.

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed**
- find the right dose on chart below. If possible, use weight to dose; otherwise, use age.
- chew or crush tablets completely before swallowing; do not swallow tablets whole
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- do not give for more than 5 days unless directed by a doctor

Weight (lb)	Age (yr)	Dose (chewable tablets)*
Under 24	Under 2	Ask a doctor
24-35	2-3	1 tablet
36-47	4-5	1 1/2 tablets
48-59	6-8	2 tablets
60-71	9-10	2 1/2 tablets
72-95	11	3 tablets

*or as directed by a doctor

Other information

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- avoid high humidity
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

citric acid, crospovidone, D&C red #27 aluminum lake, D&C red #30 aluminum lake, dextrates hydrated, ethylcellulose, FD&C blue #1 aluminum lake, flavors, magnesium stearate, mannitol, polyethylene, stearic acid, sucralose

Questions or comments?

1-855-423-2630

Principal Display Panel

EQUALINE®

**compare to
Children's Tylenol®**

active ingredient**
NDC 41163-494-08

children's
acetaminophen
160 mg, chewable tablets
pain reliever/fever reducer

for ages 2-11

grape flavor

- *aspirin free*
- *ibuprofen free*

CHEW OR CRUSH TABLETS COMPLETELY BEFORE SWALLOWING.
DO NOT SWALLOW TABLETS WHOLE.

24 chewable tablets

actual size

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS
OPENED OR IF BLISTER UNIT IS TORN, BROKEN
OR SHOWS ANY SIGNS OF TAMPERING**

**100% Quality
GUARANTEED**

Like it or let us make it right.

That's our quality promise.

855-423-2630

supervaluprivatebrands.com

DISTRIBUTED BY UNFI

PROVIDENCE, RI 02908 USA

**This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Children's Tylenol®.

50844 REV0522D44908

EQUALINE®

children's
acetaminophen
160 mg, chewable tablets

pain reliever/fever reducer

24 chewable tablets

EQUALINE®

children's
acetaminophen
160 mg, chewable tablets

pain reliever/fever reducer

for ages 2-11
grape flavor

- aspirin free
- ibuprofen free

CHEW OR CRUSH TABLETS COMPLETELY BEFORE SWALLOWING.
DO NOT SWALLOW TABLETS WHOLE.

24 chewable tablets

compare to
Children's Tylenol®
active ingredient**
NDC 41163-494-08



actual size

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS
OPENED OR IF BLISTER UNIT IS TORN, BROKEN
OR SHOWS ANY SIGNS OF TAMPERING



**100% Quality
GUARANTEED**

Like it or let us make it right.
That's our quality promise.
855-423-2630

DISTRIBUTED BY UNFI

PROVIDENCE, RI 02908 USA

**This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Children's Tylenol®.
50844 REV0522D44908

Drug Facts (continued)
Inactive ingredients citric acid, croscopolone, D&C red #27 aluminum lake, D&C red #30 aluminum lake, dextrates/hydrated, ethylcellulose, FD&C blue #1 aluminum lake, flavors, magnesium stearate, mannitol, polyethylene, stearic acid, sucralose
Questions or comments? 1-855-423-2630

No print/No varnish
Lot & Exp date

B-0019E-449-08-RR
REV0522D44908

Drug Facts

KEEP OUTER PACKAGE
FOR COMPLETE
PRODUCT INFORMATION

Active ingredient (in each chewable tablet)
Acetaminophen 160 mg Pain reliever/
fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - sore throat
 - flu
 - toothache
 - the common cold
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes more than 5 doses in 24 hours, which is the maximum daily amount. If you give your child more than 5 doses in 24 hours, which is the maximum daily amount, or if your child has ever taken acetaminophen with other drugs containing acetaminophen, severe liver damage may occur. Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin redness
- hives
- rash

 If a skin reaction occurs, stop use and seek medical help right away.
 Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Drug Facts (continued)

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if your child has

- liver disease.
- ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
 - fever gets worse or lasts more than 3 days
 - new symptoms occur
 - redness or swelling is present
- These could be signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- find the right dose on chart below. If possible, use weight to dose; otherwise, use age.
- chew or crush tablets completely before swallowing; do not swallow tablets whole
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- do not give for more than 5 days unless directed by a doctor

Weight (lb)	Age (yr)	Dose (chewable tablets)*
Under 24	Under 2	Ask a doctor
24-35	2-3	1 tablet
36-47	4-5	1 1/2 tablets
48-59	6-8	2 tablets
60-71	9-10	2 1/2 tablets
72-85	11	3 tablets

*or as directed by a doctor

Other information

TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN ■ avoid high humidity ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ see end flap for expiration date and lot number

CHILDRENS ACETAMINOPHEN

acetaminophen tablet, chewable

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41163-494
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561)	
D&C RED NO. 27 ALUMINUM LAKE (UNII: ZK64F7XSTX)	
D&C RED NO. 30 (UNII: 2S42T2808B)	
DEXTROSE MONOHYDRATE (UNII: LX22YL083G)	
ETHYLCELLULOSE, UNSPECIFIED (UNII: 7Z8S9VYZ4B)	
FD&C BLUE NO. 1 ALUMINUM LAKE (UNII: J9EQA3S2JM)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MANNITOL (UNII: 3OWL53L36A)	
HIGH DENSITY POLYETHYLENE (UNII: UG00KM4WR7)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color	purple	Score	2 pieces
Shape	ROUND	Size	16mm
Flavor	GRAPE	Imprint Code	44;449
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41163-494-08	4 in 1 CARTON	01/28/2005	
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	01/28/2005	

Labeler - United Natural Foods, Inc. dba UNFI (943556183)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(41163-494)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(41163-494) , pack(41163-494)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(41163-494)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(41163-494)

Revised: 6/2023 United Natural Foods, Inc. dba UNFI