

**POLYETHYLENE GLYCOL 3350- polyethylene glycol 3350 powder, for solution**  
**Geri-Care Pharmaceutical Corp**

-----  
**gc peg 489**

**Active ingredient (in each dose)**

Polyethylene Glycol 3350, 17 grams (cap filled to the top)

**Purpose**

Osmotic Laxative

**Uses**

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

**Warnings**

**Allergy alert:** Do not use if you are allergic to polyethylene glycol

**Do not use** if you have kidney disease, except under the advice and supervision of a doctor

**Ask a doctor before use if you have**

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug

**When using this product** you may have loose, watery, more frequent stools

**Stop use and ask a doctor if**

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

**If pregnant or breast feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- do not take more than directed unless advised by your doctor

- the bottle top is a measuring cup marked to contain 17 grams of powder when filled to the indicated “17 GRAMS” line
- adults and children 17 years of age and older:
- use once a day
- fill to indicated “17 GRAMS” line in cap which is marked to indicate the correct dose 17 grams
- stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
- do not combine with starch-based thickeners used for difficulty swallowing
- ensure that the powder is fully dissolved before drinking
- do not drink if there are any clumps
- do not use more than 7 days
- children 16 years of age or under: ask a doctor

### Other information

- store at 20° - 25°C (68° - 77°F)
- tamper-evident: do not use if foil seal under cap printed with “SEALED for YOUR PROTECTION” is missing, open or broken.



### Inactive ingredients

none

### Questions or comments?

1-800-540-3765

### Package label

 <p>LIVE LIFE WELL</p> <p>NDC 57896-489-30</p> <p><b>Polyethylene Glycol 3350 Powder for Solution</b> <b>Osmotic Laxative</b></p> <ul style="list-style-type: none"> <li>• Relieves Occasional Constipation (Irregularity)</li> <li>• Softens Stool</li> </ul> <p>Unflavored Powder • Sugar Free</p> <p>Compare to the active ingredient in MiraLAX®*</p> <p>Net Wt. 17.9 OZ (510 g) 30 ONCE-DAILY DOSES</p>	<p><i>Dissolves in Any Beverage</i></p> <p>*This product is not manufactured or distributed by the owner of the registered trademark MiraLAX®.</p> <p>Manufactured for: Geri-Care Pharmaceuticals 1295 Towbin Ave., Lakewood, NJ 08701 7116-0510-S4-BL Rev 1/2022</p>  <p>Lot No.: Exp. Date:</p>	<p><b>Polyethylene Glycol 3350 increases frequency of bowel movements and softens the stool.</b></p> <table border="1"> <tr> <td data-bbox="1023 1344 1258 1459"> <p><b>Drug Facts</b></p> <p><b>Active ingredient (in each dose)</b> Polyethylene Glycol 3350, 17 grams (cap filled to the top).....Osmotic Laxative</p> </td> <td data-bbox="1258 1344 1502 1459"> <p><b>Purpose</b> relieves occasional constipation (irregularity). generally produces a bowel movement in 1 to 3 days</p> </td> </tr> <tr> <td data-bbox="1023 1459 1258 1575"> <p><b>Use</b> relieves occasional constipation (irregularity). generally produces a bowel movement in 1 to 3 days</p> </td> <td data-bbox="1258 1459 1502 1575"> <p><b>Drug Facts (continued)</b> Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> </td> </tr> <tr> <td data-bbox="1023 1575 1258 1690"> <p><b>Warnings</b> Allergy alert: Do not use if you are allergic to polyethylene glycol. Do not use if you have kidney disease, except under the advice and supervision of a doctor. Ask a doctor before use if you have: nausea, vomiting or abdominal pain a sudden change in bowel habits that lasts over 2 weeks irritable bowel syndrome</p> </td> <td data-bbox="1258 1575 1502 1690"> <p><b>Directions</b> do not take more than directed unless advised by your doctor the bottle cap is a measuring cup designed to contain 17 grams of powder when filled to the top rim adults and children 17 years of age and older: use once a day fill to the top of the bottle cap which will provide the correct dose (17g) stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink do not combine with starch-based thickeners used for difficulty swallowing ensure that the powder is fully dissolved before drinking do not drink if there are any clumps do not use more than 7 days children 16 years of age or under: ask a doctor</p> </td> </tr> <tr> <td data-bbox="1023 1690 1258 1827"> <p><b>When using this product</b> you may have loose, watery, more frequent stools. <b>Stop use and ask a doctor if</b> you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition. you get diarrhea you need to use a laxative for longer than 1 week If pregnant or breast-feeding, ask a health professional before use.</p> </td> <td data-bbox="1258 1690 1502 1827"> <p><b>Other information</b> store at 20° - 25°C (68° - 77°F) tamper-evident: do not use if foil seal under cap, printed with “SEALED for YOUR PROTECTION” is missing, open or broken</p> <p><b>Inactive ingredients</b> none</p> <p><b>Questions or comments?</b> 1-800-540-3765</p> </td> </tr> </table>	<p><b>Drug Facts</b></p> <p><b>Active ingredient (in each dose)</b> Polyethylene Glycol 3350, 17 grams (cap filled to the top).....Osmotic Laxative</p>	<p><b>Purpose</b> relieves occasional constipation (irregularity). generally produces a bowel movement in 1 to 3 days</p>	<p><b>Use</b> relieves occasional constipation (irregularity). generally produces a bowel movement in 1 to 3 days</p>	<p><b>Drug Facts (continued)</b> Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>	<p><b>Warnings</b> Allergy alert: Do not use if you are allergic to polyethylene glycol. Do not use if you have kidney disease, except under the advice and supervision of a doctor. Ask a doctor before use if you have: nausea, vomiting or abdominal pain a sudden change in bowel habits that lasts over 2 weeks irritable bowel syndrome</p>	<p><b>Directions</b> do not take more than directed unless advised by your doctor the bottle cap is a measuring cup designed to contain 17 grams of powder when filled to the top rim adults and children 17 years of age and older: use once a day fill to the top of the bottle cap which will provide the correct dose (17g) stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink do not combine with starch-based thickeners used for difficulty swallowing ensure that the powder is fully dissolved before drinking do not drink if there are any clumps do not use more than 7 days children 16 years of age or under: ask a doctor</p>	<p><b>When using this product</b> you may have loose, watery, more frequent stools. <b>Stop use and ask a doctor if</b> you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition. you get diarrhea you need to use a laxative for longer than 1 week If pregnant or breast-feeding, ask a health professional before use.</p>	<p><b>Other information</b> store at 20° - 25°C (68° - 77°F) tamper-evident: do not use if foil seal under cap, printed with “SEALED for YOUR PROTECTION” is missing, open or broken</p> <p><b>Inactive ingredients</b> none</p> <p><b>Questions or comments?</b> 1-800-540-3765</p>
<p><b>Drug Facts</b></p> <p><b>Active ingredient (in each dose)</b> Polyethylene Glycol 3350, 17 grams (cap filled to the top).....Osmotic Laxative</p>	<p><b>Purpose</b> relieves occasional constipation (irregularity). generally produces a bowel movement in 1 to 3 days</p>									
<p><b>Use</b> relieves occasional constipation (irregularity). generally produces a bowel movement in 1 to 3 days</p>	<p><b>Drug Facts (continued)</b> Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>									
<p><b>Warnings</b> Allergy alert: Do not use if you are allergic to polyethylene glycol. Do not use if you have kidney disease, except under the advice and supervision of a doctor. Ask a doctor before use if you have: nausea, vomiting or abdominal pain a sudden change in bowel habits that lasts over 2 weeks irritable bowel syndrome</p>	<p><b>Directions</b> do not take more than directed unless advised by your doctor the bottle cap is a measuring cup designed to contain 17 grams of powder when filled to the top rim adults and children 17 years of age and older: use once a day fill to the top of the bottle cap which will provide the correct dose (17g) stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink do not combine with starch-based thickeners used for difficulty swallowing ensure that the powder is fully dissolved before drinking do not drink if there are any clumps do not use more than 7 days children 16 years of age or under: ask a doctor</p>									
<p><b>When using this product</b> you may have loose, watery, more frequent stools. <b>Stop use and ask a doctor if</b> you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition. you get diarrhea you need to use a laxative for longer than 1 week If pregnant or breast-feeding, ask a health professional before use.</p>	<p><b>Other information</b> store at 20° - 25°C (68° - 77°F) tamper-evident: do not use if foil seal under cap, printed with “SEALED for YOUR PROTECTION” is missing, open or broken</p> <p><b>Inactive ingredients</b> none</p> <p><b>Questions or comments?</b> 1-800-540-3765</p>									

## POLYETHYLENE GLYCOL 3350

polyethylene glycol 3350 powder, for solution

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:57896-489
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>POLYETHYLENE GLYCOL 3350</b> (UNII: G2M7P15E5P) (POLYETHYLENE GLYCOL 3350 - UNII:G2M7P15E5P)	POLYETHYLENE GLYCOL 3350	17 g in 17 g

**Product Characteristics**

<b>Color</b>	white	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:57896-489-14	238 g in 1 BOTTLE; Type 0: Not a Combination Product	07/01/2016	
2	NDC:57896-489-30	510 g in 1 BOTTLE; Type 0: Not a Combination Product	07/01/2016	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090812	05/01/2014	

**Labeler** - Geri-Care Pharmaceutical Corp (611196254)**Registrant** - Geri-Care Pharmaceutical Corp (611196254)