

LAXATIVE- bisacodyl tablet, delayed release
CHAIN DRUG MARKETING ASSOCIATION INC

Quality Choice 44-327

Active ingredient (in each tablet)

Bisacodyl USP, 5 mg

Purpose

Stimulant laxative

Uses

- for relief of occasional constipation and irregularity
- this product generally produces bowel movement in 6 to 12 hours

Warnings

Do not use

if you cannot swallow without chewing.

Ask a doctor before use if you have

- a sudden change in bowel habits that lasts more than 2 weeks
- stomach pain, nausea or vomiting

When using this product

- do not use within 1 hour after taking an antacid or milk
- do not chew or crush tablet(s)
- you may have stomach discomfort, faintness and cramps

Stop use and ask a doctor if

- you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- take with a glass of water

adults and children 12 years and over	take 1 to 3 tablets in a single daily dose
children 6 to under 12 years	take 1 tablet in a single daily dose
children under 6 years	ask a doctor

Other information

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- avoid excessive humidity
- see end flap for expiration date and lot number

Inactive ingredients

acacia, ammonium hydroxide, calcium carbonate, carnauba wax, colloidal anhydrous silica, corn starch, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, iron oxide black, lactose anhydrous, magnesium stearate, methylparaben, polydextrose, polyethylene glycol, polyvinyl acetate phthalate, povidone, propylene glycol, propylparaben, shellac glaze, simethicone, sodium alginate, sodium benzoate, sodium bicarbonate, stearic acid, sucrose, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments?

1-800-426-9391

Principal Display Panel

QC®

QUALITY
CHOICE

NDC 63868-572-25

*Compare to the
Active Ingredient in
Dulcolax® Laxative Tablets

Laxative

Bisacodyl USP, 5 mg - Stimulant Laxative

**Gentle, Dependable
Constipation Relief**

Actual Size

25 Comfort Coated Tablets

**TAMPER EVIDENT: DO NOT USE IF
PACKAGE IS OPENED OR IF BLISTER
UNIT IS TORN, BROKEN OR SHOWS
ANY SIGNS OF TAMPERING**

*This product is not manufactured or distributed by Sanofi-Aventis Deutschland GMBH,
owner of the registered trademark
Dulcolax® Laxative Tablets. 50844 REV0119A32756

SATISFACTION GUARANTEED
100%QC

Distributed by C.D.M.A., Inc. ©
43157 W 9 Mile Rd
Novi, MI 48375
www.qualitychoice.com
Questions: 800-935-2362



Overnight Relief

Laxative

Bisacodyl USP, 5 mg - Stimulant Laxative

NDC 63868-572-25



*Compare to the
Active Ingredient in
Dulcolax® Laxative Tablets

Laxative

Bisacodyl USP, 5 mg - Stimulant Laxative

Gentle, Dependable
Constipation Relief

Actual
Size



25 Comfort Coated Tablets

E1020R

This product is not manufactured or distributed by Sandoz-Aventis Deutschland GmbH, owner of the registered trademark Dulcolax® Laxative Tablets. 50844 REV0119A32756

Drug Facts (continued)

sodium benzoate, sodium bicarbonate, stearic acid, sucrose, talc, titanium dioxide, triethyl citrate, triethyl citrate

Drug Facts (continued)

Questions or comments?
1-800-426-9391

B-0220-327-56-R
REV0119A32756




Distributed by C.D.M.A., Inc.®
43157 W 9 Mile Rd
Novi, MI 48375
www.qualitychoice.com
Questions: 800-935-2362

OR BROKEN
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
■ avoid excessive humidity
■ see end flap for expiration date and lot number
Inactive ingredients acacia, ammonium hydroxide, calcium carbonate, carnauba wax, colloidal anhydrous silica, corn starch, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, iron oxide black, lactose anhydrous, magnesium stearate, methylparaben, polydextrose, polyethylene glycol, polyvinyl acetate phthalate, povidone, propylene glycol, propylparaben, shellac glaze, simethicone, sodium alginate.

■ a sudden change in bowel habits that lasts more than 2 weeks
■ When using this product
■ do not chew or crush tablet(s)
■ do not use within 1 hour after taking an antacid or milk
■ you may have stomach discomfort, flatulence and cramps
■ Stop use and ask a doctor if
■ you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition.
■ you need to use a laxative for more than 1 week
■ If pregnant or breast-feeding, ask a health professional before use.



5 11896 5755E 9



Quality Choice 44-327

LAXATIVE

bisacodyl tablet, delayed release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63868-572
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISACODYL (UNII: 10X0709Y6I) (DEACETYLBISACODYL - UNII:R09078E41Y)	BISACODYL	5 mg

Inactive Ingredients

Ingredient Name	Strength
ACACIA (UNII: 5C5403N26O)	
AMMONIA (UNII: 5138Q19F1X)	
CALCIUM CARBONATE (UNII: H0G9379FGK)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
FERROSO FERRIC OXIDE (UNII: XM0M87F357)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
METHYL PARABEN (UNII: A2I8C7HI9T)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
Polyvinyl Acetate Phthalate (UNII: 58QVG85GW3)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

PROPYLPARABEN (UNII: Z8IX2SC1OH)	
SHELLAC (UNII: 46N107B71O)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
SODIUM ALGINATE (UNII: C269C4G2ZQ)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCROSE (UNII: C151H8M554)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

Product Characteristics

Color	orange	Score	no score
Shape	ROUND	Size	6mm
Flavor		Imprint Code	5
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63868-572-25	1 in 1 CARTON	02/14/2020	
1		25 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	505G(a)(3)	02/14/2020	

Labeler - CHAIN DRUG MARKETING ASSOCIATION INC (011920774)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	manufacture(63868-572) , pack(63868-572)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(63868-572) , pack(63868-572)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(63868-572)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(63868-572)