

IBUPROFEN- ibuprofen tablet, film coated
IBUPROFEN- ibuprofen tablet, film coated
TIME CAP LABORATORIES,INC

IBUPROFEN TABLETS USP 200MG ROUND BROWN

DRUG FACTS

Active ingredient (in each tablet)

Ibuprofen 200 mg (NSAID)*

* nonstreoidal anti-inflammatory drug

Purpose

Pain reliever / Fever reducer

Uses

temporarily relieves minor aches and pains due to:

- o headache
- o toothache
- o backache
- o menstrual cramps
- o the common cold
- o muscular aches
- o minor pain of arthritis

temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood-thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product

- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better

you have symptoms of heart problem or stroke:

- chest pain
- trouble breathing
- weakness in one part or side of the body
- slurred speech
- leg swelling

pain gets worse or lasts more than 10 days

fever gets worse or lasts more than 3 days

redness or swelling is present in the painful area

any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely

directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away

(1-800-222-1222).

Directions

do not take more than directed

the smallest effective dose should be used

adults and children 12 years and over:

- take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor

children under 12 years:

- ask a doctor

OTHER INFORMATION

Other information

- read all warnings and directions before use
- store at 20-25°C (68-77°F)

Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide

Questions or comments? call **1-877-290-4008**

Time-Cap Labs, Inc.
NDC 49483-600-01
*Compare to the active ingredient in Advil® Caplets.

Ibuprofen

Tablets USP, 200 mg

CAPLETS

Pain Reliever/
Fever Reducer (NSAID)

100 FILM-COATED
BROWN CAPLETS
(Capsule-Shaped Tablets)

This product is not manufactured or distributed by Pfizer Consumer Healthcare, owner of the registered trademark Advil® Caplets.

Made in India 600R 0617

Distributed by:

Time-Cap Labs, Inc.

7 Michael Avenue

Farmingdale, NY 11735



49483-600-01 8

TAMPER EVIDENT DO NOT USE THIS PRODUCT IF THE UNPAINTED FOL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts

Active Ingredient (in each caplet)
Ibuprofen 200 mg (NSAID) Pain reliever/fever reducer
nonsteroidal anti-inflammatory drug

Uses ■ temporarily relieves minor aches and pains due to:
■ headache ■ backache ■ menstrual cramps
■ minor pain or arthritis ■ muscular aches
■ the common cold ■ toothache
■ temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
■ asthma (wheezing) ■ hives ■ facial swelling
■ rash ■ shock ■ skin redness ■ blisters ■ If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This drug may cause stomach bleeding, which may cause severe stomach bleeding. The chance is higher if you:
■ have had stomach ulcers or bleeding problems
■ take a blood-thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ have 3 or more alcoholic drinks every day while using this product
Do not use for more than 10 days for pain or for a longer time than directed.
Heart attack and stroke warning: This drug may increase the risk of heart attack, heart failure and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

LOT #:

EXP. DATE

PEEL HERE
TO REVEAL
DRUG
FACTS

[illegible]

Adhesive Area

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if:

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are:

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product:

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if:

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **do not take more than directed**
- the **smallest effective dose should be used**
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- read all warnings and directions before use
- store at 20-25°C (68-77°F)

Inactive ingredients: colloidal silicon dioxide, croscarmellose sodium, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, prepolimerized starch, talc, titanium dioxide

Questions or comments? Call 1-877-290-4008

Inside

1/16"
Margin

NDC 49483-114-73

timely

[†]Compare to the active ingredient in Advil® Tablets

Ibuprofen

TABLETS USP, 200 mg

PAIN RELIEVER / FEVER REDUCER (NSAID)

100 TABLETS

114
Actual Size

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts

Active ingredient (in each tablet)

Ibuprofen 200 mg (NSAID)* Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - the common cold
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

■ skin redness

■ rash

■ blisters

If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Drug Facts (continued under label)

This product is not manufactured or distributed by Haleon group of companies, owner of the registered trademark Advil® Tablets

Distributed by:

Time-Cap Labs, Inc.

7 Michael Avenue,

Farmingdale, NY 11735

Made in India 601R 1123



LOT:
EXP.:

Varnish Omit Area

Adhesive Area

Drug Facts (continued)

Do not use

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver disease, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin
- taking any other drug

When using this product:

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problem or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part of side of the body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless directly directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children
 - take 1 tablet every 4 to 6 hours while symptoms persist
 - if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor

children under 12 years: ask a doctor

Other information ■ read all warnings and directions before use ■ store at 20-25 °C (68-77 °F)

Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, silica, titanium dioxide

Questions or comments? call 1-877-290-4008

Inside

Varnish Omit Area

Lot No.:

PEEL HERE

Exp Date:



Drug Facts (continued under label)

This product is not manufactured or distributed by Pfizer Consumer Healthcare, owner of the registered trademark Advil® Tablets.

Made in India

Distributed by:

Time-Cap Labs, Inc.

7 Michael Avenue

Farmingdale, NY 11735

601R 0223

NDC 49493-114-50

timely

Compare to the Active Ingredient in Advil® Tablets

Ibuprofen

TABLETS USP, 200 mg
PAIN RELIEVER / FEVER REDUCER
(NSAID)

500 TABLETS

Actual Size

114

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts

Active ingredient (in each tablet)

Ibuprofen 200 mg (NSAID) * Pain reliever/fever reducer, nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - backache
 - muscle aches
 - menstrual cramps
- minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling (wheezing)
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood-thinning (anticoagulant) or steroid drug
- take other drugs containing aspirin or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> ■ You have problems or serious side effects from taking pain relievers or fever reducers ■ You have a history of stomach problems, such as heartburn ■ You have high blood pressure, liver disease, heart problems, kidney disease, asthma, or had a stroke ■ You are taking a diuretic <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> ■ under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease its benefit of aspirin ■ taking any other drug <p>When using this product:</p> <ul style="list-style-type: none"> ■ take with food or milk if your stomach upset occurs <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ you have stomach pain that does not get better ■ you have symptoms of heart problem or stroke: ■ chest pain trouble breathing ■ weakness in one part or side of body ■ slurred speech leg swelling ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear <hr/> <p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy, unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact Poison Control Center right away. (1-800-222-1222).</p> <p>Directions</p> <ul style="list-style-type: none"> ■ do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and over: take 1 tablet every 4 to 6 hours with symptoms persist ■ if pain or fever does not respond to 1 tablet, 2 tablets may be used ■ do not exceed 6 tablets in 24 hours, unless directed by a doctor ■ Children under 12 years: ask a doctor <p>Other information</p> <ul style="list-style-type: none"> ■ read all warnings and directions before use ■ store at 20°-25°C (68°-77°F) <p>Inactive ingredients</p> <p>croscarmellose sodium, croscarmellose sodium, iron oxide red, polyethylene glycol, hydroxypropyl methylcellulose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide.</p>	<p>Questions or comments? Call 1-877-290-4008</p>
--	--

NDC 49483-114-17

[†]Compare to the active ingredient in Advil® Tablets

Ibuprofen

TABLETS USP, 200 mg

PAIN RELIEVER / FEVER REDUCER (NSAID)

114

200

TABLETS

Actual Size

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts	Purpose
<p>Active ingredient (in each tablet) Ibuprofen 200 mg (NSAID) <small>*nonsteroidal anti-inflammatory drug</small></p>	<p>Pain relief/fever reducer</p>
<p>Uses</p> <ul style="list-style-type: none"> ■ temporarily relieves minor aches and pains due to: <ul style="list-style-type: none"> ■ headache ■ toothache ■ backache ■ menstrual cramps ■ the common cold ■ muscular aches ■ minor pain of arthritis ■ temporarily reduces fever 	
<p>Warnings</p> <p>Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:</p> <ul style="list-style-type: none"> ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin redness ■ rash ■ blisters <p>If an allergic reaction occurs, stop use and seek medical help right away.</p> <p>Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:</p> <ul style="list-style-type: none"> ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood-thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed <p>Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.</p>	
<p>Do not use</p> <ul style="list-style-type: none"> ■ if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery 	
<p>Ask a doctor before use if</p> <ul style="list-style-type: none"> ■ stomach bleeding warning applies to you 	

Drug Facts (continued under label)

This product is not manufactured or distributed by Haleon group of companies, owner of the registered trademark Advil® Tablets

Distributed by:

Time-Cap Labs, Inc.

7 Michael Avenue

Farmingdale, NY 11735

Made in India 601R 0723

LOT No.: **EXP. DATE**

NDC 49483-114-17

[†]Compare to the active ingredient in Advil® Tablets

Ibuprofen

TABLETS USP, 200 mg

PAIN RELIEVER / FEVER REDUCER (NSAID)

114

200

TABLETS

Actual Size

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts	Purpose
<p>Active ingredient (in each tablet) Ibuprofen 200 mg (NSAID) <small>*nonsteroidal anti-inflammatory drug</small></p>	<p>Pain relief/fever reducer</p>
<p>Uses</p> <ul style="list-style-type: none"> ■ temporarily relieves minor aches and pains due to: <ul style="list-style-type: none"> ■ headache ■ toothache ■ backache ■ menstrual cramps ■ the common cold ■ muscular aches ■ minor pain of arthritis ■ temporarily reduces fever 	
<p>Warnings</p> <p>Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:</p> <ul style="list-style-type: none"> ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin redness ■ rash ■ blisters <p>If an allergic reaction occurs, stop use and seek medical help right away.</p> <p>Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:</p> <ul style="list-style-type: none"> ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood-thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed <p>Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.</p>	
<p>Do not use</p> <ul style="list-style-type: none"> ■ if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery 	
<p>Ask a doctor before use if</p> <ul style="list-style-type: none"> ■ stomach bleeding warning applies to you 	

Drug Facts (continued under label)

This product is not manufactured or distributed by Haleon group of companies, owner of the registered trademark Advil® Tablets

Distributed by:


Time-Cap Labs, Inc.

7 Michael Avenue

Farmingdale, NY 11735

Made in India 601R 0723

LOT No.: **EXP. DATE**

<p>NDC 49483-114-10</p> <p>timely™ Compare to the Active Ingredient in Advil® Tablets</p> <h1>Ibuprofen</h1> <p>TABLETS USP, 200 mg</p> <p>PAIN RELIEVER / FEVER REDUCER (NSAID)</p> <p>1000 TABLETS</p>  <p>Actual Size</p>		<p>DANGER WARNING: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING</p> <table border="1"> <tr> <th>Drug Facts</th> <th>Drug Facts (continued)</th> </tr> <tr> <td> Active ingredient (in each tablet) Purpose Ibuprofen 200 mg (NSAID) ... Pain reliever/Fever reducer <i>*nonsteroidal anti-inflammatory drug</i> </td> <td> Direct or for longer than directed. Do not use ■ if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic Ask a doctor or pharmacist before use if you are ■ under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug When using this product ■ take with food or milk if stomach upset occurs Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problem or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of the body ■ slurred speech ■ leg swelling ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days </td> </tr> <tr> <th>Uses</th> <th>Directions</th> </tr> <tr> <td> ■ temporarily relieves minor aches and pains due to: ■ headache ■ toothache ■ backache ■ menstrual cramps ■ the common cold ■ muscular aches ■ minor pain of arthritis ■ temporarily reduces fever </td> <td> ■ redness or swelling is present in the painful area ■ any new symptoms appear If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). do not take more than directed the smallest effective dose should be used </td> </tr> <tr> <th>Warnings</th> <th>Other information</th> </tr> <tr> <td> Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters If an allergic reaction occurs, stop use and seek medical help right away. Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood-thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than </td> <td> Inactive ingredients: colloidal silicon dioxide, croscarmellose sodium, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide Questions or comments? call 1-877-290-4008 </td> </tr> </table>		Drug Facts	Drug Facts (continued)	Active ingredient (in each tablet) Purpose Ibuprofen 200 mg (NSAID) ... Pain reliever/Fever reducer <i>*nonsteroidal anti-inflammatory drug</i>	Direct or for longer than directed. Do not use ■ if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic Ask a doctor or pharmacist before use if you are ■ under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug When using this product ■ take with food or milk if stomach upset occurs Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problem or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of the body ■ slurred speech ■ leg swelling ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days	Uses	Directions	■ temporarily relieves minor aches and pains due to: ■ headache ■ toothache ■ backache ■ menstrual cramps ■ the common cold ■ muscular aches ■ minor pain of arthritis ■ temporarily reduces fever	■ redness or swelling is present in the painful area ■ any new symptoms appear If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). do not take more than directed the smallest effective dose should be used	Warnings	Other information	Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters If an allergic reaction occurs, stop use and seek medical help right away. Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood-thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than	Inactive ingredients: colloidal silicon dioxide, croscarmellose sodium, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide Questions or comments? call 1-877-290-4008	<p>This product is not manufactured or distributed by Pfizer Consumer Healthcare, owner of the Imprint registered trademark Advil® Tablets. 601R 0522</p> <p>Made in India Distributed by: Time-Cap Labs, Inc., 7 Michael Avenue Paramus, NY 11765 Lot No.: C-04-N-00005555</p> <p>Varnish Omit Area</p> <p>Exp Date:</p>
Drug Facts	Drug Facts (continued)															
Active ingredient (in each tablet) Purpose Ibuprofen 200 mg (NSAID) ... Pain reliever/Fever reducer <i>*nonsteroidal anti-inflammatory drug</i>	Direct or for longer than directed. Do not use ■ if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic Ask a doctor or pharmacist before use if you are ■ under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug When using this product ■ take with food or milk if stomach upset occurs Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problem or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of the body ■ slurred speech ■ leg swelling ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days															
Uses	Directions															
■ temporarily relieves minor aches and pains due to: ■ headache ■ toothache ■ backache ■ menstrual cramps ■ the common cold ■ muscular aches ■ minor pain of arthritis ■ temporarily reduces fever	■ redness or swelling is present in the painful area ■ any new symptoms appear If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). do not take more than directed the smallest effective dose should be used															
Warnings	Other information															
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters If an allergic reaction occurs, stop use and seek medical help right away. Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood-thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than	Inactive ingredients: colloidal silicon dioxide, croscarmellose sodium, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide Questions or comments? call 1-877-290-4008															

ibuprofen tablet, film coated

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49483-114	
Route of Administration	ORAL			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)		IBUPROFEN	200 mg	
Inactive Ingredients				
Ingredient Name			Strength	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)				
TALC (UNII: 7SEV7J4R1U)				
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
FERRIC OXIDE RED (UNII: 1K09F3G675)				
STARCH, CORN (UNII: O8232NY3SJ)				
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
Product Characteristics				
Color	brown	Score	no score	
Shape	ROUND (Biconvex)	Size	10mm	
Flavor		Imprint Code	114	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49483-114-50	500 in 1 BOTTLE; Type 0: Not a Combination Product	09/09/2021	
2	NDC:49483-114-10	1000 in 1 BOTTLE; Type 0: Not a Combination Product	09/09/2021	
3	NDC:49483-114-17	200 in 1 BOTTLE; Type 0: Not a Combination Product	08/19/2023	
4	NDC:49483-114-73	100 in 1 BOTTLE; Type 0: Not a Combination Product	11/18/2023	
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA		ANDA091239	03/22/2016	

IBUPROFEN

ibuprofen tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49483-601
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL (UNII: 532B59J990)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	

Product Characteristics

Color	brown	Score	2 pieces
Shape	ROUND	Size	10mm
Flavor		Imprint Code	114
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49483-601-00	6500 in 1 BAG; Type 0: Not a Combination Product	03/22/2016	
2	NDC:49483-601-05	50 in 1 BOTTLE; Type 0: Not a Combination Product	03/22/2016	
3	NDC:49483-601-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	03/22/2016	
4	NDC:49483-601-50	500 in 1 BOTTLE; Type 0: Not a Combination Product	03/22/2016	08/07/2021
5	NDC:49483-601-10	1000 in 1 BOTTLE; Type 0: Not a Combination Product	03/22/2016	08/07/2021

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091239	03/22/2016	

IBUPROFEN

ibuprofen tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49483-600
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
POLYVINYL ALCOHOL (UNII: 532B59J990)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
TALC (UNII: 7SEV7J4R1U)	

Product Characteristics

Color	brown	Score	2 pieces
Shape	CAPSULE	Size	15mm
Flavor		Imprint Code	117
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49483-600-05	50 in 1 BOTTLE; Type 0: Not a Combination Product	03/22/2016	
2	NDC:49483-600-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	03/22/2016	

3	NDC:49483-600-50	500 in 1 BOTTLE; Type 0: Not a Combination Product	11/14/2017	
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
ANDA	ANDA091239		03/22/2016	

Labeler - TIME CAP LABORATORIES,INC (037052099)

Registrant - TIME CAP LABORATORIES,INC (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LIMITED		925822975	manufacture(49483-601, 49483-600, 49483-114)

Revised: 11/2023

TIME CAP LABORATORIES,INC