ASPIRIN ENTERIC SAFETY COATED- aspirin tablet Publix Supermarkets, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredient (in each tablet)
Aspirin 81 mg (NSAID)*
*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

- for the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.
- ask your doctor about other uses for safety coated 81 mg aspirin

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- ringing in the ears or loss of hearing occurs
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor

children under 12 years: consult a doctor

Other information

store between 59-86°F (15-30°C)

Inactive ingredients

D&C yellow #10, FD&C yellow #6**, methacrylic acid copolymer, microcrystalline cellulose, pregelatinized starch corn, silicon dioxide, sodium bicarbonate, sodium lauryl sulfate, stearic acid, talc, titanium dioxide, triethyl citrate

Principal Display Panel

†Compare to the active ingredient in Bayer® Low Dose Aspirin

LOW DOSE

Aspirin

ASPIRIN REGIMEN 81 mg≠

PAIN RELIEVER (NSAID)

Safety coated≠≠

TABLETS

†This product is not manufactured or distributed by Bayer Healthcare LLC, distributor of Bayer® Low Dose Aspirin

≠Talk to your doctor or other healthcare provider before using this product for your heart. Aspirin is not right for everyone.

≠≠Coating helps protect against stomach upset

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

DISTRIBUTED BY PUBLIX SUPERMARKETS, INC., 3300 PUBLIX CORPORATE PARKWAY, LAKELAND, FL 33811 1-888-267-3037 publix.com

Product Label

Drug Facts (continued)

- ablets every 4 hours not to exceed 48 tablets in 24 drink a full glass of water with each dose
 adults and children 12 years and over: take 4 to 8 hours unless directed by a doctor
 - children under 12 years: consult a doctor

Other information

■ store between 59-86°F (15-30°C)

Inactive ingredients D&C yellow #10, FD&C yellow #6, methacrylic acid copolymer

microcrystalline cellulose, pregelatinized starch (com), silicon dioxide, sodium bicarbonate, sodium lauryl sulfate stearic acid, talc, titanium dioxide, triethyl citrate †This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Bayer® Low Dose Aspirin. *Talk to your doctor or other healthcare provider before using this product for your heart. Aspirin is not right for everyone

#Coating helps protect against stomach upset

AMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

LOW DOSE

ASPIRIN REGIMEN 81 mg[‡]

ask your doctor about other uses for safety coated

81 mg aspirin

needing immediate relief.

PAIN RELIEVER (NSAID)

Safety coated^{‡‡}

120

PUBLIX GUARANTEE: PUBLIX GUARANTEE: COMPLETE SATISFACTION OR YOUR MONEY BACK

PLD-B567A

FC005623

Exp. Date: Lot No.:

DISTRIBUTED BY PUBLIX SUPER MARKETS, INC.,

LAKELAND, FL 33811 1-888-267-3037

publix.com

3300 PUBLIX CORPORATE PARKWAY

ACTUAL SIZE

[†]Compare to the active ingredient in Bayer® Low Dose Aspirin

take other drugs containing prescription or nonprescription have 3 or more alcoholic drinks every day while using this

NSAIDs (aspirin, ibuprofen, naproxen, or others)

take more or for a longer time than directed

product

take a blood thinning (anticoagulant) or steroid drug

have had stomach ulcers or bleeding problems

chance is higher if you:

are age 60 or older

Drug Facts

Orug Facts (continued)

Do not use

Active ingredient (in each tablet)

oidal anti-inflammatory drug Spirin 81 mg (NSA)

ain reliever

if you are allergic to aspirin or any other pain reliever/fever if you have ever had an allergic reaction to this product or Purpose

Ask a doctor or pharmacist before use if you are taking a ■ you are taking a diuretic
■ you have asthma

or kidney disease

 you have a history of stomach problems, such as heartburn you have high blood pressure, heart disease, liver cirrhosis

stomach bleeding warning applies to you

Ask a doctor before use if

any of its ingredients

 an allergic reaction occurs. Seek medical help right away prescription drug for gout diabetes arthritis Stop use and ask a doctor if

 you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood have bloody or black stools

recovering from chicken pox or flu-like symptoms should not

use this product. When using this product, if changes in

Reye's syndrome: Children and teenagers who have or are

have stomach pain that does not get bette pain gets worse or lasts more than 10 days

fever gets worse or lasts more than 3 days

Illergy alert: Aspirin may cause a severe allergic reaction because these symptoms could be an early sign of Reye's behavior with nausea and vomiting occur, consult a docto

syndrome, a rare but serious illness

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The

asthma (wheezing)

shock hives

facial swelling

which may include:

 ringing in the ears or loss of hearing occurs redness or swelling is present

These could be signs of a serious condition new symptoms occur

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirir during the last 3 months of pregnancy unless definite problems in the unborn child or complications during directed to do so by a doctor because it may cause

(eep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

NDC 56062-867-12

for the temporary relief of minor aches and pains or as Because of its delayed action, this product will not provide fast relief of headaches or other symptoms

recommended by your docto

PUBLIX Low Dose Aspirin

ASPIRIN ENTERIC SAFETY COATED

aspirin tablet

Product Information	
---------------------	--

Product Type HUMAN OTC DRUG Item Code (Source) NDC:56062-867

Route of Administration ORAL

Active Ingredient/Active Moiety

· · · · · · · · · · · · · · · · · · ·		
Ingredient Name	Basis of Strength	Strength
ACDIDIN (LINII) P16CO5Y76E) (ACDIDIN LINII) P16CO5Y76E)	AC DIDINI	81 ma

Inactive Ingredients

Ingredient Name

Strength

STARCH, CORN (UNII: O8232NY3SJ)

D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)

METHACRYLIC ACID - METHYL METHACRYLATE COPOLYMER (1:1) (UNII: 74G4R6TH13)

MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)

SILICON DIOXIDE (UNII: ETJ7Z6XBU4)

SODIUM BICARBONATE (UNII: 8MDF5V39QO)
SODIUM LAURYL SULFATE (UNII: 368GB5141J)

STEARIC ACID (UNII: 4ELV7Z65AP)

TALC (UNII: 7SEV7J4R1U)

TITANIUM DIOXIDE (UNII: 15FIX9V2JP)
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)

Product Characteristics

Color	yellow	Score	no score
Shape	ROUND	Size	7mm
Flavor		Imprint Code	E
Contains			

Packaging

Ш				
	# Item Code	Package Description	Marketing Start Date	Marketing End Date
	NDC:56062- 867-12	1 in 1 BOX	07/31/2019	
	1	120 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part343	07/31/2019	

Labeler - Publix Supermarkets, Inc. (006922009)

Revised: 12/2022 Publix Supermarkets, Inc.