TRIPROLIDINE HCL DROPS- triprolidine hydrochloride liquid Brandywine Pharmaceuticals, LLC

Triprolidine HCI Drops

Triprolidine HCl Drops NDC 71321-701-50

Active ingredients (in each 2 mL)

Triprolidine HCI 1.25 mg

Purpose

Antihistamine

Uses

- temporarily relieves these symptoms due to hay fever (allergic rhinitis) or other upper respiratory allergies:
- runny nose itching of the nose or throat
- sneezing itchy, watery eyes

Warnings

Do not exceed recommended dosage.

Ask a doctor before use if the child has

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- trouble urinating due to an enlarged prostate

Ask a doctor before use if the child is

taking sedatives or tranquilizers

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur
- sedatives and tranquilizers may increase the drowsiness effect

Stop use and ask a doctor if

■ new symptoms occur

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- use only with enclosed dropper
 mL= milliliter
- do not use dropper for any other drug product

children 6 to under 12 years: 2 mL every 6 hours, not to exceed 4 doses in 24 hours, or as directed by a doctor.

children under 6: ask a doctor.

Other information

- read all product information before using
- this packaging is child-resistant.
- store at room temperature 20-30°C (68-86°F)

Inactive ingredients

citric acid anhydrous, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sodium saccharin, sorbitol solution

Questions or comments?

Call 610-314-7943 9 a.m. - 5 p.m. EST.

PRINCIPAL DISPLAY PANEL

NDC 71321-701-50

Triprolidine HCI Drops

1.69 fl oz (50 mL)

TAMPER EVIDENT: DO NOT USE IF FOIL SEAL UNDER CAP IS BROKEN **OR MISSING**

71321-701-50

Triprolidine HCl **Drops**

Antihistamine

Each dropperful (2 mL), for oral administration, contains: Triprolidine HCl 1.25 mg

Gluten Free • Dye Free Sugar Free • Alcohol Free

1.69 fl oz (50 mL)

RELIEVES:

Sneezing Runny Nose Itchy, Watery Eyes Itchy Nose **Itchy Throat**



Rev 1/2023

TEMPORARILY RELIEVES:

Sneezing **Runny Nose** Itchy, Watery Eyes **Itchy Nose Itchy Throat**

Gluten Free Dye Free Sugar Free **Alcohol Free**



71321-701-50

Triprolidine HCI Drops

Antihistamine

Each dropperful (2 mL), for oral administration, contains: Triprolidine HCl 1.25 mg

Sneezing • Runny Nose Itchy, Watery Eyes Itchy Nose • Itchy Throat

Gluten Free • Dye Free Sugar Free • Alcohol Free



1.69 fl oz (50 mL)

WARNING: KEEP OUT OF REACH OF CHILDREN

TAMPER EVIDENT: DO NOT USE IF FOIL SEAL UNDER CAP IS BROKEN OR MISSING





Drug Facts

(in each 2 mL dropperful)

Warnings
Do not exceed recommended dosage

Ask a doctor before use if the child has

temporarily relieves these symptoms due to hav fever

| Interporary refleves tress symptoms due to may revert
| Interporary refleves tress symptoms due to may revert
| Interporary refleves tress symptoms due to may revert
| Interporary refleves tress symptoms due to may revert
| Interporary refleves tress symptoms due to may revert
| Interporary refleves tress symptoms due to may revert
| Interporary refleves tress symptoms due to may revert
| Interporary refleves tress symptoms due to may revert
| Interporary refleves tress symptoms due to may revert
| Interporary refleves tress symptoms due to may revert
| Interporary refleves tress symptoms due to may revert
| Interporary refleves tress symptoms due to may revert
| Interporary refleves tress symptoms due to may revert
| Interporary refleves tress tres

■ a breathing problem such as emphysema or chronic bronchitis
■ glaucoma ■ trouble urinating due to an enlarged prostate gland

Ask a doctor before use if the child is taking sedatives

Rexcitability may occur, especially in children
 marked drowsiness may occur
 sedatives and tranquilizers may increase the drowsiness effectives.

■ mL= milliliter

2 mL every 6 hours, not to exceed 4 doses in 24 hours, or as directed by a doctor.

Stop use and ask a doctor if ■ new symptoms occur Keep out of reach of children. In case of overdose, get medical

help or contact a Poison Control Center right away.

Dose

■ read all product information before using
■ this packaging is child-resistant
■ store at room temperature 20-30°C (68-86°F)

citric acid anhydrous, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sodium saccharin, sorbitol solution

■ use only with enclosed dropper ■ mL=
■ do not use dropper for any other drug product

12 years: In 24 hours, children under 6: ask a doctor.

Inactive ingredients

Questions or comments? Call 610-314-7943 9 a.m. - 5 p.m. EST

Active ingredients

or tranquilizers
When using this product

Directions

Uses

Manufactured for: Brandywine Pharmaceuticals, LLC West Chester, PA, 19382



Use with enclosed calibrated dropper



Drug Facts

Active ingredients (in each 2 mL dropperful)

Purpose

Triprolidine HCl 1.25 mg.....Antihistamine

Uses

- temporarily relieves these symptoms due to hay fever (allergic rhinitis) or other upper respiratory allergies:
 - runny nose
- itching of the nose or throat
- sneezing
- itchy, watery eyes

Warnings

Do not exceed recommended dosage.

Ask a doctor before use if the child has

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma trouble urinating due to an enlarged prostate gland

Ask a doctor before use if the child is taking sedatives or tranquilizers

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur
- sedatives and tranquilizers may increase the drowsiness effect

Stop use and ask a doctor if new symptoms occur

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- use only with enclosed dropper
- mL= milliliter
- do not use dropper for any other drug product

Age	Dose
Children 6 to under	
12 years:	in 24 hours, or as directed by a doctor.
children under 6:	ask a doctor.

Other information

- read all product information before using
- this packaging is child-resistant.
- store at room temperature 20-30°C (68-86°F)

Inactive ingredients

citric acid anhydrous, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sodium saccharin, sorbitol solution

Questions or comments?

Call 610-314-7943 9 a.m. - 5 p.m. EST

TRIPROLIDINE HCL DROPS

triprolidine hydrochloride liquid

Product Information

HUMAN OTC DRUG NDC:71321-701 **Item Code (Source) Product Type**

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name Basis of Strength Strength TRIPROLIDINE HYDROCHLORIDE (UNII: YAN7R5L890) (TRIPROLIDINE -TRIPROLIDINE 1.25 mg in 2 mL **HYDROCHLORIDE**

UNII:2L8T9S52QM)

SORBITOL (UNII: 506T60A25R)

Inactive Ingredients Ingredient Name Strength ANHYDROUS CITRIC ACID (UNII: XF417D3PSL) GLYCERIN (UNII: PDC6A3C0OX) PROPYLENE GLYCOL (UNII: 6DC9Q167V3) WATER (UNII: 059QF0KO0R) **SODIUM BENZOATE** (UNII: OJ245FE5EU) **SODIUM CITRATE** (UNII: 1Q73Q2|ULR) **SACCHARIN SODIUM** (UNII: SB8ZUX40TY)

Packaging Marketing Start Marketing End # Item Code **Package Description** Date **Date** 1 NDC:71321-50 mL in 1 BOTTLE, DROPPER; Type 0: Not a 03/08/2023 701-50 **Combination Product**

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M012	03/08/2023		

Labeler - Brandywine Pharmaceuticals, LLC (080581956)

Revised: 1/2024 Brandywine Pharmaceuticals, LLC