# DAY COLD AND FLU PLUS AND NIGHTTIME COLD AND FLU PLUS- acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride CVS PHARMACY, INC.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

#### DAY Cold & Flu Plus and NIGHTTIME Cold & Flu Plus

### COLD & FLU FORMULA DAY NON-DROWSY Active ingredients (in each softgel)

Acetaminophen 325 mg Dextromethorphan hydrobromide 10 mg Phenylephrine hydrochloride 5 mg

## COLD & FLU FORMULA NIGHT Active ingredients (in each softgel)

Acetaminophen 325 mg Dextromethorphan hydrobromide 10 mg Doxylamine succinate 6.25 mg Phenylephrine hydrochloride 5 mg

#### **Purposes**

#### **DAY COLD & FLU FORMULA**

Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

#### **Purposes**

#### NIGHT COLD & FLU FORMULA

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Nasal decongestant

#### Uses

#### DAY COLD & FLU FORMULA

- temporarily relieves these symptoms due to a cold or flu:
  - minor aches and pains
  - headache
  - o cough
  - sore throat
  - nasal and sinus congestion

• temporarily reduces fever

#### Uses

#### NIGHT COLD & FLU FORMULA

- temporarily relieves these symptoms due to a cold or flu:
  - minor aches and pains
  - headache
  - nasal and sinus congestion
  - cough
  - sore throat
  - o runny nose
  - o sneezing
- temporarily reduces fever

#### **DAY COLD & FLU FORMULA**

#### **Warnings**

#### **Liver warning**

These products contain acetaminophen. Severe liver damage may occur if you take

- more than 10 softgels in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

#### Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

#### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

#### Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- cough with excessive phlegm (mucus)

- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

#### When using this product do not exceed recommended dosage

#### Stop use and ask a doctor if

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

• nervousness, dizziness, or sleeplessness occurs

**If pregnant or breast-feeding**, ask a health professional before use.

#### Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### NIGHTTIME COLD & FLU FORMULA Warnings

#### Liver warning

These products contain acetaminophen. Severe liver damage may occur if you take

- more than 10 softgels in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

#### Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Allergy alert**: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

#### Do not use to sedate children.

#### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

• in children under 12 years of age

#### Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough with excessive phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

#### Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

#### When using this product

- do not exceed recommended dosage
- may cause marked drowsiness
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

#### Stop use and ask a doctor if

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

• nervousness, dizziness, or sleeplessness occurs

**If pregnant or breast-feeding,** ask a health professional before use.

#### Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### **DAY COLD & FLU FORMULA**

#### Directions

- do not take more than the recommended dose
- adults and children 12 years and over: take 2 softgels with water every 4 hours. Do not exceed 10 softgels in 24 hours or as directed by a doctor.
- children under 12 years: do not use

### NIGHTTIME COLD & FLU FORMULA Directions

- do not take more than the recommended dose
- adults and children 12 years and over: take 2 softgels with water every 4 hours. Do not exceed 10 softgels in 24 hours or as directed by a doctor.
- children under 12 years: do not use

#### DAY COLD & FLU FORMULA

#### Other information

• store at room temperature. Avoid excessive heat.

#### NIGHTTIME COLD & FLU FORMULA

#### Other information

• store at room temperature. Avoid excessive heat.

#### DAY COLD & FLU FORMULA

#### **Inactive ingredients**

FD&C red #40, FD&C yellow #6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol special, and white edible ink

#### NIGHTTIME COLD & FLU FORMULA

#### **Inactive ingredients**

D&C yellow #10, FD&C blue #1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol special, and white edible ink

#### DAY COLD & FLU FORMULA

Questions or comments?

Call toll free: 1-855-215-8180

### NIGHTTIME COLD & FLU FORMULA Questions or comments?

Call toll free: 1-855-215-8180

#### **Principal Display Panel - Carton Label**

CVS NON-DROWSY DAY Cold & Flu Plus and NIGHTTIME Cold & Flu Plus 20ct

NDC 69842-069-08

<sup>\*</sup>Compare to the active ingredients in Alka-Seltzer PLUS® Day and Night Cold and Flu Formula



#### COLD & FLU FORMULA • DAY NON-DROWSY

#### Drug Facts (continued)

Sore throat warning If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

■ skin reddening
■ blisters
■ rash

If a skin reaction occurs, stop use and seek medical help right away.

#### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

#### Ask a doctor before use if you have

- liver disease heart disease high blood pressure
- thyroid disease diabetes
- cough with excessive phlegm (mucus)
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product do not exceed recommended dosage

#### COLD & FLU FORMULA • NIGHT

#### Drug Facts (continued)

#### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

#### Ask a doctor before use if you have

- liver disease
   heart disease
   high blood pressure
- thyroid disease
   diabetes
   glaucoma
- cough with excessive phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

#### Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

#### When using this product

- do not exceed recommended dosage
- may cause marked drowsiness
   avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

#### DAY COLD AND FLU PLUS AND NIGHTTIME COLD AND FLU PLUS

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride kit

#### **Product Information**

Product Type HUMAN OTC DRUG Item Code (Source) NDC:69842-069

#### **Packaging**

# Item Code	Package Description	<b>Marketing Start Date</b>	Marketing End Date
1 NDC:69842-069-08	1 in 1 CARTON; Type 0: Not a Combination Product	12/13/2016	

#### **Quantity of Parts**

Part #	Package Quantity	Total Product Quantity
Part 1	1 BLISTER PACK	12
Part 2	1 BLISTER PACK	8

#### Part 1 of 2

#### DAY COLD AND FLU PLUS

acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride capsule, liquid filled

#### **Product Information**

Route of Administration ORAL

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
ACETAMINO PHEN (UNII: 36209 ITL9D) (ACETAMINO PHEN - UNII: 36209 ITL9D)	ACETAMINOPHEN	325 mg		
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9 D2RTI9 KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg		
<b>PHENYLEPHRINE HYDRO CHLO RIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE	5 mg		

Inactive Ingredients	
Ingredient Name	Strength
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PO VIDO NE (UNII: FZ989 GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics				
Color orange (clear) Score no score				
Shape	capsule (oblong)	Size	20 mm	
Flavor		Imprint Code	PC9	
Contains				

]	ackaging				
#	# Item Code Package Description Marketing Start Date Marketing End I				
1		1 in 1 CARTON			
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product			

Marketing Info	Marketing Information				
Marketing Category Application Number or Monograph Citation Marketing Start Date Marketing End Date					
OTC monograph final	part341	12/13/2016			

#### Part 2 of 2

#### NIGHTTIME COLD AND FLU PLUS

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride capsule, liquid filled

#### **Product Information**

Route of Administration ORAL

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
ACETAMINO PHEN (UNII: 36209 ITL9D) (ACETAMINO PHEN - UNII: 36209 ITL9D)	ACETAMINOPHEN	325 mg		
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9 D2RTI9 KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg		
<b>PHENYLEPHRINE HYDRO CHLO RIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE	5 mg		
DO XYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DO XYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg		

Inactive Ingredients	
Ingredient Name	Strength
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PO VIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics				
Color green (clear) Score no score				
Shape	capsule (oblong)	Size	20 mm	
Flavor		Imprint Code	PC22	
Contains				

Packaging	Packaging		
# Item Code	Package Description	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	1 in 1 CARTON		
1	8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information					
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
OTC monograph final	part341	12/13/20 16			
<b>Marketing Info</b>	Marketing Information				
Marketing Category Application Number or Monograph Citation Marketing Start Date Marketing End Date					
OTC monograph final	part341	12/13/2016			
o r o monographi imar	Parto-ri	12/15/2010			

### Labeler - CVS PHARMACY, INC. (062312574)

Establishment			
Name	Address	ID/FEI	Business Operations
Humanwell PuraCap Pharmaceuticals (Wuhan) Co., Ltd		421293287	manufacture(69842-069), analysis(69842-069)

Revised: 11/2019 CVS PHARMACY, INC.