

NON DROWSY DAYTIME COLD AND FLU SOFTGEL- acetaminophen, dextromethorphan hydrobromide, and phenylephrine hydrochloride capsule, liquid filled
KROGER COMPANY

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Non Drowsy Daytime Cold and Flu Softgel - 48ct

Drug Facts

Active ingredients (in each Softgel)	Purposes
Acetaminophen 325 mg	Pain reliever / Fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

temporarily relieves common cold/flu symptoms:

- nasal congestion
- cough due to minor throat & bronchial irritation
- sore throat
- headache
- minor aches & pains
- fever

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hrs, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- thyroid disease
- diabetes
- high blood pressure
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion or cough get worse or last more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults & for children even if you do not notice any signs or symptoms.

Directions

- take only as directed - see Overdose warning

- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	2 softgels with water every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- store at room temperature between 20-25 °C (68-77 °F)

Inactive ingredients

FD&C Red #40, FD&C Yellow #6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution, titanium dioxide

Questions or comments

1-800-632-6900

PRINCIPAL DISPLAY PANEL - 48 Softgel Blister Pack Carton

COMPARE TO the active ingredients of VICKS[®] DAYQUIL[™]
COLD & FLU RELIEF LIQUICAPS[™] *See bottom panel

NDC 30142-768-48

POWERFUL
NON-DROWSY
RELIEF

Kroger[®]

OUR PHARMACIST RECOMMENDED

Cold & Flu
DayTime
Multi-Symptom Relief

Acetaminophen
Dextromethorphan HBr
Phenylephrine HCl

actual
size

Pain Reliever
Fever Reducer
Cough Suppressant
Nasal Decongestant

48 SOFTGELS

COMPARE TO the active ingredients of VICKS® DAYQUIL™
COLD & FLU RELIEF LIQUICAPS™ *See bottom panel

NDC 30142-768-48



**POWERFUL
NON-DROWSY
RELIEF**

Cold & Flu



DayTime

Multi-Symptom Relief

**Acetaminophen
Dextromethorphan HBr
Phenylephrine HCl**

actual
size



Pain Reliever
Fever Reducer
Cough Suppressant
Nasal Decongestant

48 SOFTGELS



Cold & Flu



DayTime

Multi-Symptom Relief

**Acetaminophen
Dextromethorphan HBr
Phenylephrine HCl**

Pain Reliever
Fever Reducer
Cough Suppressant
Nasal Decongestant

48 SOFTGELS

84440A



OUR BRANDS. OUR GUARANTEE.
Love It or Your Money Back.
www.kroger.com

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

TAMPER EVIDENT: DO NOT USE IF BLISTER UNIT IS BROKEN OR TORN

Drug Facts

Active ingredients (in each Softgel)

Purposes

Acetaminophen 325 mg _____ Pain reliever / Fever reducer
 Dextromethorphan HBr 10 mg _____ Cough suppressant
 Phenylephrine HCl 5 mg _____ Nasal decongestant

Uses:

temporarily relieves common cold/flu symptoms:

- nasal congestion
- cough due to minor throat & bronchial irritation
- sore throat
- headache
- minor aches & pains
- fever

Warnings:

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hrs, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin redness
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have ■ liver disease ■ heart disease ■ thyroid disease ■ diabetes ■ high blood pressure ■ trouble urinating due to enlarged prostate gland ■ cough that occurs with too much phlegm (mucus) ■ persistent or chronic cough as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Stop use and ask a doctor if ■ you get nervous, dizzy or sleepless ■ pain, nasal congestion or cough get worse or last more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults & for children even if you do not notice any signs or symptoms.

**DISTRIBUTED BY
 THE KROGER CO.
 CINCINNATI, OHIO 45202
 MADE IN INDIA
 For More Product
 Information, Scan UPC
 Using Your Kroger App or
 Call 800-632-6900**

*Vicks® and DayQuil® are trademarks of The Procter & Gamble Company, Cincinnati, OH 45202. The Procter & Gamble Company is not affiliated with The Kroger Co. or this product.

Drug Facts (continued)

Directions:

- take only as directed - see Overdose warning
- do not exceed 4 doses per 24 hrs
- adults & children 12 yrs & over 2 softgels with water every 4 hrs
- children 4 to under 12 yrs ask a doctor
- children under 4 yrs do not use

Other information:

- store at room temperature between 20-25 °C (68-77 °F)

Inactive ingredients:
 FD&C red #40, FD&C yellow #6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution, titanium dioxide


Questions or comments: 1-800-632-6900

READ AND KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION




PLEASE RECYCLE
 CT3014276848
 REV.00-01 2021

LOT:
EXP:



NON DROWSY DAYTIME COLD AND FLU SOFTGEL
 acetaminophen, dextromethorphan hydrobromide, and phenylephrine hydrochloride capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:30142-768
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Acetaminophen (UNII: 362O9ITL9D) (Acetaminophen - UNII:362O9ITL9D)	Acetaminophen	325 mg
Dextromethorphan Hydrobromide (UNII: 9D2RTI9KYH) (Dextromethorphan - UNII:7355X3ROTS)	Dextromethorphan Hydrobromide	10 mg
Phenylephrine Hydrochloride (UNII: 04JA59TNSJ) (Phenylephrine - UNII:1WS297W6MV)	Phenylephrine Hydrochloride	5 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
Glycerin (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
Propylene Glycol (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	ORANGE	Score	no score
Shape	OVAL	Size	20mm
Flavor		Imprint Code	786
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:30142-768-48	4 in 1 CARTON	05/01/2021	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part341	05/01/2021	

