

BUTORPHANOL- butorphanol tartrate injection, solution

Aspen Veterinary Resources

ANADA # 200-332, Approved by FDA

BUTORPHANOL CIV

(butorphanol tartrate) INJECTION


CAUTION

Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian

DESCRIPTION

Butorphanol (butorphanol tartrate injection) is a totally synthetic centrally acting, narcotic agonist-antagonist analgesic with potent antitussive activity. It is a member of the phenanthrene series. The chemical name is Morphinan-3, 14-diol, 17-(cyclobutylmethyl)-,(-)-, (S-(R*,R*)))-2,3-dihydroxybutanedioate (1:1) (salt). It is a white, crystalline, water soluble substance having a molecular weight of 477.55; its molecular formula is $C_{21}H_{29}NO_2 \cdot C_4H_6O_6$.

Chemical Structure:



Each mL of Butorphanol contains 10 mg butorphanol base (as butorphanol tartrate), 3.3 mg citric acid, 6.4 mg sodium citrate, 4.7 mg sodium chloride, and 0.1 mg benzethonium chloride, q.s, with water for injection.

CLINICAL PHARMACOLOGY

Comparative Pharmacology

In animals, butorphanol has been demonstrated to be 4 to 30 times more potent than morphine and pentazocine (Talwin[®]-V) respectively.¹ In humans, butorphanol has been shown to have 5 to 7 times the analgesic activity of morphine and 20 times that of pentazocine.^{2,3} Butorphanol has 15 to 20 times the oral antitussive activity of codeine or dextromethorphan in dogs and guinea pigs.⁴

As an antagonist, butorphanol is approximately equivalent to nalorphine and 30 times more potent than pentazocine.¹

Cardiopulmonary depressant effects are minimal after treatment with butorphanol as demonstrated in dogs,⁵ humans^{6,7} and horses.⁸ Unlike classical narcotic agonist

analgesics which are associated with decreases in blood pressure, reduction in heart rate, and concomitant release of histamine, butorphanol does not cause histamine release.¹ Furthermore, the cardiopulmonary effects of butorphanol are not distinctly dosage related but rather reach a ceiling effect beyond which further dosage increases result in relatively lesser effects.

Reproduction: Studies performed in mice and rabbits revealed no evidence of impaired fertility or harm to the fetus due to butorphanol tartrate. In the female rat, parenteral administration was associated with increased nervousness and decreased care for the newborn, resulting in a decreased survival rate of the newborn. This nervousness was seen only in the rat species.


Equine Pharmacology

Following intravenous injection in horses, butorphanol is largely eliminated from the blood within 3 to 4 hours. The drug is extensively metabolized in the liver and excreted in the urine.

In ponies, butorphanol given intramuscularly at a dosage of 0.22 mg/kg was shown to alleviate experimentally induced visceral pain for about 4 hours.⁹

In horses, intravenous dosages of butorphanol ranging from 0.05 to 0.4 mg/kg were shown to be effective in alleviating visceral and superficial pain for at least four hours, as illustrated in the following figure:

Analgesic Effects of Butorphanol Given at Various Dosages in Horses with Abdominal Pain



*Pain threshold in butorphanol-treated colicky horses relative to placebo controls

A definite dosage-response relationship was detected in that butorphanol dosage of 0.1 mg/kg was more effective than 0.05 mg/kg but not different from 0.2 mg/kg in alleviating deep abdominal pain.

Acute Equine Studies

Rapid intravenous administration of butorphanol at a dosage of 2.0 mg/kg (20 times the recommended dosage) to a previously unmedicated horse resulted in a brief episode of inability to stand, muscle fasciculation, a convulsive seizure of 6 seconds duration, and recovery within three minutes. The same dosage administered after 10 successive daily 1.0 mg/kg dosages of butorphanol resulted only in transient sedative effects. During the

10 day course of administration at 1.0 mg/kg (10 times the recommended use level) in two horses, the only detectable drug effects were transient behavioral changes typical of narcotic agonist activity. These included muscle fasciculation about the head and neck, dysphoria, lateral nystagmus, ataxia, and salivation. Repeated administration of butorphanol at 1.0 mg/kg (10 times the recommended dose) every four hours for 48 hours caused constipation in one of two horses.

Subacute Equine Studies

Horses were found to tolerate butorphanol given intravenously at dosages of 0.1, 0.3, and 0.5 mg/kg every 4 hours for 48 hours followed by once daily injections for a total of 21 days. The only detectable drug effects were slight transient ataxia observed occasionally in the high dosage group. No clinical, laboratory, or gross or histopathologic evidence of any butorphanol-related toxicity was encountered in the horses.

INDICATIONS

Butorphanol (butorphanol tartrate injection) is indicated for the relief of pain associated with colic in adult horses and yearlings. Clinical studies in the horse have shown that butorphanol tartrate alleviates abdominal pain, associated with torsion, impaction, intussusception, spasmodic and tympanic colic, and postpartum pain.

WARNINGS

NOT FOR HUMAN USE. DO NOT USE IN HORSES INTENDED FOR HUMAN CONSUMPTION.

CAUTION

Butorphanol a potent analgesic, should be used with caution with other sedative or analgesic drugs as these are likely to produce additive effects.

There are no well-controlled studies using butorphanol in breeding horses, weanlings, and foals. Therefore, the drug should not be used in these groups.

ADVERSE REACTIONS

In clinical trials in horses, the most commonly observed side effect was slight ataxia which lasted 3 to 10 minutes. Marked ataxia was reported in 1.5% of the 327 horses treated. Mild sedation was reported in 9% of the horses.

DOSAGE

The recommended dosage in the horse is 0.1 mg of butorphanol per kilogram of body weight (0.05 mg/lb) by intravenous injection. This is equivalent to 5 mL of Butorphanol for each 1000 lbs body weight. The dose may be repeated within 3 to 4 hours but treatment should not exceed 48 hours. Pre-clinical model studies and clinical field trials in horses demonstrate that the analgesic effects of butorphanol tartrate are seen within 15 minutes following injection and persist for about 4 hours.

HOW SUPPLIED

Butorphanol (butorphanol tartrate injection) 10 mg base activity per mL.

NDC 46066-752-01 20 mL vial in package of one

NDC 46066-752-02 50 mL vial in package of one

STORAGE

Store at controlled room temperature 20° to 25°C (68° to 77°F). Protect from light.

REFERENCES

1. Pircio, A.W. *et al*: The Pharmacology of Butorphanol. *Arch. Int. Pharmacodyn. Ther.* 220(2):231-257, 1976.
2. Dobkin, A.B. *et al*: Butorphanol and Pentazocine in Patients with Severe Postoperative Pain. *Clin. Pharmacol. Ther.* 18:547-553, 1975.
3. Gilbert, M.S. *et al*: Intramuscular Butorphanol and Meperidine in Postoperative Pain. *Clin. Pharmacol. Ther.* 20:359-364, 1976.
4. Cavanagh, R.L. *et al*: Antitussive Properties of Butorphanol, *Arch. Int. Pharmacodyn. Ther.* 220 258-268, 1976.
5. Shurig, J.E. *et al*: Effect of Butorphanol and Morphine on Pulmonary Mechanics, Arterial Blood Pressure, and Venous Plasma Histamine in the Anesthetized Dog. *Arch. Int. Pharmacodyn. Ther.* 233:296-304, 1978.
6. Nagashmina, H. *et al*: Respiratory and Circulatory Effects of Intravenous Butorphanol and Morphine *Clin. Pharmacol. Ther.* 19:735-745, 1976.
7. Popio, K.A. *et al*: Hemodynamic and Respiratory Effects of Morphine and Butorphanol. *Clin. Pharm. Ther.* 23:281-287, 1978.
8. Robertson, J.T. and Muir, W.W.: Cardiopulmonary Effects of Butorphanol Tartrate in Horses. *Am. J. Vet. Res.* 42:41-44, 1981.
9. Kalpravidh, M. *et al*: Effects of Butorphanol, Flunixin, Levorphanol, Morphine, Pentazocine and Xylazine in Ponies. *Am. J. Vet. Res.* 45:217-223, 1984.

Approved by FDA under ANADA 200-332

Pivetal™

Manufactured for:

Aspen Veterinary Resources®, LTD.,

Liberty, MO 64068

www.pivetalvet.com

PVBUA0N Rev. 06/22

Principal Display Panel Text for Container Label:

NDC 46066-752-01

BUTORPHANOL CIV

(butorphanol tartrate)

INJECTION

Contains 10 mg butorphanol base

per mL as butorphanol tartrate, USP

Caution: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

ANADA 200-332, Approved by FDA

20 mL Pivetal™ Logo

NDC 46066-752-01

BUTORPHANOL CIV
(butorphanol tartrate)

INJECTION

Contains 10 mg butorphanol base per mL as butorphanol tartrate, USP

Caution: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.
ANADA 200-332, Approved by FDA

20 mL

Pivetal™

DOSAGE: By intravenous injection, 0.1 mg/kg body weight (0.05 mg/lb). This is equivalent to 5 mL for each 1000 lbs of body weight. Dose may be repeated within 3 to 4 hours. Treatment should not exceed 48 hours. Store at controlled room temperature 20° to 25°C (68° to 77°F). Protect from light.

FOR USE IN HORSES ONLY
DO NOT USE IN HORSES INTENDED FOR HUMAN CONSUMPTION. READ PACKAGE INSERT.

Each mL of solution contains: Butorphanol base as butorphanol tartrate, USP 10 mg, Citric acid, USP 3.3 mg, Sodium citrate, USP 6.4 mg, Sodium chloride, USP 4.7 mg, Benzethonium chloride, USP 0.1 mg, Water for injection, USP q.s.

Manufactured for: Aspen Veterinary Resources®, LTD.,
Liberty, MO 64068 | Pivetalvet.com
P147A Rev. 06/19 PVBUAAL Rev. 06/19

(01)00099355017382

LOT
EXP.

Principal Display Panel Text for Carton Label:

NDC 46066-752-01

Pivetal™ Logo

BUTORPHANOL CIV
(butorphanol tartrate)

INJECTION

Contains 10 mg butorphanol base per mL as butorphanol tartrate, USP

Caution: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

ANADA 200-332, Approved by FDA

20 mL



BUTORPHANOL

butorphanol tartrate injection, solution

Product Information

Product Type	PRESCRIPTION ANIMAL DRUG	Item Code (Source)	NDC:46066-752
Route of Administration	INTRAVENOUS	DEA Schedule	CIV

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Butorphanol tartrate (UNII: 2L7I72RUHN) (Butorphanol - UNII:QV897JC36D)	Butorphanol tartrate	10 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
Citric Acid Monohydrate (UNII: 2968PHW8QP)	
Sodium Citrate (UNII: 1Q73Q2JULR)	
Sodium Chloride (UNII: 451W47IQ8X)	
Benzethonium Chloride (UNII: PH41D05744)	

Water (UNII: 059QF0KO0R)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:46066-752-01	1 in 1 CARTON		
1		20 mL in 1 VIAL		
2	NDC:46066-752-02	1 in 1 CARTON		
2		50 mL in 1 VIAL		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANADA	ANADA200332	10/14/2019	

Labeler - Aspen Veterinary Resources (627265361)

Registrant - Akorn Operating Company LLC (117693100)

Establishment

Name	Address	ID/FEI	Business Operations
Akorn		117696790	PACK, LABEL

Establishment

Name	Address	ID/FEI	Business Operations
Akorn		117696832	MANUFACTURE, ANALYSIS, STERILIZE

Revised: 2/2022

Aspen Veterinary Resources