GENTAMICIN SULFATE- gentamicin sulfate solution/ drops REMEDYREPACK INC.

Gentamicin Sulfate Ophthalmic Solution, USP 0.3% (Sterile) Rx only

DESCRIPTION

Gentamicin sulfate ophthalmic solution, USP is a sterile, aqueous solution buffered to approximately pH 7.0 and formulated for ophthalmic use.

Each mL contains:

Active:gentamicin sulfate, USP (equivalent to 3 mg gentamicin).

Inactives:dibasic sodium phosphate, sodium chloride, monobasic sodium phosphate, purified water. Hydrochloric acid and/or sodium hydroxide may be added to adjust pH (6.5 - 7.5).

Preservative:benzalkonium chloride 0.01%.

Gentamicin is an aminoglycoside antibiotic obtained from cultures of *Micromonospora* purpurea. It is a mixture of the sulfate salts of gentamicin C1, C2, C1a and C2a. All three components appear to have similar antimicrobial activity. Gentamicin sulfate occurs as a white to buff powder and is soluble in water and insoluble in alcohol.

The structural formula is as follows:

CLINICAL PHARMACOLOGY

Microbiology

Gentamicin sulfate is active in vitroagainst many strains of the following microorganisms:

Staphylococcus aureus, Staphylococcus epidermidis, Streptococcus pyogenes, Streptococcus pneumoniae, Enterobacter aerogenes, Escherichia coli, Haemophilus influenzae, Klebsiella pneumoniae, Neisseria gonorrhoeae, Pseudomonas aeruginosa, and Serratia marcescens.

INDICATIONS AND USAGE

Gentamicin sulfate ophthalmic solution is indicated in the topical treatment of ocular bacterial infections including conjunctivitis, keratitis, keratoconjunctivitis, corneal ulcers, blepharitis, blepharoconjunctivitis, acute meibomianitis, and dacryocystitis, caused by susceptible strains of the following microorganisms:

Staphylococcus aureus, Staphylococcus epidermidis, Streptococcus pyogenes, Streptococcus pneumoniae, Enterobacter aerogenes, Escherichia coli, Haemophilus influenzae, Klebsiella pneumoniae, Neisseria gonorrhoeae, Pseudomonas aeruginosa, and Serratia marcescens.

CONTRAINDICATIONS

Gentamicin sulfate ophthalmic solution is contraindicated in patients with known hypersensitivity to any of its components.

WARNINGS

NOT FOR INJECTION INTO THE EYE

Gentamicin sulfate ophthalmic solution is not for injection. It should never be injected subconjunctivally, nor should it be directly introduced into the anterior chamber of the eye.

PRECAUTIONS

General

Prolonged use of topical antibiotics may give rise to overgrowth of nonsusceptible organisms including fungi. Bacterial resistance to gentamicin may also develop. If purulent discharge, inflammation or pain becomes aggravated, the patient should discontinue use of the medication and consult a physician. If irritation or hypersensitivity to any component of the drug develops, the patient should discontinue use of this preparation and appropriate therapy should be instituted.

Information for Patients

To avoid contamination, do not touch tip of container to the eye, eyelid or any surface.

Repackaged By / Distributed By: RemedyRepack Inc.

625 Kolter Drive, Indiana, PA 15701

Carcinogenesis, Mutagenesis, Impairment of Fertility

There are no published carcinogenicity or impairment of fertility studies on gentamicin. Aminoglycoside antibiotics have been found to be non-mutagenic.

Pregnancy

Gentamicin has been shown to depress body weights, kidney weights and median glomerular counts in newborn rats when administered systemically to pregnant rats in daily doses approximately 500 times the maximum recommended ophthalmic human dose. There are no adequate and well-controlled studies in pregnant women. Gentamicin should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

ADVERSE REACTIONS

Bacterial and fungal corneal ulcers have developed during treatment with gentamicin ophthalmic preparations.

The most frequently reported adverse reactions are ocular burning and irritation upon drug instillation, non-specific conjunctivitis, conjunctival epithelial defects and conjunctival hyperemia.

Other adverse reactions which have occurred rarely are allergic reactions, thrombocytopenic purpura and hallucinations.

To report SUSPECTED ADVERSE REACTIONS, contact Bausch & Lomb Incorporated at 1-800-553-5340 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DOSAGE AND ADMINISTRATION

Instill one or two drops into the affected eye(s) every four hours. In severe infections dosage may be increased to as much as two drops every hour.

Gentamicin sulfate ophthalmic solution USP, 0.3% is supplied sterile in a white LDPE plastic bottle, with a white LLDPE extended controlled drop tip and white polypropylene cap in the following size:

NDC: 70518-3785-00

PACKAGING: 1 in 1 CARTON, 5 mL in 1 BOTTLE, DROPPER TYPE 0

NOT FOR INJECTION INTO THE EYE

FOR OPHTHALMIC USE ONLY

Storage: Store between 2°C to 25°C (36°F to 77°F). Avoid exposure to excessive heat.

KEEP OUT OF REACH OF CHILDREN.

Repackaged and Distributed By:

Remedy Repack, Inc.

625 Kolter Dr. Suite #4 Indiana, PA 1-724-465-8762

DRUG: Gentamicin Sulfate

GENERIC: Gentamicin Sulfate

DOSAGE: SOLUTION/ DROPS

ADMINSTRATION: OPHTHALMIC

NDC: 70518-3785-0

PACKAGING: 5 mL in 1 BOTTLE, DROPPER

OUTER PACKAGING: 1 in 1 CARTON

ACTIVE INGREDIENT(S):

• GENTAMICIN SULFATE 3mg in 1mL

INACTIVE INGREDIENT(S):

- SODIUM CHLORIDE
- SODIUM PHOSPHATE, MONOBASIC, UNSPECIFIED FORM
- WATER
- HYDROCHLORIC ACID
- SODIUM HYDROXIDE
- BENZALKONIUM CHLORIDE
- SODIUM PHOSPHATE, DIBASIC, UNSPECIFIED FORM

Gentamicin Sulfate

0.3 %

Ophthalmic Solution OTY: 5 mL

FOR OPHTHALMIC USE ONLY; DO NOT USE IF IMPRINTED NECKBAND IS NOT INTACT.; (Sterile)



RX ONLY

NDC #: 70518-3785-00

Expires: LOT #:

Source NDC: 24208-0580-60

MFG: Bausch & Lomb, Bridgewater, NJ 08807

Keep this and all medication out of the reach of children

WARNING: NOT FOR INJECTION INTO THE EYE

Directions For Use: See Package Insert

Store at 20-25°C (68-77°F); excursions permitted to 15-30°C (59-86°F) [See USP]

Repackaged by: RemedyRepack Inc., Indiana, PA 15701, 724.465.8762

GENTAMICIN SULFATE

gentamicin sulfate solution/ drops

Product Information

Product Type

HUMAN PRESCRIPTION DRUG

Item Code (Source) NDC:70518-3785(NDC:24208-580)

Active Ingredient/Active Moiety

| . , | | |
|--|--------------------------|--------------|
| Ingredient Name | Basis of Strength | Strength |
| GENTAMICIN SULFATE (UNII: 8X7386QRLV) (GENTAMICIN - UNII:T6Z9V48IKG) | GENTAMICIN | 3 mg in 1 mL |

| Inactive Ingredients | | | | |
|--|----------|--|--|--|
| Ingredient Name | Strength | | | |
| SODIUM CHLORIDE (UNII: 451W47IQ8X) | | | | |
| SODIUM PHOSPHATE, MONOBASIC, UNSPECIFIED FORM (UNII: 3980JIH2SW) | | | | |
| WATER (UNII: 059QF0KO0R) | | | | |
| HYDROCHLORIC ACID (UNII: QTT17582CB) | | | | |
| SODIUM HYDROXIDE (UNII: 55X04QC32I) | | | | |
| BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7) | | | | |
| SODIUM PHOSPHATE, DIBASIC, UNSPECIFIED FORM (UNII: GR686LBA74) | | | | |

| F | Packaging | | | | | |
|---|----------------------|--|-------------------------|-----------------------|--|--|
| # | t Item Code | Package Description | Marketing Start Date | Marketing End Date | | |
| 1 | NDC:70518- 3785-0 | 1 in 1 CARTON | 07/11/2023 | | | |
| 1 | L | 5 mL in 1 BOTTLE, DROPPER; Type 0: Not a Combination Product | | | | |

| Marketing Information | | | |
|-----------------------|---|-------------------------|-----------------------|
| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
| ANDA | ANDA064048 | 07/11/2023 | |
| | | | |

Labeler - REMEDYREPACK INC. (829572556)

Revised: 3/2024 REMEDYREPACK INC.