

IBUPROFEN- ibuprofen capsule, liquid filled
EQUATE (Wal-Mart Stores, Inc.) (see also WAL-MART INC)

Drug Facts

Active ingredient (in each capsule)

Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)*

(present as the free acid and potassium salt)

***nonsteroidal anti-inflammatory drug**

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscular aches
 - minor pain of arthritis
 - toothache
 - backache
 - the common cold
 - menstrual cramps
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - vomit blood
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - slurred speech
 - trouble breathing
 - leg swelling
 - weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over:
 - take 1 capsule every 4 to 6 hours while symptoms persist
 - if pain or fever does not respond to 1 capsule, 2 capsules may be used
 - do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- **each capsule contains:** potassium 20 mg
- read all warnings and directions before use. Keep carton
- store between 20° to 25°C (68°to 77°F)
- avoid excessive heat above 40°C (104°F)
- swallow whole; do not crush, chew, or dissolve

Inactive ingredients

FD&C green #3, gelatin, medium-chain triglycerides, polyethylene glycol, potassium hydroxide, purified water, sorbitan, sorbitol, solution, white ink

Questions or comments?

Call **1-888-287-1915**

Principal Display Panel

Compare to Advil® Liqui-Gels® Active ingredient†

Ibuprofen

Capsules, 200 mg

Pain Reliever/Fever Reducer (NSAID)

SOFTGELS

**Liquid-Filled Capsules

TAMPER-EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS

BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

†This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Advil® Liqui-Gels®.

DISTRIBUTED BY: Walmart Inc., Bentonville, AR 72716

Product Label

Exp. Date:

PLD-DS248 F000599

Lot No.:

6 81131 24791 7

483574

DISTRIBUTED BY: Weimant Inc., Bentonville, AR 72716
This product is not manufactured or distributed by Pfizer
Consumer Healthcare, distributor of Advil® Liqui-Gels®.

Satisfaction guaranteed - For questions
or comments please call 1-888-287-1915.

Drug Facts

Active ingredient (in each capsule)

Solubilized ibuprofen equal to
200 mg Ibuprofen (NSAID)*
(present as the free acid and potassium salt)
*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/
fever reducer

Uses ■ temporarily relieves minor aches and pains
due to: ■ headache ■ muscular aches ■ minor
pain of arthritis ■ toothache ■ backache ■ the
common cold ■ menstrual cramps ■ temporarily
reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic
reaction, especially in people allergic to aspirin. Symptoms
may include: ■ hives ■ facial swelling ■ asthma
(wheezing) ■ shock ■ skin reddening ■ rash
■ blisters. If an allergic reaction occurs, stop use and seek
medical help right away.

Stomach bleeding warning: This product contains an
NSAID, which may cause severe stomach bleeding.
The chance is higher if you: ■ are age 60 or older
■ have had stomach ulcers or bleeding problems ■ take
a blood thinning (anticoagulant) or steroid drug ■ take
other drugs containing prescription or nonprescription
NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have
3 or more alcoholic drinks every day while using this

Drug Facts (continued)

product ■ take more or for a longer time than directed
Heart attack and stroke warning: NSAIDs, except
aspirin, increase the risk of heart attack, heart failure, and
stroke. These can be fatal. The risk is higher if you use
more than directed or for longer than directed.

Do not use ■ if you have ever had an allergic reaction
to any other pain reliever/fever reducer ■ right before
or after heart surgery

Ask a doctor before use if ■ the stomach bleeding
warning applies to you ■ you have problems or serious
side effects from taking pain relievers or fever reducers
■ you have a history of stomach problems, such as
heartburn ■ you have high blood pressure, heart
disease, liver cirrhosis, kidney disease, asthma, or had a
stroke ■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are
■ under a doctor's care for any serious condition
■ taking aspirin for heart attack or stroke, because
ibuprofen may decrease this benefit of aspirin
■ taking any other drug

When using this product, take with food or milk if
stomach upset occurs.

Stop use and ask a doctor if ■ you experience any of
the following signs of stomach bleeding: ■ feel faint
■ vomit blood ■ have bloody or black stools ■ have
stomach pain that does not get better

Drug Facts (continued)

■ store between 20° to 25°C (68° to 77°F)
■ avoid excessive heat above 40°C (104°F)
■ swallow whole; do not crush, chew, or dissolve

Inactive ingredients FD&C green #3, gelatin,
medium-chain triglycerides, polyethylene glycol,
potassium hydroxide, purified water, sorbitol sorbitan
solution, white ink

Questions or comments?

1-888-287-1915

TAMPER EVIDENCE DO NOT USE IF PRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

KEEP OUTER CARTON FOR COMPLETE
WARNINGS AND PRODUCT INFORMATION

Drug Facts (continued)
■ you have symptoms of heart problems or stroke:
■ chest pain ■ slurred speech ■ trouble breathing
■ leg swelling ■ weakness in one part or side of body
■ pain gets worse or lasts more than 10 days ■ fever
■ gets worse or lasts more than 3 days ■ redness or
swelling is present in the painful area ■ any new
symptoms appear
if pregnant or breast-feeding, ask a health professional
before use. It is especially important not to use ibuprofen
during the last 3 months of pregnancy unless definitely
directed to do so by a doctor because it may cause
problems in the unborn child or complications during
delivery. **Keep out of reach of children.** In case of
overdose, get medical help or contact a Poison Control
Center right away (1-800-222-1222).

Directions ■ do not take more than directed
■ the smallest effective dose should be used
■ adults and children 12 years and over: ■ take
1 capsule every 4 to 6 hours while symptoms persist
■ if pain or fever does not respond to 1 capsule, 2
capsules may be used ■ do not exceed 6 capsules in
24 hours, unless directed by a doctor ■ children under
12 years: ask a doctor

Other information ■ each capsule contains:
potassium 20 mg ■ read all warnings and directions
before use. Keep carton.

equate™
Ibuprofen
Capsules, 200mg



NDC 49035-745-20

Pain Reliever/ Fever Reducer (NSAID)



Actual Size

200
mg

20
SOFTGELS**

**Liquid-Filled Capsules

EQUATE Ibuprofen 200 mg Softgels

IBUPROFEN

ibuprofen capsule, liquid filled

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:49035-745

Route of Administration

ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
GELATIN (UNII: 2G86QN327L)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
SORBITAN (UNII: 6O92ICV9RU)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	

Product Characteristics

Color	blue	Score	no score
Shape	CAPSULE	Size	19mm
Flavor		Imprint Code	PC16
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49035-745-30	300 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/31/2018	03/31/2025
2	NDC:49035-745-60	160 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/31/2018	03/31/2025
3	NDC:49035-745-40	1 in 1 BOX	03/31/2018	03/31/2025
3		40 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
4	NDC:49035-745-80	80 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/31/2018	03/31/2025
5	NDC:49035-745-20	1 in 1 BOX	03/31/2018	03/31/2025
5		20 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA206568	03/31/2018	03/31/2025

Labeler - EQUATE (Wal-Mart Stores, Inc.) (see also WAL-MART INC) (051957769)

Revised: 10/2022

EQUATE (Wal-Mart Stores, Inc.) (see also WAL-MART INC)