

**VERAPAMIL HYDROCHLORIDE - verapamil hydrochloride capsule, extended release**  
**Physicians Total Care, Inc.**

**HIGHLIGHTS OF PRESCRIBING INFORMATION**

These highlights do not include all the information needed to use verapamil hydrochloride extended-release capsules (PM) safely and effectively. See full prescribing information for verapamil hydrochloride extended-release capsules (PM).

**Verapamil Hydrochloride Extended-release Capsules (PM) (Controlled Onset) for Oral use**

**Initial U.S. Approval: 1998**

**INDICATIONS AND USAGE**

Verapamil hydrochloride is a calcium channel blocker indicated for the management of essential hypertension. (1)

**DOSAGE AND ADMINISTRATION**

- Do not crush or chew capsule contents; swallow capsule whole or sprinkle entire contents onto applesauce. (2.2, 17)
- Usual dosage: 200 mg once daily at bedtime; if inadequate response, titrate upward to 300 mg, then 400 mg once daily at bedtime. (2.1)
- Initial dose of 100 mg once daily at bedtime in patients with renal or hepatic impairment, elderly or low-weight patients. (2.1)

**DOSAGE FORMS AND STRENGTHS**

- Extended-release capsules controlled-onset: 100 mg, 200 mg, 300 mg. (3)

**CONTRAINDICATIONS**

- Severe left ventricular dysfunction. (4)
- Hypotension (< 90 mm Hg systolic pressure) or cardiogenic shock. (4)
- Sick sinus syndrome (except in patients with pacemaker). (4)
- 2nd- or 3rd-degree AV block (except in patients with pacemaker). (4)
- Atrial flutter or atrial fibrillation and an accessory bypass tract. (4)

**WARNINGS AND PRECAUTIONS**

- Congestive heart failure or pulmonary edema may develop. (5.1)
- Hypotension/dizziness may occur. (5.2)
- Elevated transaminases have occurred; monitor liver function. (5.3)
- Ventricular fibrillation has occurred in patients with atrial flutter or atrial fibrillation and an accessory bypass tract. (5.4)
- Reduce dose or discontinue therapy if marked first-degree AV block or progression to second- or third-degree AV block occurs. (5.5)
- Sinus bradycardia, pulmonary edema, severe hypotension, second-degree AV block, sinus arrest and death occurred in patients with hypertrophic cardiomyopathy. (5.6)

**ADVERSE REACTIONS**

Most common adverse reactions (incidence  $\geq$  3% and more common than in patients treated with placebo) are headache, infection, constipation, flu syndrome, peripheral edema, dizziness, pharyngitis and sinusitis. (6.1)

**To report SUSPECTED ADVERSE REACTIONS, contact Mylan Pharmaceuticals Inc. at 1-877-446-3769 (1-877-4-INFO-RX) or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch)**

**DRUG INTERACTIONS**

- CYP3A4 inhibitors increase verapamil levels. (7.1)
- CYP3A4 inducers decrease verapamil levels. (7.1)
- Grapefruit juice may significantly increase verapamil levels. (7.2)
- Beta blockers: reports of excess bradycardia and AV block, including complete heart block; monitor closely. (7.3)
- Digoxin levels can increase by 50 to 75%; reduce digoxin dose. (7.4)
- Alcohol elimination inhibited resulting in elevated ethanol levels. (7.5)

**USE IN SPECIFIC POPULATIONS**

- Use during pregnancy only if potential benefit justifies the potential risk to the fetus. (8.1)
- Discontinue nursing while verapamil is administered. (8.3)
- Safety and effectiveness in pediatric patients have not been established. (8.4)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 6/2012

**FULL PRESCRIBING INFORMATION: CONTENTS\***

**1 INDICATIONS AND USAGE**

**2 DOSAGE AND ADMINISTRATION**

2.1 Essential Hypertension

2.2 Sprinkling the Capsule Contents on Food

**3 DOSAGE FORMS AND STRENGTHS**

**4 CONTRAINDICATIONS**

**5 WARNINGS AND PRECAUTIONS**

5.1 Heart Failure

5.2 Hypotension

5.3 Elevated Liver Enzymes

5.4 Accessory Bypass Tract (Wolff-Parkinson-White or Lown-Ganong Levine)

5.5 Atrioventricular Block

5.6 Patients with Hypertrophic Cardiomyopathy

## **6 ADVERSE REACTIONS**

6.1 Clinical Trials Experience

6.2 Open Trials / Post-marketing Experience

6.3 Treatment of Acute Cardiovascular Adverse Reactions

## **7 DRUG INTERACTIONS**

7.1 CYP3A4 Inhibitors and Inducers

7.2 Grapefruit Juice

7.3 Beta-Blockers

7.4 Digitalis

7.5 Alcohol

7.6 Clonidine

7.7 Telithromycin

7.8 Antineoplastic Agents

7.9 Quinidine

7.10 Aspirin

7.11 Antihypertensive agents

7.12 Disopyramide

7.13 Flecainide

7.14 Carbamazepine

7.15 Cyclosporine

7.16 Lithium

7.17 Inhalation Anesthetics

7.18 Neuromuscular Blocking Agents

7.19 Phenobarbital

7.20 Rifampin

7.21 Theophylline

7.22 Cimetidine

7.23 Nitrates

## **8 USE IN SPECIFIC POPULATIONS**

8.1 Pregnancy

8.2 Labor and Delivery

8.3 Nursing Mothers

8.4 Pediatric Use

8.5 Geriatric Use

8.6 Impaired Hepatic Function

8.7 Impaired Renal Function

8.8 Attenuated (decreased) Neuromuscular Transmission

## **10 OVERDOSAGE**

## **11 DESCRIPTION**

## **12 CLINICAL PHARMACOLOGY**

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

## **13 NONCLINICAL TOXICOLOGY**

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

13.2 Animal Toxicology and/or Pharmacology

## **14 CLINICAL STUDIES**

## **16 HOW SUPPLIED/STORAGE AND HANDLING**

## **17 PATIENT COUNSELING INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## **FULL PRESCRIBING INFORMATION**

### **1 INDICATIONS AND USAGE**

Verapamil hydrochloride extended-release capsules (PM) for oral use are indicated for the management of essential hypertension.

### **2 DOSAGE AND ADMINISTRATION**

**THE CONTENTS OF THE VERAPAMIL HYDROCHLORIDE EXTENDED-RELEASE CAPSULE (PM) SHOULD NOT BE CRUSHED OR CHEWED. VERAPAMIL HYDROCHLORIDE EXTENDED-RELEASE CAPSULES (PM) ARE TO BE SWALLOWED WHOLE OR THE ENTIRE CONTENTS OF THE CAPSULE SPRINKLED ONTO APPLESAUCE.**

#### **2.1 Essential Hypertension**

Administer verapamil hydrochloride extended-release capsules (PM) once daily at bedtime. Clinical trials studied doses of 100 mg, 200 mg, 300 mg and 400 mg. The usual daily dose of verapamil hydrochloride extended-release capsules (PM) in clinical trials has been 200 mg given by mouth once daily at bedtime. In rare instances, initial doses of 100 mg a day may be warranted in patients who have an increased response to verapamil, [e.g., patients with impaired renal function, impaired hepatic function, elderly, low-weight patients, etc. (*see Use in Specific Populations (8.5, 8.6, 8.7)*)]. Base upward titration on therapeutic efficacy and safety evaluated approximately 24 hours after dosing. The antihypertensive effects of verapamil hydrochloride extended-release capsules (PM) are evident within the first week of therapy.

If an adequate response is not obtained with 200 mg of verapamil hydrochloride extended-release capsules (PM), the dose may be titrated upward in the following manner:

1. 300 mg each evening
2. 400 mg each evening (2 x 200 mg)

When verapamil hydrochloride extended-release capsules (PM) are administered at bedtime, office evaluation of blood pressure during morning and early afternoon hours is essentially a measure of peak effect. The usual evaluation of trough effect, which sometimes might be needed to evaluate the appropriateness of any given dose of verapamil hydrochloride extended-release capsules (PM), would be just prior to bedtime.

## 2.2 Sprinkling the Capsule Contents on Food

Verapamil hydrochloride extended-release capsules (PM) may also be administered by carefully opening the capsule and sprinkling the beads onto one tablespoonful of applesauce. Swallow the applesauce immediately without chewing and follow with a glass of cool water to ensure complete swallowing of the beads. The applesauce used should not be hot, and it should be soft enough to be swallowed without chewing. Use any bead/applesauce mixture immediately and do not store for future use. Absorption of the beads sprinkled onto other foods has not been tested. This method of administration may be beneficial for patients who have difficulty swallowing whole capsules. Subdividing the contents of a verapamil hydrochloride extended-release capsule (PM) is not recommended.

## 3 DOSAGE FORMS AND STRENGTHS

Verapamil hydrochloride extended-release capsules (PM), 100 mg, are hard-shell gelatin capsules with a red opaque cap and a white opaque body filled with white to off-white beads. The capsule is axially printed with **MYLAN** over **6201** in black ink on both the cap and the body.

Verapamil hydrochloride extended-release capsules (PM), 200 mg, are hard-shell gelatin capsules with a red opaque cap and a light orange opaque body filled with white to off-white beads. The capsule is axially printed with **MYLAN** over **6202** in black ink on both the cap and the body.

Verapamil hydrochloride extended-release capsules (PM), 300 mg, are hard-shell gelatin capsules with a red opaque cap and a red opaque body filled with white to off-white beads. The capsule is axially printed with **MYLAN** over **6203** in black ink on both the cap and the body.

## 4 CONTRAINDICATIONS

Verapamil is contraindicated in:

- Severe left ventricular dysfunction [*see Warnings and Precautions (5.1)*].
- Hypotension (less than 90 mm Hg systolic pressure) or cardiogenic shock.
- Sick sinus syndrome (except in patients with a functioning artificial ventricular pacemaker).
- Second- or third-degree AV block (except in patients with a functioning artificial ventricular pacemaker).
- Patients with atrial flutter or atrial fibrillation and an accessory bypass tract (e.g., Wolff-Parkinson-White, Lown-Ganong-Levine syndromes) [*see Warnings and Precautions (5.4)*].

## 5 WARNINGS AND PRECAUTIONS

### 5.1 Heart Failure

Verapamil has a negative inotropic effect which, in most patients, is compensated by its afterload reduction (decreased systemic vascular resistance) properties without a net impairment of ventricular performance. In previous clinical experience with 4,954 patients primarily with immediate-release verapamil, 87 (1.8%) developed congestive heart failure or pulmonary edema. Avoid verapamil in patients with severe left ventricular dysfunction (e.g., ejection fraction less than 30% or moderate to severe symptoms of cardiac failure) and in patients with any degree of ventricular dysfunction if they are receiving a beta-adrenergic blocker [*see Drug Interactions (7.3)*]. Control patients with milder ventricular dysfunction, if possible, with optimum doses of digitalis and/or diuretics before verapamil treatment is started [*see Drug Interactions (7.4)*].

## 5.2 Hypotension

Occasionally, the pharmacologic action of verapamil may produce a decrease in blood pressure below normal levels which may result in dizziness or symptomatic hypotension. In hypertensive patients, decreases in blood pressure below normal are unusual. The incidence of hypotension observed in 4,954 patients enrolled in clinical trials of other verapamil formulations was 2.5% [see *Adverse Reactions* (6.1)]. In clinical studies of verapamil hydrochloride extended-release capsules (PM), 1.7% of the patients developed significant hypotension. Tilt table testing (60 degrees) was not able to induce orthostatic hypotension.

## 5.3 Elevated Liver Enzymes

Elevations of transaminases with and without concomitant elevations in alkaline phosphatase and bilirubin have been reported. Such elevations have sometimes been transient and may disappear even in the face of continued verapamil treatment.

Several cases of hepatocellular injury related to verapamil have been proven by rechallenge; half of these had clinical symptoms (malaise, fever and/or right upper quadrant pain) in addition to elevations of SGOT, SGPT and alkaline phosphatase. Periodic monitoring of liver function in patients receiving verapamil is therefore prudent.

## 5.4 Accessory Bypass Tract (Wolff-Parkinson-White or Lown-Ganong Levine)

Some patients with paroxysmal and/or chronic atrial flutter or atrial fibrillation and a coexisting accessory AV pathway have developed increased antegrade conduction across the accessory pathway bypassing the AV node, producing a very rapid ventricular response or ventricular fibrillation after receiving intravenous verapamil (or digitalis). Although a risk of this occurring with oral verapamil has not been established, such patients receiving oral verapamil may be at risk and its use in these patients is contraindicated [see *Contraindications* (4)]. Treatment is usually DC-cardioversion. Cardioversion has been used safely and effectively after oral verapamil.

## 5.5 Atrioventricular Block

The effect of verapamil on AV conduction and the SA node may lead to asymptomatic first-degree AV block and transient bradycardia, sometimes accompanied by nodal escape rhythms. PR interval prolongation is correlated with verapamil plasma concentrations, especially during the early titration phase of therapy. Higher degrees of AV block, however, were infrequently (0.8%) observed in previous verapamil clinical trials [see *Adverse Reactions* (6.1)]. Marked first-degree block or progressive development to second- or third-degree AV block requires a reduction in dosage or, in rare instances, discontinuation of verapamil and institution of appropriate therapy depending upon the clinical situation.

## 5.6 Patients with Hypertrophic Cardiomyopathy

In 120 patients with hypertrophic cardiomyopathy, idiopathic hypertrophic subaortic stenosis (IHSS) (most of them refractory or intolerant to propranolol) who received therapy with verapamil at doses up to 720 mg/day, a variety of serious adverse effects were seen. Three patients died in pulmonary edema; all had severe left ventricular outflow obstruction and a history of left ventricular dysfunction. Eight other patients had pulmonary edema and/or severe hypotension; abnormally high (over 20 mm Hg) pulmonary capillary wedge pressure and a marked left ventricular outflow obstruction were present in most of these patients. Concomitant administration of quinidine [see *Drug Interactions* (7.9)] preceded the severe hypotension in three of the eight patients (two of whom developed pulmonary edema). Sinus bradycardia occurred in 11% of the patients, second-degree AV block in 4% and sinus arrest in 2% [see *Adverse Reactions* (6)]. It must be appreciated that this group of patients had a serious disease with a high mortality rate. Most adverse effects responded well to dose reduction and only rarely did verapamil have to be discontinued.

# 6 ADVERSE REACTIONS

## 6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice. The adverse reaction information from clinical trials does, however, provide a basis for identifying the adverse events that appear to be related to drug use and for approximating rates. Serious adverse reactions are uncommon when verapamil therapy is initiated with upward dose titration within the recommended single and total daily dose. See *Warnings and Precautions* (5.1, 5.2, 5.3, 5.4, 5.5) for discussion of heart failure, hypotension, elevated liver enzymes, AV block and rapid ventricular response. Reversible (upon discontinuation of verapamil) non-obstructive, paralytic ileus has been infrequently reported in association with the use of verapamil. The following reactions (Table 1) to orally administered verapamil hydrochloride extended-release capsules (PM) occurred at rates of 2% or greater or occurred at lower rates but appeared to be drug-related in clinical trials in hypertension.

**Table 1. Adverse Events Occurring in  $\geq 2\%$  of Verapamil Hydrochloride Extended-release Capsules (PM) Patients in Placebo-Controlled Clinical Trials**

	All Doses Studied N = 297 %	Placebo N = 116 %		All Doses Studied N = 297 %	Placebo N = 116 %
Headache	12.1	11.2	Dyspepsia	2.7	1.7
Infection	12.1*	6.9	Rhinitis	2.7	2.6
Constipation	8.8*	0.9	Diarrhea	2.4	1.7
Flu Syndrome	3.7	2.6	Pain	2.4	1.7
Peripheral edema	3.7	0.9	Edema	1.7	0
Dizziness	3	0.9	Nausea	1.7	0
Pharyngitis	3	2.6	Accidental Injury	1.5	0
Sinusitis	3	2.6			

\* Infection, primarily upper respiratory infection (URI) and unrelated to study medication. Constipation was typically mild and easily manageable. At the usual once daily dose of 200 mg, the observed incidence of constipation was 3.9%.

In previous experience with other formulations of verapamil (N = 4,954) the following reactions (Table 2) have occurred at rates greater than 1% or occurred at lower rates but appeared clearly drug-related in clinical trials in 4,954 patients.

**Table 2. Adverse Events Occurring in  $> 1\%$  (or lower rates and clearly drug-related) of Patients with Other Verapamil Formulations**

Constipation	7.3%	Fatigue	1.7%
Dizziness	3.3%	Bradycardia (HR<50/min)	1.4%
Nausea	2.7%	Rash	1.2%
Hypotension	2.5%	AV block (total 1°, 2°, 3°)	1.2%
Headache	2.2%	AV block (2° and 3°)	0.8%
Edema	1.9%	Flushing	0.6%
CHF/Pulmonary Edema	1.8%		

In clinical trials related to the control of ventricular response in patients taking digoxin who had atrial fibrillation or atrial flutter, ventricular rate below 50/min at rest occurred in 15% of patients and asymptomatic hypotension occurred in 5% of patients.

## 6.2 Open Trials / Post-marketing Experience

The following reactions, reported with orally administered verapamil in 2% or less of patients, occurred under conditions [open verapamil trials, post-marketing experience (reactions added since the initial US approval of verapamil hydrochloride extended-release capsules (PM) in 1998 are marked with an asterisk)] where a causal relationship is uncertain; they are listed to alert the physician to a possible relationship:

**Cardiovascular:** angina pectoris, atrioventricular dissociation, ECG Abnormal\*, chest pain, claudication, hypertension\*, myocardial infarction, palpitations, purpura (vasculitis), syncope.

**Digestive System:** diarrhea, dry mouth, elevated liver enzymes\* [see Warnings and Precautions (5.3)], gastrointestinal distress, gingival hyperplasia.

**Hemic and Lymphatic:** ecchymosis or bruising.

**Nervous System:** cerebrovascular accident, confusion, equilibrium disorders, extrapyramidal symptoms, insomnia, muscle cramps, paresthesia, psychotic symptoms, shakiness, somnolence.

**Respiratory:** dyspnea.

**Skin:** arthralgia and rash, exanthema, hair loss, hyperkeratosis, macules, sweating, urticaria, Stevens-Johnson Syndrome, erythema multiforme.

**Special Senses:** blurred vision, tinnitus.

**Urogenital:** gynecomastia, galactorrhea/hyperprolactinemia, impotence, increased urination, spotty menstruation.

**Other:** allergy aggravated, asthenia\*.

## 6.3 Treatment of Acute Cardiovascular Adverse Reactions

The frequency of cardiovascular adverse reactions that require therapy is rare; hence, experience with their treatment is limited. Whenever severe hypotension or complete AV block occurs following oral administration of verapamil, apply the appropriate emergency measures immediately; e.g., intravenously administered norepinephrine bitartrate, atropine sulfate, isoproterenol hydrochloride (all in the usual doses), or calcium gluconate (10% solution). In patients with hypertrophic cardiomyopathy, use alpha-adrenergic agents (phenylephrine hydrochloride, metaraminol bitartrate, or methoxamine hydrochloride

to maintain blood pressure, and isoproterenol and avoid norepinephrine. If further support is necessary, inotropic agents (dopamine hydrochloride or dobutamine hydrochloride) may be administered. Actual treatment and dosage depends on the severity of the clinical situation and the judgment and experience of the treating physician.

## **7 DRUG INTERACTIONS**

### **7.1 CYP3A4 Inhibitors and Inducers**

*In vitro* metabolic studies indicate that verapamil is metabolized by cytochrome P450 CYP3A4, CYP1A2 and CYP2C. Clinically significant interactions have been reported with inhibitors of CYP3A4 (e.g., erythromycin, ritonavir) causing elevation of plasma levels of verapamil. Hypotension, bradyarrhythmias and lactic acidosis have been observed in patients receiving concurrent telithromycin, an antibiotic in the ketolide class of antibiotics. Inducers of CYP3A4 (e.g., rifampin) have caused a lowering of plasma levels of verapamil.

### **7.2 Grapefruit Juice**

Grapefruit juice may significantly increase concentrations of verapamil. Grapefruit juice given to nine healthy volunteers increased S- and R-verapamil AUC<sub>0-12</sub> by 36% and 28%, respectively. Steady-state C<sub>max</sub> and C<sub>min</sub>, of S-verapamil increased by 57% and 16.7%, respectively with grapefruit juice compared to control. Similarly, C<sub>max</sub> and C<sub>min</sub>, of R-verapamil increased by 40% and 13%, respectively. Grapefruit juice did not affect half-life, nor was there a significant change in AUC<sub>0-12</sub> ratio R/S compared to control. Grapefruit juice did not cause a significant difference in the pharmacokinetics of norverapamil. This increase in verapamil plasma concentration is not expected to have any clinical consequences.

### **7.3 Beta-Blockers**

Concomitant therapy with beta-adrenergic blockers and verapamil may result in additive negative effects on heart rate, atrioventricular conduction and/or cardiac contractility. The combination of extended-release verapamil and beta-adrenergic blocking agents has not been studied. However, there have been reports of excess bradycardia and AV block, including complete heart block, when the combination has been used for the treatment of hypertension. For hypertensive patients, the risk of combined therapy may outweigh the potential benefits. The combination should be used only with caution and close monitoring. Asymptomatic bradycardia (36 beats/min) with a wandering atrial pacemaker has been observed in a patient receiving concomitant timolol (a beta-adrenergic blocker) eyedrops and oral verapamil.

A decrease in metoprolol and propranolol clearance has been observed when either drug is administered concomitantly with verapamil. A variable effect has been seen when verapamil and atenolol were given together.

### **7.4 Digitalis**

Consider reducing digoxin dose when verapamil and digoxin are to be given together. Monitor digoxin level periodically during therapy. Chronic verapamil treatment can increase serum digoxin levels by 50% to 75% during the first week of therapy, and this can result in digitalis toxicity. In patients with hepatic cirrhosis the influence of verapamil on digoxin pharmacokinetics is magnified. Verapamil may reduce total body clearance and extrarenal clearance of digitoxin by 27% and 29%, respectively. If digoxin toxicity is suspected, suspend or discontinue digoxin therapy.

In previous clinical trials with other verapamil formulations related to the control of ventricular response in patients taking digoxin who had atrial fibrillation or atrial flutter, ventricular rates below 50/min at rest occurred in 15% of patients, and asymptomatic hypotension occurred in 5% of patients.

### **7.5 Alcohol**

Verapamil has been found to significantly inhibit ethanol elimination resulting in elevated blood ethanol concentrations that may prolong the intoxicating effects of alcohol.

### **7.6 Clonidine**

Sinus bradycardia resulting in hospitalization and pacemaker insertion has been reported in association with the use of clonidine concurrently with verapamil. Monitor heart rate in patients receiving concomitant verapamil and clonidine.

### **7.7 Telithromycin**

Hypotension, bradyarrhythmias and lactic acidosis have been observed in patients receiving concurrent telithromycin, an antibiotic in the ketolide class of antibiotics.

### **7.8 Antineoplastic Agents**

Verapamil can increase doxorubicin levels. The absorption of verapamil can be reduced by the cyclophosphamide, oncovin, procarbazine, prednisone (COPP) and the vindesine, adriamycin, cisplatin (VAC) cytotoxic drug regimens. Concomitant administration of R-verapamil can decrease the clearance

of paclitaxel.

### **7.9 Quinidine**

In a small number of patients with hypertrophic cardiomyopathy, concomitant use of verapamil and quinidine resulted in significant hypotension. Until further data are obtained, avoid combined therapy of verapamil and quinidine in patients with hypertrophic cardiomyopathy.

The electrophysiological effects of quinidine and verapamil on AV conduction were studied in eight patients. Verapamil significantly counteracted the effects of quinidine on AV conduction. There has been a report of increased quinidine levels during verapamil therapy.

### **7.10 Aspirin**

In a few reported cases, coadministration of verapamil with aspirin has led to increased bleeding times greater than observed with aspirin alone.

### **7.11 Antihypertensive agents**

Verapamil administered concomitantly with oral antihypertensive agents (e.g., vasodilators, angiotensin-converting enzyme inhibitors, diuretics, beta-blockers) will usually have an additive effect on lowering blood pressure. Monitor patients receiving these combinations appropriately. Concomitant use of agents that attenuate alpha-adrenergic function with verapamil may result in reduction in blood pressure that is excessive in some patients. Such an effect was observed in one study following the concomitant administration of verapamil and prazosin.

### **7.12 Disopyramide**

Until data on possible interactions between verapamil and disopyramide are obtained, do not administer disopyramide within 48 hours before or 24 hours after verapamil administration.

### **7.13 Flecainide**

A study in healthy volunteers showed that the concomitant administration of flecainide and verapamil may have additive effects on myocardial contractility, AV conduction and repolarization. Concomitant therapy with flecainide and verapamil may result in additive negative inotropic effect and prolongation of atrioventricular conduction.

### **7.14 Carbamazepine**

Verapamil therapy may increase carbamazepine concentrations during combined therapy. This may produce carbamazepine side effects such as diplopia, headache, ataxia or dizziness.

### **7.15 Cyclosporine**

Verapamil therapy may increase serum levels of cyclosporine.

### **7.16 Lithium**

Increased sensitivity to the effects of lithium (neurotoxicity) has been reported during concomitant verapamil-lithium therapy with either no change or an increase in serum lithium levels. However, the addition of verapamil has also resulted in the lowering of serum lithium levels in patients receiving chronic stable oral lithium. Patients receiving both drugs must be monitored carefully.

### **7.17 Inhalation Anesthetics**

Animal experiments have shown that inhalation anesthetics depress cardiovascular activity by decreasing the inward movement of calcium ions. When used concomitantly, inhalation anesthetics and calcium antagonists, such as verapamil, titrate slowly to avoid excessive cardiovascular depression.

### **7.18 Neuromuscular Blocking Agents**

Clinical data and animal studies suggest that verapamil may potentiate the activity of neuromuscular blocking agents (curare-like and depolarizing). It may be necessary to decrease the dose of verapamil and/or the dose of the neuromuscular blocking agent when the drugs are used concomitantly.

### **7.19 Phenobarbital**

Phenobarbital therapy may increase verapamil clearance.

### **7.20 Rifampin**

Therapy with rifampin may markedly reduce oral verapamil bioavailability.

### **7.21 Theophylline**

Verapamil may inhibit the clearance and increase the plasma levels of theophylline.

### **7.22 Cimetidine**

The interaction between cimetidine and chronically administered verapamil has not been studied.

Variable results on clearance have been obtained in acute studies of healthy volunteers; clearance of verapamil was either reduced or unchanged.

### **7.23 Nitrates**

Verapamil has been given concomitantly with short- and long-acting nitrates without any undesirable drug interactions. The pharmacologic profile of both drugs and the clinical experience suggest beneficial interactions.

## **8 USE IN SPECIFIC POPULATIONS**

### **8.1 Pregnancy**

Teratogenic Effects. Pregnancy Category C

Reproduction studies have been performed in rabbits and rats at oral doses up to 1.9 (15 mg/kg/day) and 7.5 (60 mg/kg/day) times the human oral daily dose, respectively, and have revealed no evidence of teratogenicity. In the rat, however, this multiple of the human dose was embryocidal and retarded fetal growth and development, probably because of adverse maternal effects reflected in reduced weight gains of the dams. This oral dose has also been shown to cause hypotension in rats. There are no adequate and well controlled studies in pregnant women. Verapamil should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus. Verapamil crosses the placental barrier and can be detected in umbilical vein blood at delivery.

### **8.2 Labor and Delivery**

It is not known whether the use of verapamil during labor or delivery has immediate or delayed adverse effects on the fetus, or whether it prolongs the duration of labor or increases the need for forceps delivery or other obstetric intervention. Such adverse experiences have not been reported in the literature, despite a long history of use of verapamil in Europe in the treatment of cardiac side effects of beta-adrenergic agonist agents used to treat premature labor.

### **8.3 Nursing Mothers**

Verapamil is excreted into human milk. In case studies where verapamil concentration in human milk was calculated, the nursing infant doses ranged from less than 0.01% to 0.1% of the mother's verapamil dose. Consider possible infant exposure when verapamil is administered to a nursing woman.

### **8.4 Pediatric Use**

Safety and effectiveness in pediatric patients have not been established.

### **8.5 Geriatric Use**

Clinical studies of verapamil hydrochloride extended-release capsules (PM) were not adequate to determine if subjects aged 65 or over respond differently from younger patients. Other reported clinical experience has not identified differences in response between the elderly and younger patients; however, greater sensitivity to verapamil hydrochloride extended-release capsules (PM) by some older individuals cannot be ruled out.

Aging may affect the pharmacokinetics of verapamil. Elimination half-life may be prolonged in the elderly [*see Clinical Pharmacology (12.3)*].

Verapamil is highly metabolized by the liver, and about 70% of the administered dose is excreted as metabolites in the urine. Clinical circumstances, some of which may be more common in the elderly, such as hepatic or renal impairment, should be considered [*see Use in Specific Populations (8.6, 8.7)*]. In general, lower initial doses of verapamil hydrochloride extended-release capsules (PM) may be warranted in the elderly [*see Dosage and Administration (2.1)*].

### **8.6 Impaired Hepatic Function**

Since verapamil is highly metabolized by the liver, consider lower dosages and closely monitor responses to the drug in patients with impaired hepatic function. Severe liver dysfunction prolongs the elimination half-life of immediate-release verapamil to about 14 to 16 hours; hence, approximately 30% of the dose given to patients with normal liver function should be administered to these patients. Monitor for abnormal prolongation of the PR interval or other signs of excessive pharmacologic effects [*see Overdosage (10)*]. In patients with hepatic cirrhosis, the influence of verapamil on digoxin pharmacokinetics is magnified. Verapamil may reduce total body clearance and extrarenal clearance of digitoxin by 27% and 29%, respectively.

### **8.7 Impaired Renal Function**

About 70% of an administered dose of verapamil is excreted as metabolites in the urine. Until further data are available, monitor these patients for abnormal prolongation of the PR interval or other signs of overdosage [*see Overdosage (10)*].

### **8.8 Attenuated (decreased) Neuromuscular Transmission**



It has been reported that verapamil decreases neuromuscular transmission in patients with Duchenne's muscular dystrophy, and that verapamil prolongs recovery from the neuromuscular blocking agent vecuronium and causes a worsening of myasthenia gravis. It may be necessary to decrease the dosage of verapamil when it is administered to patients with attenuated neuromuscular transmission.

## 10 OVERDOSAGE

There is no specific antidote for verapamil overdosage; treatment is supportive. Delayed pharmacodynamic consequences may occur with extended-release formulations, observe patients for at least 48 hours, preferably under continuous hospital care. Reported effects include hypotension, bradycardia, cardiac conduction defects, arrhythmias, hyperglycemia and decreased mental status. In addition, there have been literature reports of noncardiogenic pulmonary edema in patients taking large overdoses of verapamil (up to approximately 9 g).


In acute overdosage, consider gastrointestinal decontamination with cathartics and whole bowel irrigation. Calcium, inotropes (i.e., isoproterenol hydrochloride, dopamine hydrochloride and glucagon), atropine sulfate, vasopressors (i.e., norepinephrine and epinephrine), and cardiac pacing have been used with variable results to reverse hypotension and myocardial depression. In a few reported cases, overdose with calcium channel blockers that was initially refractory to atropine became more responsive to this treatment when the patients received large doses (close to 1 gram/hour for more than 24 hours) of calcium chloride.

Calcium chloride is preferred to calcium gluconate since it provides 3 times more calcium per volume. Asystole should be handled by the usual measures including cardiopulmonary resuscitation. Verapamil cannot be removed by hemodialysis.

## 11 DESCRIPTION

Verapamil hydrochloride is a calcium ion influx inhibitor (slow channel blocker or calcium ion antagonist). Verapamil hydrochloride extended-release capsules (PM) are available for oral administration as a 100 mg hard gelatin capsule (red opaque cap/white opaque body), a 200 mg hard gelatin capsule (red opaque cap/light orange opaque body), and as a 300 mg hard gelatin capsule (red opaque cap/red opaque body). Verapamil is administered as a racemic mixture of the R and S enantiomers.

The structural formula of verapamil hydrochloride is:



Chemical Name: Benzeneacetonitrile,  $\alpha$ -[3-[[2-(3,4-dimethoxyphenyl) ethyl]methylamino]propyl]-3,4-dimethoxy- $\alpha$ -(1-methylethyl)-, monohydrochloride, ( $\pm$ )-

Verapamil hydrochloride, USP is an almost white, crystalline powder, practically free of odor, with a bitter taste. It is soluble in water, chloroform and methanol. Verapamil hydrochloride is not structurally related to other cardioactive drugs.

In addition to verapamil hydrochloride, the verapamil hydrochloride extended-release capsules (PM) contain the following inactive ingredients: ammonium hydroxide, colloidal anhydrous silica, dibutyl sebacate, diethyl phthalate, ethylcellulose, hypromellose, maltodextrin, methacrylic acid copolymer, oleic acid, polyethylene glycol, povidone, silicon dioxide, sugar spheres, sodium lauryl sulfate and talc.

Each of the empty gelatin capsules for the 100 mg, 200 mg and 300 mg capsule strengths contain the following: D&C Red No. 28, D&C Red No. 33, D&C Yellow No. 10, FD&C Red No. 40, gelatin and titanium dioxide. In addition, the 200 mg empty gelatin capsule contains FD&C Yellow No. 6.

The imprinting ink contains the following: black iron oxide, D&C Yellow No. 10 Aluminum Lake, FD&C Blue No. 1 Aluminum Lake, FD&C Blue No. 2 Aluminum Lake, FD&C Red No. 40 Aluminum Lake, propylene glycol and shellac glaze.

### **System Components and Performance**

Verapamil hydrochloride extended-release capsules (PM) are designed for bedtime dosing, and it incorporates a delayed drug delivery system. This slow onset results in an average maximum plasma

concentration ( $C_{\max}$ ) of verapamil in the morning hours (approximately 9 to 12 hours following product administration). These bead-filled capsules provide for extended-release of the drug in the gastrointestinal tract. This delay is introduced by the release-controlling polymers applied to drug loaded beads. The release-controlling polymer is a combination of water soluble and water insoluble polymers. As water from the gastrointestinal tract comes into contact with the polymer coated beads, the water soluble polymer slowly dissolves and the drug diffuses through the resulting pores in the coating. The water insoluble polymer continues to act as a barrier, maintaining the controlled release of drug. The rate of release is essentially independent of posture and food. Multiparticulate systems such as verapamil capsules (PM) have been shown to be independent of gastrointestinal motility.

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

Verapamil is a calcium ion influx inhibitor (L-type calcium channel blocker or calcium channel antagonist). Verapamil exerts its pharmacologic effects by selectively inhibiting the transmembrane influx of ionic calcium into arterial smooth muscle as well as in conductile and contractile myocardial cells without altering serum calcium concentrations. Verapamil binding is voltage-dependent with affinity increasing as the vascular smooth muscle membrane potential is reduced. In addition, verapamil binding is frequency dependent and apparent affinity increases with increased frequency of depolarizing stimulus.

The L-type calcium channel is an oligomeric structure consisting of five putative subunits designated alpha-1, alpha-2, beta, tau and epsilon. Biochemical evidence points to separate binding sites for 1, 4-dihydropyridines, phenylalkylamines and the benzothiazepines (all located on the alpha-1 subunit). Although they share a similar mechanism of action, calcium channel blockers represent three heterogeneous categories of drugs with differing vascular-cardiac selectivity ratios.

### 12.2 Pharmacodynamics

#### Essential Hypertension

Verapamil produces its antihypertensive effect by a combination of vascular and cardiac effects. It acts as a vasodilator with selectivity for the arterial portion of the peripheral vasculature. As a result the systemic vascular resistance is reduced and usually without orthostatic hypotension or reflex tachycardia. Bradycardia (rate less than 50 beats/min) is uncommon. During isometric or dynamic exercise verapamil does not alter systolic cardiac function in patients with normal ventricular function.

Verapamil does not alter total serum calcium levels. However, one report has suggested that calcium levels above the normal range may alter the therapeutic effect of verapamil.

Verapamil regularly reduces the total systemic resistance (afterload) against which the heart works both at rest and at a given level of exercise by dilating peripheral arterioles.

#### Electrophysiologic Effects

Electrical activity through the AV node depends, to a significant degree, upon the transmembrane influx of extracellular calcium through the L-type (slow) channel. By decreasing the influx of calcium, verapamil prolongs the effective refractory period within the AV node and slows AV conduction in a rate-related manner.

Normal sinus rhythm is usually not affected, but in patients with sick sinus syndrome, verapamil may interfere with sinus-node impulse generation and may induce sinus arrest or sinoatrial block. Atrioventricular block can occur in patients without preexisting conduction defects [*see Warnings and Precautions (5.5)*].

Verapamil does not alter the normal atrial action potential or intraventricular conduction time, but depresses amplitude, velocity of depolarization and conduction in depressed atrial fibers. Verapamil may shorten the antegrade effective refractory period of the accessory bypass tract. Acceleration of ventricular rate and/or ventricular fibrillation has been reported in patients with atrial flutter or atrial fibrillation and a coexisting accessory AV pathway following administration of verapamil [*see Warnings and Precautions (5.4)*].

Verapamil has a local anesthetic action that is 1.6 times that of procaine on an equimolar basis. It is not known whether this action is important at the doses used in man.

#### Hemodynamics

Verapamil reduces afterload and myocardial contractility. In most patients, including those with organic cardiac disease, the negative inotropic action of verapamil is countered by reduction of afterload and cardiac index remains unchanged. During isometric or dynamic exercise, verapamil does not alter systolic cardiac function in patients with normal ventricular function. In patients with severe left ventricular dysfunction (e.g., pulmonary wedge pressure above 20 mm Hg or ejection fraction less than 30%), or in patients taking beta-adrenergic blocking agents or other cardiodepressant drugs, deterioration of ventricular function may occur [*see Drug Interactions (7.3)*].

## Pulmonary Function

Verapamil does not induce bronchoconstriction and, hence, does not impair ventilatory function.

Verapamil has been shown to have either a neutral or relaxant effect on bronchial smooth muscle.

### 12.3 Pharmacokinetics

Verapamil is administered as a racemic mixture of the R and S enantiomers. The systemic concentrations of R and S enantiomers, as well as overall bioavailability, are dependent upon the route of administration and the rate and extent of release from the dosage forms. Upon oral administration, there is rapid stereoselective biotransformation during the first pass of verapamil through the portal circulation.

#### Absorption

In a study in five subjects with oral immediate-release verapamil, the systemic bioavailability was from 33% to 65% for the R enantiomer and from 13% to 34% for the S enantiomer. Following oral administration of an immediately releasing formulation every 8 hours in 24 subjects, the relative systemic availability of the S enantiomer compared to the R enantiomer was approximately 13% following a single day's administration and approximately 18% following administration to steady-state. The degree of stereoselectivity of metabolism for verapamil hydrochloride extended-release capsules (PM) was similar to that for the immediate-release formulation. The R and S enantiomers have differing levels of pharmacologic activity. In studies in animals and humans, the S enantiomer has 8 to 20 times the activity of the R enantiomer in slowing AV conduction. In animal studies, the S enantiomer has 15 to 50 times the activity of the R enantiomer in reducing myocardial contractility in isolated blood-perfused dog papillary muscle, respectively, and twice the effect in reducing peripheral resistance. In isolated septal strip preparations from five patients, the S enantiomer was 8 times more potent than the R in reducing myocardial contractility. Dose escalation study data indicate that verapamil concentrations increase disproportionately to dose as measured by relative peak plasma concentrations ( $C_{max}$ ) or areas under the plasma concentration vs. time curves (AUC).

Consumption of a high fat meal just prior to dosing in the morning had no effect on the extent of absorption and a modest effect on the rate of absorption from verapamil hydrochloride extended-release capsules (PM). The rate of absorption was not affected by whether the volunteers were supine 2 hours after night-time dosing or nonsupine for 4 hours following morning dosing. Administering verapamil hydrochloride extended-release capsules (PM) in the morning increased the extent of absorption of verapamil and/or decreased the metabolism to norverapamil.

When the contents of the verapamil hydrochloride extended-release capsules (PM) were administered by sprinkling onto one tablespoonful of applesauce, the rate and extent of verapamil absorption were found to be bioequivalent to the same dose when administered as an intact capsule. Similar results were observed with norverapamil.

#### Distribution

Although some evidence of lack of dose linearity was observed for verapamil hydrochloride extended-release capsules (PM), this non-linearity was enantiomer specific, with the R enantiomer showing the greatest degree of nonlinearity.

**Table 3. Pharmacokinetic Characteristics of Verapamil Enantiomers After Administration of Escalating Doses of Verapamil Hydrochloride Extended-release Capsules (PM)**

	ISOMER	200	300	400
Dose Ratio		1	1.5	2
Relative $C_{max}$	R	1	1.89	2.34
	S	1	1.88	2.5
Relative AUC	R	1	1.67	2.34
	S	1	1.35	2.20

Racemic verapamil is released from verapamil hydrochloride extended-release capsules (PM) by diffusion following the gradual solubilization of the water soluble polymer. The rate of solubilization of the water soluble polymer produces a lag period in drug release for approximately 4 to 5 hours. The drug release phase is prolonged with the peak plasma concentration ( $C_{max}$ ) occurring approximately 11 hours after administration. Trough concentrations occur approximately 4 hours after bedtime dosing while the patient is sleeping. Steady-state pharmacokinetics were determined in healthy volunteers. Steady-state concentration is achieved by day 5 of dosing.

In healthy volunteers, following administration of verapamil hydrochloride extended-release capsules (PM) (200 mg per day), steady-state pharmacokinetics of the R and S enantiomers of verapamil is as follows: Mean  $C_{max}$  of the R isomer was 77.8 ng/ml and 16.8 ng/ml for the S isomer;  $AUC_{(0-24h)}$  of the R isomer was 1037 ng•h/ml and 195 ng•h/ml for the S isomer.

In general, bioavailability of verapamil is higher and half-life longer in older (> 65 yrs) subjects. Lean body weight also affects its pharmacokinetics inversely. It was not possible to observe a gender difference in the clinical trials of verapamil hydrochloride extended-release capsules (PM) due to the small sample size. However, there are conflicting data in the literature suggesting that verapamil

clearance decreased with age in women to a greater degree than in men.

### Metabolism and Excretion

Orally administered verapamil undergoes extensive metabolism in the liver. Verapamil is metabolized by O-demethylation (25%) and N-dealkylation (40%), and is subject to pre-systemic hepatic metabolism with elimination of up to 80% of the dose. The metabolism is mediated by hepatic cytochrome P450, and animal studies have implied that the mono-oxygenase is the specific isoenzyme of the P450 family. Thirteen metabolites have been identified in urine. Norverapamil enantiomers can reach steady-state plasma concentrations approximately equal to those of the enantiomers of the parent drug. For verapamil hydrochloride extended-release capsules (PM), the norverapamil R enantiomer reached steady-state plasma concentrations similar to the verapamil R enantiomer, but the norverapamil S enantiomer concentrations were approximately twice that of the verapamil S enantiomer concentrations. The cardiovascular activity of norverapamil appears to be approximately 20% that of verapamil. Approximately 70% of an administered dose is excreted as metabolites in the urine and 16% or more in the feces within 5 days. About 3% to 4% is excreted in the urine as unchanged drug.

R-verapamil is 94% bound to plasma albumin, while S-verapamil is 88% bound. In addition, R-verapamil is 92% and S-verapamil 86% bound to alpha-1 acid glycoprotein. In patients with hepatic insufficiency, metabolism of immediate-release verapamil is delayed and elimination half-life prolonged up to 14 to 16 hours because of the extensive hepatic metabolism [see *Use in Specific Populations* (8.6)]. In addition, in these patients there is a reduced first pass effect, and verapamil is more bioavailable. Verapamil clearance values suggest that patients with liver dysfunction may attain therapeutic verapamil plasma concentrations with one third of the oral daily dose required for patients with normal liver function.

After 4 weeks of oral dosing of immediate-release verapamil (120 mg q.i.d.), verapamil and norverapamil levels were noted in the cerebrospinal fluid with estimated partition coefficient of 0.06 for verapamil and 0.04 for norverapamil.

### Geriatric Use

The pharmacokinetics of verapamil GITS were studied after 5 consecutive nights of dosing 180 mg in 30 healthy young (19 to 43 years) vs. 30 healthy elderly (65 to 80 years) male and female subjects. Older subjects had significantly higher mean verapamil  $C_{max}$ ,  $C_{min}$  and  $AUC_{(0-24h)}$  compared to younger subjects. Older subjects had mean AUCs that were approximately 1.7 to 2 times higher than those of younger subjects as well as a longer average verapamil  $t_{1/2}$  (approximately 20 hr vs. 13 hr).

## 13 NONCLINICAL TOXICOLOGY

### 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

An 18-month toxicity study in rats, at a low multiple (6-fold) of the maximum recommended human dose, and not the maximum tolerated dose, did not suggest a tumorigenic potential. There was no evidence of a carcinogenic potential of verapamil administered in the diet of rats for 2 years at doses of 10, 35 and 120 mg/kg/day or approximately 1.3, 4.4 and 15 times, respectively, the maximum recommended human daily dose (400 mg/day or 8 mg/kg/day).

Verapamil was not mutagenic in the Ames test in five test strains at 3 mg per plate, with or without metabolic activation.

Studies in female rats at daily dietary doses up to 6.9 times (55 mg/kg/day) the maximum recommended human dose did not show impaired fertility. Effects on male fertility have not been determined.

### 13.2 Animal Toxicology and/or Pharmacology

In chronic animal toxicology studies verapamil caused lenticular and/or suture line changes at 30 mg/kg/day or greater and frank cataracts at 62.5 mg/kg/day or greater in the beagle dog but not in the rat. Development of cataracts due to verapamil has not been reported in man.

## 14 CLINICAL STUDIES

Verapamil hydrochloride extended-release capsules (PM) was evaluated in two placebo-controlled, parallel design, double-blind studies of patients with mild to moderate hypertension. In the clinical trials, 413 evaluable patients were randomized to either placebo, 100 mg, 200 mg, 300 mg or 400 mg and treated for up to 8 weeks. Verapamil hydrochloride extended-release capsules (PM) or placebo was given once daily between 9 pm and 11 pm (nighttime) and blood pressure changes were measured with 36-hour ambulatory blood pressure monitoring (ABPM). The results of these studies demonstrate that verapamil hydrochloride extended-release capsules (PM), at 200 mg, 300 mg and 400 mg, is a consistently and significantly more effective antihypertensive agent than placebo in reducing ambulatory blood pressures. Over this dose range, the placebo-subtracted net decreases in diastolic BP at trough (averaged over 6 to 10 pm) were dose related, and ranged from 3.8 to 10 mm Hg after 8 weeks of therapy. Although verapamil hydrochloride extended-release capsules (PM) 100 mg was not effective in reducing diastolic BP at trough when measured by ABPM, efficacy was demonstrated in reducing diastolic BP when measured manually at trough and peak and, from 6 am to 12 noon and over 24 hours

when measured by ABPM [see Dosage and Administration (2.1) for titration schedule]).

There were no apparent treatment differences between patient subgroups of different age (older or younger than 65 years), sex and race. For severity of hypertension, "moderate" hypertensives (mean daytime diastolic BP  $\geq 105$  mm Hg and  $\geq 114$  mm Hg) appeared to respond better than "mild" hypertensives (mean daytime diastolic BP  $\geq 90$  mm Hg and  $\leq 104$  mm Hg). However, sample size for the subgroup comparisons were limited.

## 16 HOW SUPPLIED/STORAGE AND HANDLING

Verapamil Hydrochloride Extended-Release Capsules (PM) are available containing 100 mg, 200 mg or 300 mg of verapamil hydrochloride, USP.

The 100 mg capsule is a hard-shell gelatin capsule with a red opaque cap and a white opaque body filled with white to off-white beads. The capsule is axially printed with **MYLAN** over **6201** in black ink on both the cap and the body. They are available as follows:

NDC 54868-2211-0 bottles of 30 capsules

The 200 mg capsule is a hard-shell gelatin capsule with a red opaque cap and a light orange opaque body filled with white to off-white beads. The capsule is axially printed with **MYLAN** over **6202** in black ink on both the cap and the body. They are available as follows:

NDC 54868-6124-0 bottles of 10 capsules

NDC 54868-6124-1 bottles of 30 capsules

The 300 mg capsule is a hard-shell gelatin capsule with a red opaque cap and a red opaque body filled with white to off-white beads. The capsule is axially printed with **MYLAN** over **6203** in black ink on both the cap and the body. They are available as follows:

NDC 54868-5841-0 bottles of 20 capsules

NDC 54868-5841-2 bottles of 30 capsules

NDC 54868-5841-1 bottles of 60 capsules

**Store at 20° to 25°C (68° to 77°F). [See USP for Controlled Room Temperature.]**

**Protect from moisture.**

Dispense in a tight, light-resistant container as defined in the USP using a child-resistant closure.

## 17 PATIENT COUNSELING INFORMATION

**THE CONTENTS OF THE VERAPAMIL HYDROCHLORIDE EXTENDED-RELEASE CAPSULES (PM) SHOULD NOT BE CRUSHED OR CHEWED. VERAPAMIL HYDROCHLORIDE EXTENDED-RELEASE CAPSULES (PM) ARE TO BE SWALLOWED WHOLE OR THE ENTIRE CONTENTS OF THE CAPSULE SPRINKLED ONTO APPLESAUCE [see Dosage and Administration (2.2)].**

When the sprinkle method of administration is prescribed, explain the details of the proper technique to the patient [see Dosage and Administration (2.2)].


Mylan Pharmaceuticals Inc.  
Morgantown, WV 26505

REVISED JUNE 2010  
VPMLC:R3

### **Relabeling and Repackaging by:**

Physicians Total Care, Inc.  
Tulsa, Oklahoma 74146

**PRINCIPAL DISPLAY PANEL - 100 mg**



**Verapamil HCL  
Extended-release  
Capsules (PM)  
100 mg  
Controlled-onset**

**Rx only**

Each capsule contains:

Verapamil

hydrochloride, USP 100 mg

Dispense in a tight, light-resistant container as defined in the USP using a child-resistant closure.

Keep container tightly closed.

**Keep this and all medication out of the reach of children.**


**Store at 20° to 25°C (68° to 77°F).**

**[See USP Controlled Room Temperature.]**

**Protect from moisture.**

**Usual Adult Dosage:** See accompanying prescribing information.

**PRINCIPAL DISPLAY PANEL - 200 mg**



**VERAPAMIL  
HYDROCHLORIDE  
EXTENDED-RELEASE  
CAPSULES (PM)  
200 mg  
CONTROLLED-ONSET**

**(Rx only)**

Each capsule contains:

Verapamil

hydrochloride, USP 200 mg

Dispense in a tight, light-resistant container as defined in the USP using a child-resistant closure.

Keep container tightly closed.

**Keep this and all medication out of the reach of children.**


**Store at 20° to 25°C (68° to 77°F).**

**[See USP Controlled Room Temperature.]**

**Protect from moisture.**

**Usual Adult Dosage:** See accompanying prescribing information.

**PRINCIPAL DISPLAY PANEL - 300 mg**



**VERAPAMIL  
HYDROCHLORIDE  
EXTENDED-RELEASE  
CAPSULES (PM)  
300 mg  
CONTROLLED-ONSET**

**(Rx only)**

Each capsule contains:  
Verapamil  
hydrochloride, USP 300 mg

Dispense in a tight, light-resistant  
container as defined in the USP  
using a child-resistant closure.

Keep container tightly closed.

**Keep this and all medication  
out of the reach of children.**

**Store at 20° to 25°C (68° to 77°F).**  
**[See USP Controlled Room  
Temperature.]**

**Protect from moisture.**

**Usual Adult Dosage:** See accom-  
panying prescribing information.

VERAPAMIL HYDROCHLORIDE			
verapamil hydrochloride capsule, extended release			
Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:54868-2211(NDC:0378-6201)
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
VERAPAMIL HYDROCHLORIDE (UNII: V3888OEY5R) (VERAPAMIL - UNII:CJ0O37KU29)		VERAPAMIL HYDROCHLORIDE	100 mg
Inactive Ingredients			
Ingredient Name			Strength
AMMONIA (UNII: 5138Q19F1X)			
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)			
DIBUTYL SEBACATE (UNII: 4W5IH7FLNY)			
DIETHYL PHTHALATE (UNII: UF064M00AF)			
ETHYLCELLULOSES (UNII: 7Z8S9VYZ4B)			
HYPROMELLOSES (UNII: 3NXW29V3WO)			
MALTODEXTRIN (UNII: 7CVR7L4A2D)			
METHACRYLIC ACID - METHYL METHACRYLATE COPOLYMER (1:1) (UNII: 74G4R6TH13)			
METHACRYLIC ACID - METHYL METHACRYLATE COPOLYMER (1:2) (UNII: 5KY68S2577)			
OLEIC ACID (UNII: 2UMI9U37CP)			
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)			

POVIDONE (UNII: FZ989GH94E)				
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
SODIUM LAURYL SULFATE (UNII: 368GB5141J)				
TALC (UNII: 7SEV7J4R1U)				
D&C RED NO. 28 (UNII: 767IP0Y5NH)				
D&C RED NO. 33 (UNII: 9DBA0SBB0L)				
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)				
FD&C RED NO. 40 (UNII: WZB9127XOA)				
GELATIN (UNII: 2G86QN327L)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
FERROUS FERRIC OXIDE (UNII: XM0M87F357)				
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)				
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
SHELLAC (UNII: 46N107B71O)				
Product Characteristics				
Color	RED (red opaque) , WHITE (white opaque)		Score	no score
Shape	CAPSULE		Size	18 mm
Flavor			Imprint Code	MYLAN;6201
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:54868-2211-0	30 in 1 BOTTLE		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
ANDA	ANDA078306		08/07/2008	

VERAPAMIL HYDROCHLORIDE

verapamil hydrochloride capsule, extended release

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:54868-6124(NDC:0378-6202)
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
VERAPAMIL HYDROCHLORIDE (UNII: V3888OEY5R) (VERAPAMIL - UNII:CJ0O37KU29)	VERAPAMIL HYDROCHLORIDE	200 mg

Inactive Ingredients	
Ingredient Name	Strength
AMMONIA (UNII: 5138Q19F1X)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
DIBUTYL SEBACATE (UNII: 4W5IH7FLNY)	
DIETHYL PHTHALATE (UNII: UF064M00AF)	
ETHYLCELLULOSES (UNII: 7Z8S9VYZ4B)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
METHACRYLIC ACID - METHYL METHACRYLATE COPOLYMER (1:1) (UNII: 74G4R6TH13)	
METHACRYLIC ACID - METHYL METHACRYLATE COPOLYMER (1:2) (UNII: 5KY68S2577)	
OLEIC ACID (UNII: 2UMI9U37CP)	
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	



TALC (UNII: 7SEV7J4R1U)					
D&C RED NO. 28 (UNII: 767IP0Y5NH)					
D&C RED NO. 33 (UNII: 9DBA0SBB0L)					
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)					
FD&C RED NO. 40 (UNII: WZB9127XOA)					
GELATIN (UNII: 2G86QN327L)					
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)					
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)					
FERROSO FERRIC OXIDE (UNII: XM0M87F357)					
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)					
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)					
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)					
SHELLAC (UNII: 46N107B71O)					
Product Characteristics					
Color	RED (red opaque) , ORANGE (light orange opaque)			Score	no score
Shape	CAPSULE			Size	22mm
Flavor				Imprint Code	MYLAN;6202
Contains					
Packaging					
#	Item Code	Package Description	Marketing Start Date		Marketing End Date
1	NDC:54868-6124-0	10 in 1 BOTTLE			
2	NDC:54868-6124-1	30 in 1 BOTTLE			
Marketing Information					
Marketing Category	Application Number or Monograph Citation		Marketing Start Date		Marketing End Date
ANDA	ANDA078306		05/17/2010		

VERAPAMIL HYDROCHLORIDE

verapamil hydrochloride capsule, extended release

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:54868-5841(NDC:0378-6203)
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
VERAPAMIL HYDROCHLORIDE (UNII: V3888OEY5R) (VERAPAMIL - UNII:CJ0O37KU29)		VERAPAMIL HYDROCHLORIDE	300 mg
Inactive Ingredients			
Ingredient Name			Strength
AMMONIA (UNII: 5138Q19F1X)			
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)			
DIBUTYL SEBACATE (UNII: 4W5IH7FLNY)			
DIETHYL PHTHALATE (UNII: UF064M00AF)			
ETHYLCELLULOSES (UNII: 7Z8S9VYZ4B)			
HYPROMELLOSES (UNII: 3NXW29V3WO)			
MALTODEXTRIN (UNII: 7CVR7L4A2D)			
METHACRYLIC ACID - METHYL METHACRYLATE COPOLYMER (1:1) (UNII: 74G4R6TH13)			
METHACRYLIC ACID - METHYL METHACRYLATE COPOLYMER (1:2) (UNII: 5KY68S2577)			
OLEIC ACID (UNII: 2UMI9U37CP)			
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)			
POVIDONE (UNII: FZ989GH94E)			
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)			
SODIUM LAURYL SULFATE (UNII: 368GB5141J)			

TALC (UNII: 7SEV7J4R1U)				
D&C RED NO. 28 (UNII: 767IP0Y5NH)				
D&C RED NO. 33 (UNII: 9DBA0SBB0L)				
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)				
FD&C RED NO. 40 (UNII: WZB9I27XOA)				
GELATIN (UNII: 2G86QN327L)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
FERROSO FERRIC OXIDE (UNII: XM0M87F357)				
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)				
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
SHELLAC (UNII: 46N107B71O)				
Product Characteristics				
Color	RED (red opaque)	Score	no score	
Shape	CAPSULE	Size	23mm	
Flavor		Imprint Code	MYLAN;6203	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:54868-5841-0	20 in 1 BOTTLE		
2	NDC:54868-5841-1	60 in 1 BOTTLE		
3	NDC:54868-5841-2	30 in 1 BOTTLE		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA078306	12/20/2007		

<b>Labeler</b> - Physicians Total Care, Inc. (194123980)
--

<b>Establishment</b>			
Name	Address	ID/FEI	Business Operations
Physicians Total Care, Inc.		194123980	relabel, repack