

**COLD AND FLU RELIEF- acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride liquid**  
**Walgreens**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Drug Facts**

**Active ingredients (in each 15 mL)**

**Acetaminophen 325 mg**

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

**Purpose**

**Pain reliever/fever reducer**

Cough suppressant

Nasal decongestant

**Uses**

- temporarily relieves common cold and flu symptoms:
  - minor aches and pains
  - headache
  - sore throat
  - nasal congestion
  - fever
  - cough due to minor throat and bronchial irritation

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- adult take more than 4 doses (30 mL each) of acetaminophen in 24 hours, which is the maximum daily amount
- child takes more than 4 doses (15 mL each) in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blistere
- rash.

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- a sodium-restricted diet
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough as occurs with smoking, asthma, or emphysema

### **Ask a doctor or pharmacist before use if user is**

taking the blood thinning drug warfarin

### **When using this product**

**do not exceed recommended dosage.**

### **Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse, or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use.

## Keep out of reach of children.

**Overdose warning:** Taking more than the recommended dose can cause serious liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults and for children even if you do not notice any signs or symptoms

## Directions

- do not take more than directed (see overdose warning)
- do not take more than 4 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device
- mL = milliliter
- keep dosing cup with product

adults and children 12 years and over	30 mL every 4 hours
children 6 to under 12 years	15 mL every 4 hours
children 4 to under 6 years	ask a doctor
children under 4 years	do not use

- **when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing**

## Other information

- each 15 mL contains: sodium 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

## Inactive ingredients

citric acid, FD&C Yellow #6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol sucralose, xanthan gum

## Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

## Principal Display Panel

Compare to the active ingredients in Vicks® DayQuil® Cold & Flu††

DAYTIME • NON-DROWSY

## Cold & Flu

ACETAMINOPHEN 325 mg / PAIN RELIEVER / FEVER REDUCER

DEXTROMETHORPHAN HBr 10 mg / COUGH SUPPRESSANT

PHENYLEPHRINE HCl 5 mg / NASAL CONGESTION

Multi-Symptom

- Relieves aches, fever & sore throat , cough & nasal congestant
- For ages 6 years & over
- Alcohol free
- Antihistamine free

FL OZ (mL)

††This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® and DayQuil® are registered trademarks of The Procter & Gamble Company.

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.**

DISTRIBUTED BY: WALGREEN CO.

200 WILMOT RD., DEERFIELD, IL 60015

**Product Label**

## Drug Facts

### Active ingredients (in each 15 mL) Purposes

Dextromethorphan HBr 10 mg..... Cough suppressant  
Phenylephrine HCl 5 mg ..... Nasal decongestant

**Uses** ■ temporarily relieves common cold and flu symptoms  
■ minor aches and pains ■ headache ■ sore throat  
■ nasal congestion ■ fever  
■ cough due to minor throat and bronchial irritation

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if: ■ adult takes more than 4 doses (30 mL each) of acetaminophen in 24 hours, which is the maximum daily amount ■ child takes more than 4 doses (15 mL each) in 24 hours ■ taken with other drugs containing acetaminophen ■ adult has 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away.  
**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

← PEEL HERE **Drug Facts** (continued under label)

NDC 0363-9974-08

**Walgreens**

Compare to the active ingredients  
in Vicks® DayQuil® Cold & Flu\*



## DAYTIME • NON-DROWSY Cold & Flu

**ACETAMINOPHEN 325 mg / PAIN RELIEVER /  
FEVER REDUCER  
DEXTROMETHORPHAN HBr 10 mg / COUGH SUPPRESSANT  
PHENYLEPHRINE HCl 5 mg / NASAL DECONGESTANT**

### Multi-Symptom

- Relieves aches, fever, sore throat, cough & nasal congestion
- For ages 6 years & over
- Antihistamine free
- Alcohol free

8 FL OZ (237 mL)

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL  
AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING

### PARENTS:

Learn about teen medicine abuse  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

\*Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.

†This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® and DayQuil® are registered trademarks of The Procter & Gamble Company.

DISTRIBUTED BY:

**WALGREEN CO.**  
200 WILMOT RD.,  
DEERFIELD, IL 60015  
**100% SATISFACTION  
GUARANTEED**  
[walgreens.com](http://walgreens.com)  
©2022 Walgreen Co.

PLD-B380D LB009015



W3ORG1022-F  
REV-0323

ITEM 816520 W11042-0423-F



### Drug Facts (continued)

■ If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have** ■ liver disease  
■ heart disease ■ high blood pressure ■ thyroid disease  
■ diabetes ■ a sodium-restricted diet ■ trouble urinating due to an enlarged prostate gland ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema ■ cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.**

**When using this product, do not exceed recommended dosage.**

**Stop use and ask a doctor if** ■ nervousness, dizziness, or sleeplessness occur ■ pain, nasal congestion, or cough gets worse, or lasts more than 5 days (children) or 7 days (adult) ■ fever gets worse, or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use.**

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended

### Drug Facts (continued)

dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** ■ do not take more than directed (see Overdose warning) ■ do not take more than 4 doses in any 24-hour period ■ measure only with dosing cup provided. Do not use any other dosing device.  
■ keep dosing cup with product ■ mL = milliliter

adults and children 12 years and over	30 mL every 4 hours
children 6 to under 12 years	15 mL every 4 hours
children 4 to under 6 years	ask a doctor
children under 4 years	do not use

■ when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

### Other information

■ each 15 mL contains: sodium 12 mg  
■ store between 20-25°C (68-77°F). Do not refrigerate.

**Inactive ingredients** citric acid, FD&C yellow #6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose, xanthan gum

### Questions or comments?

Call 1-877-753-3535 Monday-Friday 9AM-5PM EST

## WALGREENS Daytime Cold & Flu

## COLD AND FLU RELIEF

acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0363-9974
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X)	
<b>SODIUM CITRATE</b> (UNII: 1Q73Q2JULR)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-9974-08	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	09/30/2022	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	09/30/2022	

**Labeler** - Walgreens (008965063)

Revised: 4/2023

Walgreens