MEIJER KIDS TEAR FREE SPF 50 - avobenzone, homosalate, octisalate, octocrylene spray Meijer Distribution, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredients

Avobenzone 3.0%, Homosalate 10.0%, Octisalate 5.0%, Octocrylene 6.0%

Purpose

Sunscreen

Uses

- · helps prevent sunburn
- if used as directed with other sun protection measures (see **Directions**), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Flammable: Do not puncture or incinerate. Contents under pressure. Do not store at temperature above 120^{0} F. Do not use in the presence of a flame or spark. Keep away from sources of ignition - No smoking.

Do not use

on damaged or broken skin

When using this product

- Keep away from face to avoid breathing it.
- Keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor

if rash occurs

Keep out of reach of children.

- If product is swallowed, get medical help or contact a Poison Control center right away.
- Intentional misuse by deliberately concentrating and inhaling the contents can be harmful or fatal.
- May stain some fabrics.

Directions

- hold container 4-6 inches from the skin to apply
- spray liberally and spread evenly by hand 15 minutes before sun exposure
- **Do not spray directly on face**. Spray on hands then apply to face.
- do not apply in windy conditions

- use in well-ventilated area
- reapply:
- after 80 minutes of swimming or sweating
- immediately after towel drying
- at least every 2 hours
- **Sun Protection Measures**. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun production measures including:
- limit time in the sun, especially from 10 a.m. to 2 p.m.
- wear long-sleeved shirts, pants, hats, and sunglasses
- children under 6 months of age: Ask a Doctor

Inactive ingredients

Isobutane, Water, Isohexadecane, Ethylhexyl Benzoate, Cetyl PEG/PPG-10/1 Dimethicone, Dibutyl Adipate, Lauryl PEG-8 Dimethicone, Ethylhexyl Methoxycrylene, Sodium Chloride, PEG-8, Caprylyl Glycol, Glycerin, Sodium Citrate, PEG-8 Dimethicone, Aloe Barbadensis Leaf Juice, Octyldodecanol, Retinyl Palmitate, Silica, Tocopheryl Acetate, Ascorbic Acid, Sodium Propoxyhydroxypropyl Thiosulfate Silica, Phenoxyethanol, Methylparaben, Butylparaben, Ethylparaben, Isobutylparaben, Propylparaben.

Other information

protect this product from excessing heat and direct sun

Questions or comments?

Call 1-800-697-9868 Mon-Fri

Principal Display Panel

NDC: 41250-928-55

meijer

tear free

Kids Sunscreen

continuous spray

pediatrician tested

UVA/UVB protection

50 +

Broad Spectrum

SPF 50+

Water Resistant

(80 minutes)

NET WT 6 OZ. (170 g)





pediatrition tested **UVA/UVB** protection

Broad Spectrum SPF 50+

Water Resistant (80 Minutes)

NET WT 6 OZ. (170 g)

Drug Facts

Active ingredients

Avobenzone 3.0%, Homosalate 10.0%, Octisalate 5.0%

Uses • helps prevent sunbum • if used as directed with other sun protection measures (see Directions), decreases the risk of skin cancer and early skin aging caused by the sun

Purpose

Warnings
For external use only
Hammable: Do not puncture or incinerate. Contents under pressure. Do not store at temperatures above 120°F. Do not use in the presence of a flame or spark. Keep away from sources of ignition-No smoking. **Do not use** on damaged or broken skin

When using this product . Keep away from face to avoid breathing it Keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs

Keep out of the reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away Intentional misuse by deliberately concentrating and inhaling the contents can be harmful or fatal. May stain some fabrics.

Directions • hold container 4-6 inches from the skin to apply . spray liberally and spread evenly by hand 15 minutes before sun exposure • Do not spray directly on face. Spray on hands then apply to face. . do not apply in windy conditions . use in well-ventilated area · reapply: · after 80 minutes of swimming or sweating · immediately after towel drying • at least every 2 hours

Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including: • limit time in the sun, especially from 10 a.m. - 2 p.m. • wear long-sleeved shirts, pants, hats and sunglasses • children under 6 months of age: Ask a doctor

Inactive ingredients Isobutane, Water, Isobexadecane, Ethylhex Benzoate, Cetyl PEG/PPG-10/1 Dimethicone, Dibutyl Adipate, Lauryl PEG-Dimethicone, Ethylhexyl Methoxycrylene, Sodium Chloride, PEG-8, Phenoxyethanol, Caprylyl Glycol, Glycerin, Sodium Citrate, Methylparaben, Butylparaben, Ethylparaben, Isobutylparaben, Propylparaben, PEG-8 Dimethicone, Aloe Barbadensis Leaf Juice, Octyldodecanol, Retinyl Palmitate, Silica, Tocopheryl Acetate, Ascorbic Acid, Sodium Propoxyhydroxypropyl Thiosulfate Silica.

Other information protect this product from excessive heat and direct sur

MEIJER KIDS TEAR FREE SPF 50

avobenzone, homosalate, octisalate, octocrylene spray

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41250-928
--------------	----------------	--------------------	---------------

TOPICAL **Route of Administration**

Active Ingredient/Active Moiety

redice ingredient redicery			
Ingredient Name	Basis of Strength	Strength	
AVOBENZONE (UNII: G63QQF2NOX) (AVOBENZONE - UNII:G63QQF2NOX)	AVOBENZONE	3 g in 100 g	
HOMOSALATE (UNII: V06SV4M95S) (HOMOSALATE - UNII:V06SV4M95S)	HOMOSALATE	10 g in 100 g	
OCTISALATE (UNII: 4X49 Y0596W) (OCTISALATE - UNII:4X49 Y0596W)	OCTISALATE	5 g in 100 g	
OCTOCRYLENE (UNII: 5A68WGF6WM) (OCTOCRYLENE - UNII:5A68WGF6WM)	OCTOCRYLENE	6 g in 100 g	

Inactive Ingredients

Ingredient Name	Strength

ISOBUTANE (UNII: BXR49TP611)	
WATER (UNII: 059QF0KO0R)	
ISOHEXADECANE (UNII: 9 18 X10 UF1E)	
2-ETHYLHEXYL BENZOATE (UNII: R63ZWW1A13)	
DIBUTYL ADIPATE (UNII: F4K100DXP3)	
ETHYLHEXYL METHO XYCRYLENE (UNII: S3KFG6Q5X8)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
CAPRYLYL GLYCOL (UNII: 00 YIU5438 U)	
GLYCERIN (UNII: PDC6A3C0OX)	
SO DIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
PEG-8 DIMETHICONE (UNII: GIA7T764OD)	
ALOE VERA LEAF (UNII: ZY81Z83H0X)	
OCTYLDODECANOL (UNII: 46 1N1O 6 14Y)	
VITAMIN A PALMITATE (UNII: 1D1K0N0 VVC)	
SILICON DIO XIDE (UNII: ETJ7Z6 XBU4)	
.ALPHATO COPHEROL ACETATE (UNII: 9E8X80D2L0)	
ASCORBIC ACID (UNII: PQ6CK8PD0R)	
SODIUM PROPOXYHYDROXYPROPYL THIOSULFATE SILICA (UNII: 208G222332)	
PHENO XYETHANOL (UNII: HIE492ZZ3T)	
METHYLPARABEN (UNII: A218 C7HI9 T)	
BUTYLPARABEN (UNII: 3QPI1U3FV8)	
ETHYLPARABEN (UNII: 14255EXE39)	
ISOBUTYLPARABEN (UNII: 0QQJ25X58G)	
PROPYLPARABEN (UNII: Z8 IX2SC1OH)	

l	Pack	Packaging			
	#]	Item Code	Package Description	Marketing Start Date	Marketing End Date
	1 ND 55	C:41250-928-	170 g in 1 BOTTLE, SPRAY; Type 0: Not a Combination Product	04/06/2016	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part352	03/11/2016	

Labeler - Meijer Distribution, Inc. (006959555)

Revised: 3/2016 Meijer Distribution, Inc.