

**ASPIRIN 325- aspirin tablet, delayed release**  
**Walgreen Company**

-----  
**Walgreens 44-691**

***Active ingredient (in each caplet)***

Aspirin 325 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever

***Uses***

for the temporary relief of minor aches and pains or as recommended by your doctor.  
**Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.**

***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- shock
- hives
- facial swelling
- asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- are age 60 or older
- take more or for a longer time than directed
- take a blood thinning (anticoagulant) or steroid drug
- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product

**Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug for

- gout
- diabetes
- arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - vomit blood
  - feel faint
  - have bloody or black stools
  - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

***Directions***

- **do not take more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 caplets every 4 hours not to exceed 12 caplets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

***Other information***

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

## ***Inactive ingredients***

cellulose, colloidal anhydrous silica, corn starch, D&C yellow #10, FD&C yellow #6, hypromellose, methacrylic acid, polydextrose, polyethylene glycol, shellac wax, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate

## ***Questions or comments?***

**1-800-426-9391**

## ***Principal display panel***

### ***Walgreens***

Compare to the active ingredient in  
Bayer® Regular Strength Aspirin 325 mg  
Safety Coated Caplets††

**WALGREENS PHARMACIST RECOMMENDED†**

NDC 0363-0691-12

**Aspirin 325**  
**325 mg / PAIN RELIEVER (NSAID)**

**Regular Strength**

**Enteric-Coated**

ACTUAL SIZE

**100**  
CAPLETS

**HEART**  
**HEALTH\*\***

\*\*Talk to your doctor or other healthcare provider  
before using this product for your heart.

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED**  
**SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

†Our pharmacists recommend the  
Walgreens brand. We invite you to  
compare to national brands.

††This product is not manufactured or  
distributed by Bayer AG, owner of the  
registered trademark Bayer® Regular  
Strength Aspirin 325 mg Safety  
Coated Caplets.

50844

ORG012269112

DISTRIBUTED BY: **WALGREEN CO.**  
**200 WILMOT RD., DEERFIELD, IL 60015**

**Walgreens**  
**100% SATISFACTION**  
**GUARANTEED**

**walgreens.com**  
 ©2021 Walgreen Co.

**No Print Area**  
 Lot no. & Exp. Date



**ACTUAL SIZE**

**HEART HEALTH**

**Compare to the active ingredient in Bayer® Regular Strength Aspirin 325 mg Safety Coated Caplets\***

**Aspirin 325**  
 325 mg / PAIN RELIEVER (NSAID)    Regular Strength    Enteric-Coated

**100**  
CAPLETS

**\*\*Talk to your doctor or other healthcare provider before using this product for your heart.**

**Walgreens**

Compare to the active ingredient in Bayer® Regular Strength Aspirin 325 mg Safety Coated Caplets\*



NDC 0363-0691-12



how2recycle.info  
 PAPER BOX PLASTIC BOTTLE

**Drug Facts**

**Active ingredient (in each caplet)**  
 Aspirin 325 mg (NSAID)\*  
\*nonsteroidal anti-inflammatory drug

**Purpose**  
 Pain reliever

**Uses**  
 For the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.

**Warnings**  
 Reye's syndrome. Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:  
 ■ shock  
 ■ hives  
 ■ facial swelling  
 ■ asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  
 ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ are age 60 or older  
 ■ take more or for a longer time than directed  
 ■ take a blood thinning (anticoagulant) or steroid drug  
 ■ have had stomach ulcers or bleeding problems  
 ■ have 3 or more alcoholic drinks every day while using this product

**Drug Facts (continued)**

**Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

**Ask a doctor or pharmacist before use if you are taking a prescription drug for**

- gout ■ diabetes ■ arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding: ■ vomit blood ■ feel faint ■ have bloody or black stools ■ have stomach pain that does not get better ■ ringing in the ears or a loss of hearing occurs ■ pain gets worse or lasts more than 3 days ■ fever gets worse or lasts more than 10 days ■ redness or swelling is present ■ new symptoms occur

These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

**Drug Facts (continued)**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
 ■ do not take more than directed  
 ■ drink a full glass of water with each dose  
 ■ adults and children 12 years and over: take 1 or 2 caplets every 4 hours not to exceed 12 caplets in 24 hours unless directed by a doctor  
 ■ children under 12 years: do not use unless directed by a doctor

**Other information**

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

**Inactive ingredients** cellulose, colloidal anhydrous silica, corn starch, D&C yellow #10, FD&C yellow #6, hypromellose, methacrylic acid, polydextrose, polyethylene glycol, stearic wax, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate

**Questions or comments? 1-800-426-9391**

Your pharmacists recommend the Walgreens brand. We invite you to compare to national brands.  
 \*This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Bayer® Regular Strength Aspirin 325 mg Safety Coated Caplets.

50844    ORG012269112

DISTRIBUTED BY: **WALGREEN CO.**  
 200 WILMOT RD., DEERFIELD, IL 60015

**Walgreens**

**100% SATISFACTION**  
**GUARANTEED**

**walgreens.com**  
 ©2021 Walgreen Co.

ITEM 988239    W00000-0000-0  
 REV0322

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**



3 1917 119317 5

B-2201-691-12-HR  
 ORG012269112

**Walgreens 44-691**

# ASPIRIN 325

aspirin tablet, delayed release

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0363-0691
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

## Inactive Ingredients

Ingredient Name	Strength
POWDERED CELLULOSE (UNII: SMD1X3XO9M)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
METHACRYLIC ACID (UNII: 1CS02G8656)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SHELLAC (UNII: 46N107B71O)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

## Product Characteristics

<b>Color</b>	yellow	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	15mm
<b>Flavor</b>		<b>Imprint Code</b>	691
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0691-12	1 in 1 CARTON	07/01/2017	
1		100 in 1 BOTTLE, PLASTIC; Type 0: Not a		

Combination Product			
<b>Marketing Information</b>			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	07/01/2017	

**Labeler** - Walgreen Company (008965063)

<b>Establishment</b>			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(0363-0691)

<b>Establishment</b>			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(0363-0691)

<b>Establishment</b>			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(0363-0691)

<b>Establishment</b>			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(0363-0691)

<b>Establishment</b>			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(0363-0691)

<b>Establishment</b>			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(0363-0691)