

**IBUPROFEN- ibuprofen tablet**  
**The Kroger Co.**

-----  
**DRUG FACTS**

**Active Ingredient (in each caplet)**

Ibuprofen USP, 200 mg (NSAID)\*\*

\*\*nonsteroidal anti-inflammatory drug

**Purpose**

Pain Reliever/ Fever Reducer

**Uses**

- temporarily relieves minor aches and pains due to:
- Headache
- muscular aches
- minor pain of arthritis
- toothache
- backache
- the common cold
- menstrual cramps
- temporarily reduces fever

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or non-prescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks everyday while using this product
- take more or for a longer time than directed.

**Do not use**

- if you ever had an allergic reaction to any other pain reliever/fever reducer

- right before or after heart surgery

### **Ask a doctor before use if**

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic.

### **Ask a doctor or pharmacist before use if you are**

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug.

### **When using this product**

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed.

### **Stop use and ask a doctor if**

- You experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
  - Pain gets worse or last more than 10 days
  - Fever gets worse or last more than 3 days
  - Redness or swelling is present in the painful area
  - Any new symptoms appear

### **If pregnant or breast-feeding**

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

### **Directions**

- **do not take more than directed**
- **the smallest effective dose should be used**
- do not take longer than 10 days, unless directed by a doctor (see **Warnings**)
- Adults and children 12 years and older:
  - take 1 caplet every 4 to 6 hours while symptoms persist
  - if pain or fever does not respond to 1 caplet, 2 caplets may be used.
  - do not exceed 6 caplets in 24 hours, unless directed by a doctor
- Children under 12 years: ask a doctor
- **do not use** in children under 4 years of age

### **Other information**

- store between 20°- 25° C (68°- 77° F)
- read all warnings and directions before use.
- do not use if inner seal under bottle cap imprinted with "Sealed for your protection" is broken or missing

### **Inactive Ingredients**

Anhydrous Lactose, Carnauba Wax, Colloidal Silicon Dioxide, Hypromellose, Iron Oxide Red, Magnesium Stearate, Microcrystalline Cellulose, Polydextrose, Polyethylene Glycol, Povidone, Sodium Lauryl Sulfate, Sodium Starch Glycolate, Starch (Corn), Titanium Dioxide

### **Questions or comments?**

Call **1-800-632-6900**

### **Principal Display Panel**

Compare to the active ingredient of ADVIL® †See back panel

SEE NEW WARNINGS INFORMATION

Ibuprofen Tablets USP, 200 mg

Pain Reliever/ Fever Reducer (NSAID)\*\*

\*\*CAPSULE SHAPED TABLETS

†Advil® is a registered trademark of Wyeth corporation, Madison, NJ 07940. Wyeth corporation is not affiliated with the Kroger Co. or this product

**DISTRIBUTED BY THE KROGER CO.**

CINCINNATI, OHIO 45202

### **Product Label**

**COMPARE TO active ingredient of ADVIL®** †See back panel

NDC 30142-167-05

**SEE NEW WARNINGS INFORMATION**

**Kroger**  
QUALITY GUARANTEED

**Ibuprofen**  
Tablets, USP 200 mg

**Pain Reliever/Fever Reducer (NSAID)\*\***

**Our Pharmacists Recommend**

**500 CAPLETS\***  
\*capsule-shaped tablets



**Drug Facts**

**Active ingredient (in each caplet)**  
Ibuprofen USP 200 mg (NSAID)\*\* .....Pain reliever/fever reducer  
\*\* nonsteroidal anti-inflammatory drug

**Uses**

- temporarily relieves minor aches and pains due to: ■ headache
- muscular aches ■ minor pain of arthritis ■ toothache ■ backache
- the common cold ■ menstrual cramps ■ temporarily reduces fever

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chances are higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery

**Ask a doctor before use if**

- you have problems or serious side effects from taking pain relievers or fever reducers ■ the stomach bleeding warning applies to you ■ you have a history of stomach problems such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you have asthma ■ you are taking a diuretic

**Drug Facts** (continued under label)

†Myliv® is a registered trademark of Wyeth Corporation, Madison, NJ 07940. Wyeth Corporation is not affiliated with The Kroger Co. or this product.

DISTRIBUTED BY THE KROGER CO.  
CINCINNATI, OHIO 45202

**QUALITY GUARANTEED**  
If you are not completely satisfied with this product, return it for a refund or replacement.

Comments or questions? [www.Kroger.com](http://www.Kroger.com)

PLD-A LB000066

Lot No.:  
Exp. Date:

0 11110 36153 0

**PEEL HERE**

**Drug Facts** (continued)

**Ask a doctor or pharmacist before use if you are** ■ taking aspirin to prevent heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ under a doctor's care for any serious condition ■ taking any other drug

**When using this product** ■ take with food or milk if stomach upset occurs ■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

**Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear

**If pregnant or breast-feeding** ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- do not take more than directed ■ the smallest effective dose should be used ■ do not take longer than 10 days, unless directed by a doctor (see Warnings)
- Adults and children 12 years and older: ■ take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor ■ Children under 12 years: ask a doctor ■ do not use in children under 4 years of age

**Other information**

- Store between 20°-25°C (68°-77°F) ■ Read all warnings and directions before use. ■ do not use if inner seal under bottle cap imprinted with "Sealed for Your Protection" is broken or missing

**Inactive ingredients** anhydrous lactose, Canauba Wax, Colloidal Silicon Dioxide, Hypromellose, Iron Oxide Red, Magnesium Stearate, Microcrystalline Cellulose, Polydextrose, Polyethylene Glycol, Povidone, Sodium Lauryl Sulfate, Sodium Starch Glycolate, Starch (Corn), Titanium Dioxide

**Questions or comments?**  
1-800-632-6900.

Ibuprofen 200 mg tablets

<b>IBUPROFEN</b>			
ibuprofen tablet			
<b>Product Information</b>			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:30 142-167
<b>Route of Administration</b>	ORAL		
<b>Active Ingredient/Active Moiety</b>			
<b>Ingredient Name</b>		<b>Basis of Strength</b>	<b>Strength</b>

IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)		IBUPROFEN	200 mg	
<b>Inactive Ingredients</b>				
<b>Ingredient Name</b>			<b>Strength</b>	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)				
CARNAUBA WAX (UNII: R12CBM0EIZ)				
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
HYPROMELLOSES (UNII: 3NXW29V3WO)				
FERRIC OXIDE RED (UNII: 1K09F3G675)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)				
POLYDEXTROSE (UNII: VH2XOU12IE)				
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)				
POVIDONES (UNII: FZ989GH94E)				
SODIUM LAURYL SULFATE (UNII: 368GB5141J)				
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
<b>Product Characteristics</b>				
<b>Color</b>	BROWN	<b>Score</b>	no score	
<b>Shape</b>	CAPSULE	<b>Size</b>	14mm	
<b>Flavor</b>		<b>Imprint Code</b>	IP142	
<b>Contains</b>				
<b>Packaging</b>				
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:30142-167-05	500 in 1 BOTTLE		
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>	
ANDA	ANDA072199	02/02/2011		

**Labeler** - The Kroger Co. (006999528)

Revised: 9/2012

The Kroger Co.