

CONRX EXTRA STRENGTH- acetaminophen tablet
Eagle Distributors, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

ConRx™ Extra Strength

Important: Read all product information before using. Keep this box for important information.

Drug Facts

Active ingredient (in each caplet)

Acetaminophen 500 mg

Purpose

Pain reliever/fever reducer

Use

- temporarily relieves minor aches and pains due to:
 - the common cold
 - headache
 - backache
 - minor pain of arthritis
 - toothache
 - muscular aches
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. The maximum daily dose of this product is 6 caplets (3,000 mg) in 24 hours. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days

- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**
adults and children 12 years and over
 - take 2 caplets every 6 hours while symptoms last
 - do not take more than 6 caplets in 24 hours, unless directed by a doctor
 - do not use for more than 10 days unless directed by a doctor
- children under 12 years, ask a doctor

Other information

- store between 20-25°C (68-77°F)
- **do not use if pouch is torn or open**
- see side panel for lot number and expiration date

Inactive ingredients

carnauba wax*, castor oil*, corn starch, FD&C red #40 aluminum lake, hypromellose, magnesium stearate, polyethylene glycol*, powdered cellulose, pregelatinized starch, propylene glycol, shellac, sodium starch glycolate, titanium dioxide

- contains one or more of these ingredients

Questions or comments?

1-800-570-8650

PRINCIPAL DISPLAY PANEL - 50 Pouch Box

See New Warnings Information & Directions

**Compare to the Active Ingredients in
Tylenol® Extra Strength ***

ConRx™

**EXTRA
STRENGTH**

DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN
Acetaminophen
Pain Reliever - Fever Reducer

See New Warnings Information & Directions
**Compare to the Active Ingredients in
 Tylenol® Extra Strength**

ConRx™ EXTRA STRENGTH

DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN
Acetaminophen
 Pain Reliever - Fever Reducer

See New Warnings Information & Directions
**Compare to the Active Ingredients in
 Tylenol® Extra Strength**

ConRx™ EXTRA STRENGTH

DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN
Acetaminophen
 Pain Reliever - Fever Reducer

▲
 TO OPEN
 PUSH IN TAB AND PULL OUT

**Compare to the Active Ingredients in
 Tylenol® Extra Strength**

50 Pouches of 2 Caplets Each



50 Pouches of 2 Caplets Each

Acetaminophen
 Pain Reliever - Fever Reducer

DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN

ConRx™ EXTRA STRENGTH

**Compare to the Active Ingredients in
 Tylenol® Extra Strength**

See New Warnings Information & Directions

Important: Read all product information before using. Keep this box for important information.

Drug Facts	
Active ingredient (in each caplet)	Purpose
Acetaminophen 500 mg	Pain reliever/fever reducer
Uses	
<ul style="list-style-type: none"> ■ temporarily relieves minor aches and pains due to: ■ the common cold ■ headache ■ backache ■ minor pain of arthritis ■ toothache ■ muscular aches ■ premenstrual and menstrual cramps 	

See New Warnings Information & Directions
**Compare to the Active Ingredients in
 Tylenol® Extra Strength**

ConRx™ EXTRA STRENGTH


<ul style="list-style-type: none"> temporarily reduces fever 	
Warnings Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 6 caplets (3,000 mg) in 24 hours. Severe liver damage may occur if you take <ul style="list-style-type: none"> more than 4,000 mg of acetaminophen in 24 hours with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product 	
Do not use <ul style="list-style-type: none"> with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. if you are allergic to acetaminophen or any of the inactive ingredients in this product 	
Ask a doctor before use if you have liver disease	
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin	
Stop use and ask a doctor if <ul style="list-style-type: none"> pain gets worse or lasts more than 10 days fever gets worse or lasts more than 3 days new symptoms occur redness or swelling is present These could be signs of a serious condition.	
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.	
Directions do not take more than directed (see overdose warning)	
adults and children 12 years and over	<ul style="list-style-type: none"> take 2 caplets every 6 hours while symptoms last do not take more than 6 caplets in 24 hours, unless directed by a doctor do not use for more than 10 days unless directed by a doctor
children under 12 years	ask a doctor
Other information <ul style="list-style-type: none"> store between 20-25°C (68-77°F) do not use if card or pouch is torn or open see side panel for lot number and expiration date 	
Inactive ingredients carnauba wax*, castor oil*, corn starch, FD&C red #40 aluminum lake, hypromellose, magnesium stearate, polyethylene glycol*, powdered cellulose, pregelatinized starch, propylene glycol, shellac, sodium starch glycolate, titanium dioxide * contains one or more of these ingredients	
Questions or comments? 1-800-570-8650	

CONRX EXTRA STRENGTH

DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN

Acetaminophen
Pain Reliever - Fever Reducer

*This product is not manufactured or distributed by:
 McNeil Consumer Speciality Pharmaceuticals Division of McNeil PPC, Inc.
 Eagle Distributors, Inc./ConRx™ does not own the Tylenol® Extra Strength trademark.



Product manufactured for:
Eagle Distributors, Inc.
 Los Angeles, CA 90011

CONRX EXTRA STRENGTH

acetaminophen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68737-233
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Acetaminophen (UNII: 362O9ITL9D) (Acetaminophen - UNII:362O9ITL9D)	Acetaminophen	500 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
CASTOR OIL (UNII: D5340Y2I9G)	
STARCH, CORN (UNII: O8232NY3SJ)	
FD&C Red NO. 40 (UNII: WZB9127XOA)	
ALUMINUM OXIDE (UNII: LM26O6933)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYETHYLENE GLYCOL 6000 (UNII: 30IQX730WE)	
POWDERED CELLULOSE (UNII: SMD1X3XO9M)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

SHELLAC (UNII: 46N107B71O)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	RED	Score	2 pieces
Shape	OVAL	Size	18mm
Flavor		Imprint Code	CRX
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68737-233-17	50 in 1 BOX		
1		2 in 1 POUCH		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	02/15/2013	

Labeler - Eagle Distributors, Inc. (929837425)

Revised: 2/2013

Eagle Distributors, Inc.