

**GOOD SENSE NIGHTTIME COUGH- dextromethorphan hydrobromide,  
doxylamine succinate solution**

**L. Perrigo Company**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Perrigo Night Time Cough Drug Facts**

**Active ingredients (in each 30 mL)**

Dextromethorphan HBr 30 mg

Doxylamine succinate 12.5 mg

**Purpose**

Cough suppressant

Antihistamine

**Uses**

temporarily relieves cold symptoms:

- cough due to minor throat and bronchial irritation
- runny nose and sneezing

**Warnings**

**Do not use**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough as occurs with smoking, asthma, or emphysema
- trouble urinating due to an enlarged prostate gland

**Ask a doctor or pharmacist before use if you are**

taking sedatives or tranquilizers

### **When using this product**

- excitability may occur, especially in children
- may cause marked drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

### **Stop use and ask a doctor if**

cough persists for more than 1 week, tends to recur, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### **Directions**

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

### **Other information**

- **each 30 mL contains:** sodium 32 mg
- store at 20-25°C (68-77°F)

### **Inactive ingredients**

alcohol, anhydrous citric acid, FD&C blue no. 1, FD&C red no. 40, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

### **Questions or comments?**

**1-800-719-9260**

## **Principal Display Panel**

GoodSense®

All Night Cough Relief

Cough Suppressant, Antihistamine

Night Time Cough

Dextromethorphan HBr

Doxylamine Succinate

- Cough
- Sneezing
- Runny Nose

Cherry Flavor

Compare to active ingredients of Vicks® NyQuil® Cough

12 FL OZ (354 mL)

ALCOHOL 10%

NDC 0113-0668-40

GOODSENSE®

All Night Cough Relief

Cough Suppressant, Antihistamine

# Night Time Cough

Dextromethorphan HBr  
Doxylamine Succinate

- Cough
- Sneezing
- Runny Nose

**Cherry Flavor**

Compare to active ingredients of  
Vicks® NyQuil® Cough

12 FL OZ (354 mL) ALCOHOL 10%

: 66840 C2 F2

Distributed By

**Perrigo®**

Allegan, MI 49010

Empty & Recycle Cap



PLASTIC BOTTLE



PLASTIC CUP

how2recycle.info

Gluten Free

DO NOT USE IF PRINTED NECKBAND IS  
BROKEN OR MISSING

**PARENTS:**

Learn about teen medicine abuse

[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

N  
0113-0668-40  
6



### Drug Facts

Active ingredients (in each 30 mL)	Purpose
Dextromethorphan HBr 30 mg.....	Cough suppressant
Doxylamine succinate 12.5 mg.....	Antihistamine

**Uses** temporarily relieves cold symptoms:

- cough due to minor throat and bronchial irritation
- runny nose and sneezing

### Warnings

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

PEEL BACK AT  
CORNER FOR MORE  
INFORMATION

: 66840 C2 B1

### Drug Facts (continued)

Ask a doctor before use if you have

- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough as occurs with smoking, asthma, or emphysema
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

**When using this product**

- excitability may occur, especially in children
- may cause marked drowsiness
- avoid alcoholic drinks
- be careful when driving a motor

ADHESIVE AREA  
NO VARNISH • NO TYPE

### Drug Facts (continued)

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use


### Other information

- each 30 mL contains: sodium 32 mg
- store at 20-25°C (68-77°F)

### Inactive ingredients

alcohol, anhydrous citric acid, FD&C blue no. 1, FD&C red no. 40, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol,

ADHESIVE AREA  
NO VARNISH • NO TYPE



## GOOD SENSE NIGHTTIME COUGH

dextromethorphan hydrobromide, doxylamine succinate solution

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0113-0668
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>ALCOHOL</b> (UNII: 3K9958V90M)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>HIGH FRUCTOSE CORN SYRUP</b> (UNII: XY6UN3QB6S)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ05DW1A)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SODIUM CITRATE, UNSPECIFIED FORM</b> (UNII: 1Q73Q2JULR)	

### Product Characteristics

<b>Color</b>	RED (Dark Red)	<b>Score</b>	
--------------	----------------	--------------	--

<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0113-0668-30	177 mL in 1 BOTTLE; Type 0: Not a Combination Product	11/03/2002	11/14/2013
2	NDC:0113-0668-34	237 mL in 1 BOTTLE; Type 0: Not a Combination Product	09/12/2012	
3	NDC:0113-0668-40	354 mL in 1 BOTTLE; Type 0: Not a Combination Product	02/01/2022	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	11/03/2002	

**Labeler** - L. Perrigo Company (006013346)

Revised: 2/2022

L. Perrigo Company