

MEDI FIRST PAIN RELIEF- acetaminophen, aspirin, salicylamide and caffeine tablet

MEDI FIRST PLUS PAIN ZAPPER- acetaminophen, aspirin, salicylamide and caffeine tablet

Unifirst First Aid Corporation

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Medi-First[®] Pain Relief

Drug Facts

Active ingredient (in each tablet)

Acetaminophen 110 mg

Aspirin (NSAID)* 162 mg

Caffeine 32.4 mg

Salicylamide (NSAID)* 152 mg

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Pain reliever/fever reducer

Adjuvant

Pain reliever/fever reducer

Uses

temporarily relieves minor aches and pains associated with

- headache
- muscular aches
- minor arthritis pain
- back pain
- colds
- toothache
- menstrual cramps

Temporarily reduces fever.

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin and Salicylamide may cause a severe allergic reaction which may

include:

■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

■ skin reddening ■ blisters ■ rash

If a skin or allergic reaction occurs, stop use and seek medical help right away.

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains nonsteroidal anti-inflammatory drugs (NSAIDs), which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine- containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for pain for more than 10 days unless directed by a doctor
- for more than 3 days for fever unless directed by a doctor
- with any other pain releiver/fever reducer

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems (such as heartburn, upset stomach or stomach pain)
- you have heart disease, liver cirrhosis, or kidney disease
- you have asthma or high blood pressure
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- anticoagulation

- diabetes
- gout
- arthritis

Ask a doctor or pharmacist before use if you have

- liver disease

When using this product do not exceed recommended dose

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding: • feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better
- symptoms do not improve
- pain or fever gets worse
- ringing in the ears or hearing loss occurs
- new symptoms appear

If pregnant or breast-feeding, ask a health care professional before use. **It is especially important not to use aspirin or salicyamide during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

Keep out of reach of children.

Overdose warning:

Taking more than the recommended dose (overdose) may cause liver damage.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not use more than directed**

Adults and children(12 years and older): Take 2 tablets every 4 hours as needed with a full glass of water. Do not exceed 8 tablets in 24 hours or as directed by a doctor

Children under 12 years of age: Ask a doctor

Other information

- store at room temperature 59°-86°F (15°-30°C)
- avoid excessive heat and humidity
- tamper evident sealed packets
- do not use any opened or torn packets

Inactive ingredients

FD&C yellow #6, magnesium stearate, microcrystalline cellulose, povidone, starch, stearic acid.

Question or comments? 1-800-634-7680

Medi-First Pain Relief Label

100 Tablets

(50 x 2)

Medi-First®

Pain Relief extra strength

Fast Acting Advanced Formula

Pull to Open

Aches, & Fever • Acetaminophen 110 mg Aspirin (NSAID) 162 mg

Pain Reliever • Salicylamide (NSAID) 152 mg

Pain Reliever Aid • Caffeine 32.4 mg

This Package is for Households without Young Children.

Tamper Evident Unit Dose Packets

Drug Facts

Active ingredients (in each tablet) Purpose

- Acetaminophen 110 mg Pain reliever/fever reducer
- Aspirin (NSAID)* 162 mg Pain reliever/fever reducer
- Caffeine 32.4 mg Adjunct
- Salicylamide (NSAID)† 152 mg Pain reliever/fever reducer

Uses

temporarily relieves minor aches and pains associated with:

- headaches
- muscular aches
- minor arthritis pain
- back pain
- colds
- toothache
- menstrual cramps

Temporarily reduces fever.

Warnings

Roy's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Roy's syndrome, a rare but serious illness.

Allergy alert: Aspirin and Salicylamide may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin redness
- blisters
- itch

Drug Facts (continued)

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for pain for more than 10 days unless directed by a doctor.
- for more than 3 days for fever unless directed by a doctor.
- with any other pain reliever/fever reducer.

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems (such as heartburn, upset stomach or stomach pain)
- you have heart disease, liver cirrhosis, or kidney disease
- you have asthma or high blood pressure
- you are taking a diuretic.

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- anticoagulation
- diabetes
- gout
- arthritis

Ask a doctor or pharmacist before use if you have

- liver disease

When using this product do not exceed recommended dose

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better

Drug Facts (continued)

if symptoms do not improve

- pain or fever gets worse
- ringing in the ears or loss of hearing occurs
- new symptoms occur

If pregnant or breast-feeding, ask a health care professional before use. It is especially important not to use aspirin or salicylamide during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed

Adults and children (12 years and older)

Take 2 tablets every 4 hours as needed with a full glass of water. Do not exceed 8 tablets in 24 hours, or as directed by a doctor.

Children under 12 years:

Ask a doctor

Other information

- store at room temperature (59-86°F) (15-30°C)
- avoid excessive heat and humidity
- tamper evident sealed packets
- do not use any opened or torn packets

Inactive ingredients

FD&C yellow #6, magnesium stearate, microcrystalline cellulose, polydioxane, starch, stearic acid

Questions or Comments? 1-800-634-7680

Medi-First

Reorder #61133
Manufactured for
Medique Products
Fort Myers, FL 33967
1-800-634-7680
www.medi-oneproducts.com

3 47682 81133 4

Retain carton for complete product information

Medi-First Plus Pain Zapper Label

250 Tablets

(125 x 2's)

Medi-First® Plus

Pain Zapper

This Package is for Households without Young Children.

Pull To Open

Acetaminophen 110 mg,

Aspirin (NSAID) 162 mg,

Caffeine 32.4 mg, Salicylamide (NSAID) 152 mg,

Extra Strength Pain Reliever/Fever Reducer

Tamper Evident Unit Dose Packets

Drug Facts

Active Ingredients (in each tablet)

Active Ingredient	Purpose
Acetaminophen 110 mg	Pain reliever/fever reducer
Aspirin (NSAID) ¹ 162 mg	Pain reliever/fever reducer
Caffeine 32.4 mg	Adjuvant
Salicylamide (NSAID) ² 152 mg	Pain reliever/fever reducer
Nonsteroidal anti-inflammatory drug	

Uses
Temporarily relieves minor aches and pains associated with:
 ■ headaches ■ muscular aches ■ minor arthritis pain ■ back pain ■ colds ■ toothache ■ menstrual cramps
 Temporarily reduces fever.

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin and Salicylamide may cause a severe allergic reaction which may include:
 ■ hives ■ facial swelling ■ asthma (breathing) ■ shock

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 ■ skin redness ■ blisters ■ rash

If a skin or allergic reaction occurs, stop use and seek medical help right away.

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
 ■ more than 4,000 mg of acetaminophen in 24 hours, which is the maximum daily amount
 ■ with other drugs containing acetaminophen
 ■ 3 or more alcoholic drinks every day while using this product
 ■ are age 60 or older
 ■ have had stomach issues or bleeding problems
 ■ take a blood thinning (anticoagulant) or steroid drug
 ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 ■ have 3 or more alcoholic drinks every day while using this product
 ■ take more or for a longer time than directed. (Continued on opposite panel)



250 Tablets (125 x 2's)

THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.

Acetaminophen 110 mg, Aspirin (NSAID) 162 mg, Caffeine 32.4 mg, Salicylamide (NSAID) 152 mg

Extra Strength Pain Reliever/
Fever Reducer

TAMPER EVIDENT UNIT DOSE PACKETS

Drug Facts (continued)

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and occasionally, rapid heart beat.

Do not use
 ■ with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ for pain for more than 10 days unless directed by a doctor
 ■ for more than 3 days for fever unless directed by a doctor
 ■ with any other pain reliever/fever reducer

Ask a doctor before use if
 ■ stomach bleeding warning applies to you
 ■ you have a history of stomach problems (such as heartburn, upset stomach or stomach pain)
 ■ you have heart disease, heart trouble, or kidney disease
 ■ you have asthma or high blood pressure
 ■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are taking a prescription drug for
 ■ anticoagulation ■ diabetes ■ gout ■ arthritis

Ask a doctor or pharmacist before use if you have ■ liver disease

When using this product do not exceed recommended dose

Stop use and ask a doctor if
 ■ you experience any of the following signs of stomach bleeding:
 ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better
 ■ symptoms do not improve
 ■ pain or fever gets worse
 ■ ringing in the ears or loss of hearing occurs
 ■ new symptoms occur

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. Get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions ■ do not use more than directed

Adults and children (12 years and older) ■ Take 2 tablets every 4 hours as needed with a full glass of water. Do not exceed 8 tablets in 24 hours, or as directed by a doctor.

Children under 12 years: Ask a doctor

Other information
 ■ store at room temperature 68°-86°F (20°-32°C)
 ■ avoid excessive heat and humidity
 ■ tamper evident sealed packets
 ■ do not use any opened or torn packets

Inactive ingredients FD&C yellow #6, magnesium stearate, microcrystalline cellulose, povidone, starch, stearic acid

Questions or Comments? 1-800-634-7680

Retain carton for complete product information

Retain carton for complete product information

MEDI FIRST PAIN RELIEF

acetaminophen, aspirin, salicylamide and caffeine tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:47682-811
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	110 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	162 mg
SALICYLAMIDE (UNII: EM8BM710ZC) (SALICYLAMIDE - UNII:EM8BM710ZC)	SALICYLAMIDE	152 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	32.4 mg

Inactive Ingredients

Ingredient Name	Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
MAGNESIUM STEARATE (UNII: 70097M6130)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
STARCH, CORN (UNII: O8232NY3SJ)	

Product Characteristics

Color	orange (orange)	Score	no score
Shape	ROUND (ROUND)	Size	11mm
Flavor		Imprint Code	FR;2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-811-33	50 in 1 BOX	12/30/2008	
1		2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:47682-811-48	125 in 1 BOX	12/30/2008	
2		2 in 1 PACKET; Type 0: Not a Combination Product		
3	NDC:47682-811-13	250 in 1 BOX	12/30/2008	
3	NDC:47682-811-99	2 in 1 PACKET; Type 0: Not a Combination Product		
4	NDC:47682-811-99	2 in 1 PACKET; Type 0: Not a Combination Product	12/30/2008	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	12/30/2008	

MEDI FIRST PLUS PAIN ZAPPER

acetaminophen, aspirin, salicylamide and caffeine tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:47682-911
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	110 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	162 mg
SALICYLAMIDE (UNII: EM8BM710ZC) (SALICYLAMIDE - UNII:EM8BM710ZC)	SALICYLAMIDE	152 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	32.4 mg

Inactive Ingredients

Ingredient Name	Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
STARCH, CORN (UNII: O8232NY3SJ)	

Product Characteristics

Color	orange (orange)	Score	no score
Shape	ROUND (ROUND)	Size	11mm
Flavor		Imprint Code	FR;2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-911-33	50 in 1 BOX	12/30/2008	
1		2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:47682-911-48	125 in 1 BOX	12/30/2008	
2		2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	12/30/2008	

Labeler - Unifirst First Aid Corporation (832947092)

Revised: 3/2022

Unifirst First Aid Corporation