

**VANQUISH- acetaminophen, aspirin, and caffeine tablet, coated**  
**DSE Healthcare Solutions, LLC**

-----  
**Vanquish**

**Drug Facts**

**Active ingredients**

<b>Active ingredients (in each caplet)</b>	<b>Purpose</b>
Acetaminophen 194 mg	Pain reliever
Aspirin 227 mg (NSAID)*	Pain reliever
Caffeine 33 mg	Pain reliever aid

\* nonsteroidal anti-inflammatory drug

**Uses** temporarily relieves minor pain due to

- headache
- arthritis
- backache
- colds
- menstrual cramps
- muscle aches

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters

- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 caplets in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to aspirin or any other pain reliever/fever reducer

### **Ask a doctor before use if**

- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

**Ask a doctor or pharmacist before use if you are** taking a prescription drug for

- gout
- diabetes
- arthritis

### **Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- redness or swelling is present
- ringing in the ears or loss of hearing occurs
- new symptoms occur

**If pregnant or breast-feeding**, ask a health professional before use.

**It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- do not take more than the recommended dose
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 caplets every 6 hours, not to exceed 8 caplets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

### **Other information**

- save carton for full directions and warnings
- store at room temperature

### **Inactive ingredients**

Colloidal Silicon Dioxide, Corn Starch, Hypromellose, Maltodextrin, Microcrystalline Cellulose, Polydextrose, Povidone, Sodium Carboxymethyl Starch, Stearic Acid, Talc, Titanium Dioxide and Triglycerides

Each tablet contains 0.11 mg. of Sodium

### **Questions or comments?**

Call: 1-800- 214-2379

[www.vanquishheadacherelief.com](http://www.vanquishheadacherelief.com)

VANQUISH® is a registered trademark of DSE Healthcare Solutions

Distributed by DSE Healthcare Solutions, LLC  
27070 Miles Rd, Suite A, Solon, OH 44139

## **Principal Display Panel**

**NEW LOOK!**  
**Same Trusted Formula**

**MADE IN USA**  
*with Domestic and Imported Ingredients*

**Effective Headache Relief**

**VANQUISH®**  
*Aspirin (NSAID) / Acetaminophen / Caffeine Pain Reliever / Pain Reliever Aid*

**100** Analgesic COATED CAPLETS

R51019  
011-2201 C  
02/22

### **Top Panel**

Do not use this product if printed safety seal under cap is torn or missing.

**Effective Headache Relief**

**VANQUISH® 50 Years of Proven Pain Relief**  
**Refreshed for Today's Digital Lifestyle**

**EXPIRES**

### **Bottom Panel**

Questions or comments?

Call: 1-800-214-2379

VANQUISH® is a registered trademark of DSE Healthcare Solutions  
Distributed by DSE Healthcare Solutions, LLC  
27070 Miles Rd, Suite A, Solon OH 44139

OMIT A

EXPIRES

**VANQUISH**  
Effective Headache Relief

Do not use this product if printed safety seal under cap is torn or missing.

R51019  
011-2201 C  
02/22

### Drug Facts

#### Active Ingredients (in each caplet)

Active Ingredients (in each caplet)	Purpose
Acetaminophen 154 mg	Pain reliever
Aspirin 227 mg (NSAID)*	Pain reliever
Caffeine 33 mg	Pain reliever aid

**Uses** temporarily relieves minor pain due to

- headache
- backache
- menstrual cramps
- arthritis
- colic
- muscle aches

#### Warnings

**Peyer's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Peyer's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Allergy alert:** Acetaminophen may cause severe skin reactions.

Symptoms may include: • skin redness/itching • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

**Liver warning:** This product contains acetaminophen.

Severe liver damage may occur if you take

- more than 8 caplets in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 2 or more alcoholic drinks every day while using this product

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinner (anticoagulant) or steroid drug
- take other drugs containing prescription or over-the-counter NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 2 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

Do not use

### Drug Facts (continued)

- with any other drug containing acetaminophen (prescription or over-the-counter). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to aspirin or any other pain reliever/fever reducer

**Ask a doctor before use if**

- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, low circulation, or kidney disease
- you are taking a diuretic
- you have asthma

**Ask a doctor or pharmacist before use if you are taking a prescription drug for**

- gout
- diabetes
- arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- redness or swelling is present
- ringing in the ears or loss of hearing occurs
- new symptoms occur

**If pregnant or breast-feeding, ask a health professional before use.**

It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### Directions

- do not take more than the recommended dose
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 caplets every 6 hours, not to exceed 8 caplets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

**Other information**

- save carton for full directions and warnings
- store at room temperature

#### Inactive Ingredients

Colloidal Silicon Dioxide, Corn Starch, Hydroxyethylcellulose, Maltodextrin, Microcrystalline Cellulose, Polydextrose, Povidone, Sodium Carboxymethyl Starch, Stearic Acid, Talc, Titanium Dioxide and Trihydroxy

NEW Same Great Formula

NEW LOOK! Same Trusted Formula

MADE IN USA with Domestic and Imported Ingredients

# VANQUISH

### Effective Headache Relief

Aspirin (NSAID) /Acetaminophen /Caffeine  
Pain Reliever/  
Pain Reliever Aid

100 Analgesic COATED CAPLETS

100 Analgesic COATED CAPLETS

Effective Headache Relief  
**VANQUISH**  
Aspirin (NSAID) /Acetaminophen /Caffeine  
Pain Reliever/Pain Reliever Aid

MADE IN USA with Domestic and Imported Ingredients

Questions or comments?  
Call: 1-800-214-2379



8 8941112201 9

VANQUISH® is a registered trademark of OSE Healthcare Solutions  
Distributed by OSE Healthcare Solutions, LLC  
27070 Miles Rd., Suite A, Solon, OH 44139

---LEB6736P---



**See Carton for Full Information** **Active ingredients (in each caplet)** Acetaminophen 194 mg, Aspirin 227 mg, Caffeine 33 mg **Warnings** **Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. **Allergy alert:** Aspirin may cause a severe allergic reaction which may include: hives, facial swelling, asthma (wheezing), and shock. **Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening, blisters and rash. If a skin reaction occurs, stop use and seek medical help right away. **Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take more than 8 caplets in 24 hours, which is the maximum daily amount for this product with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product. **Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older have had stomach ulcers or bleeding problems take a blood thinning (anticoagulant) or steroid drug take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) have 3 or more alcoholic drinks every day while using this product take more or for a longer time than directed. **Do not use** with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. **Stop use and ask a doctor if** an allergic reaction occurs (seek medical help right away) you experience any of the following signs of stomach bleeding: (feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better) pain gets worse or lasts more than 10 days redness or swelling is present ringing in the ears or loss of hearing occurs new symptoms occur. **If pregnant or breast-feeding,** ask a health professional before use. **Directions:** Do not take more than the recommended dose. Drink a full glass of water with each dose. Adults and children 12 years and over: take 2 caplets every 6 hours, not to exceed 8 caplets in 24 hours unless directed by a doctor. Children under 12 years: consult a doctor. **Questions or Comments?** 1-800-214-2378 or [www.vanquishheadacherelief.com](http://www.vanquishheadacherelief.com)

011-2201 C  
11043-03-17  
Distributed by DSE Healthcare Solutions, LLC  
27070 Miles Rd, Suite A, Solon OH 44139  
EXPIRES 08-375210

PRINTS ONLINE  
MM/YYYY  
XXXXXX

**MADE IN USA**

**Effective Headache Relief**

**VANQUISH®**

**Aspirin (NSAID) / Acetaminophen / Caffeine Pain Reliever / Pain Reliever Aid**

**100 Analgesic COATED CAPLETS**

VANQUISH® is a registered trademark of SVADS Holdings SA  
Distributed by DSE Healthcare Solutions, LLC  
27070 Miles Rd, Suite A, Solon OH 44139

[www.vanquishheadacherelief.com](http://www.vanquishheadacherelief.com)

EXPIRES

11043-03-17

011-2201 C

**MADE IN USA**

Do not use this product if printed safety seal under cap is torn or missing.

**Effective Headache Relief**

**VANQUISH®**

**Aspirin (NSAID) / Acetaminophen / Caffeine Pain Reliever / Pain Reliever Aid**

**120 Analgesic COATED CAPLETS**

Distributed by DSE Healthcare Solutions, LLC

27070 Miles Rd, Suite A, Solon OH 44139

011-2202 L

EXPIRES 82-375210 11044-03-17



**See Carton for Full Information** **Active ingredients (in each caplet)** Acetaminophen 194 mg, Aspirin 227 mg, Caffeine 33 mg **Warnings** **Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. **Allergy alert:** Aspirin may cause a severe allergic reaction which may include: hives, facial swelling, asthma (wheezing), and shock. **Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening, blisters and rash. If a skin reaction occurs, stop use and seek medical help right away. **Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take more than 8 caplets in 24 hours, which is the maximum daily amount for this product with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product. **Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older have had stomach ulcers or bleeding problems take a blood thinning (anticoagulant) or steroid drug take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) have 3 or more alcoholic drinks every day while using this product take more or for a longer time than directed. **Do not use** with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. if you are allergic to aspirin or any other pain reliever/fever reducer. **Stop use and ask a doctor if** an allergic reaction occurs (seek medical help right away) you experience any of the following signs of stomach bleeding: (feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better) pain gets worse or lasts more than 10 days redness or swelling is present ringing in the ears or loss of hearing occurs new symptoms occur. **If pregnant or breast-feeding,** ask a health professional before use. **Directions** Do not take more than the recommended dose. Drink a full glass of water with each dose. Adults and children 12 years and over: take 2 caplets every 6 hours, not to exceed 8 caplets in 24 hours unless directed by a doctor. Children under 12 years: consult a doctor.

**Questions or Comments?** 1-800-214-2379 or [www.vanquishheadacherelief.com](http://www.vanquishheadacherelief.com)

011-2202 L  
11044-03-17  
Distributed by: DSE Healthcare Solutions, LLC  
27070 Miles Rd, Suite A, Solon OH 44139  
82-375210  
EXPIRES

PRINTS ONLINE

MM/YYYY  
XXXXXX

**MADE IN USA**

**- BONUS - 20% MORE!**

**Effective Headache Relief**

**VANQUISH®**

**Aspirin (NSAID) / Acetaminophen / Caffeine Pain Reliever / Pain Reliever Aid**

**120 Analgesic COATED CAPLETS**

VANQUISH® is a registered trademark of SVADS Holdings SA  
Distributed by DSE Healthcare Solutions, LLC  
27070 Miles Rd, Suite A, Solon OH 44139

[www.vanquishheadacherelief.com](http://www.vanquishheadacherelief.com)

EXPIRES

11045-03-17

011-2202 C



# VANQUISH

acetaminophen, aspirin, and caffeine tablet, coated

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:42364-0011
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
-----------------	-------------------	----------

<b>Acetaminophen</b> (UNII: 362O9ITL9D) (Acetaminophen - UNII:362O9ITL9D)	Acetaminophen	194 mg
<b>Aspirin</b> (UNII: R16CO5Y76E) (Aspirin - UNII:R16CO5Y76E)	Aspirin	227 mg
<b>Caffeine</b> (UNII: 3G6A5W338E) (Caffeine - UNII:3G6A5W338E)	Caffeine	33 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>Microcrystalline Cellulose</b> (UNII: OP1R32D61U)	
<b>Hypromellose, Unspecified</b> (UNII: 3NXW29V3WO)	
<b>Maltodextrin</b> (UNII: 7CVR7L4A2D)	
<b>Medium-Chain Triglycerides</b> (UNII: C9H2L21V7U)	
<b>Polydextrose</b> (UNII: VH2XOU12IE)	
<b>Povidone K30</b> (UNII: U725QWY32X)	
<b>Silicon Dioxide</b> (UNII: ETJ7Z6XBU4)	
<b>Sodium Starch Glycolate Type A Corn</b> (UNII: AG9B65PV6B)	
<b>Starch, Corn</b> (UNII: O8232NY3SJ)	
<b>Stearic Acid</b> (UNII: 4ELV7Z65AP)	
<b>Talc</b> (UNII: 7SEV7J4R1U)	
<b>Titanium Dioxide</b> (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	WHITE	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	17mm
<b>Flavor</b>		<b>Imprint Code</b>	VANQUISH
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:42364-0011-1	1 in 1 CARTON	07/15/2018	
1		100 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:42364-0011-2	1 in 1 CARTON	07/15/2018	
2		120 in 1 BOTTLE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	07/15/2018	

**Labeler** - DSE Healthcare Solutions, LLC (603016069)

