

VICKS NYQUIL KIDS BERRY COLD AND COUGH PLUS RUNNY NOSE-
chlorpheniramine maleate, dextromethorphan hbr liquid
The Procter & Gamble Manufacturing Company

VICKS® NyQuil™ Kids BERRY
COLD & COUGH + RUNNY NOSE

Drug Facts

Active ingredients (in each 15 mL)

Chlorpheniramine maleate 2 mg
Dextromethorphan HBr 15 mg

Purpose

Antihistamine
Cough suppressant

Uses

temporarily relieves cold symptoms:

- cough due to minor throat & bronchial irritation
- sneezing
- runny nose

Warnings

Do not use

- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- to make a child sleepy

Ask a doctor before use if you have

- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are

taking sedatives or tranquilizers.

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, & tranquilizers may increase drowsiness

Stop use and ask a doctor if

- cough lasts more than 1 week, tends to recur, or is accompanied by fever, rash, or persistent headache.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 6 hrs
children 6 to under 12 yrs	15 mL every 6 hrs
children 4 to under 6 yrs	do not use unless directed by a doctor
children under 4 yrs	do not use

Other information

- **each 15 mL contains:** sodium 13 mg
- Store at no greater than 25°C

Inactive ingredients

citric acid, FD&C Red No. 40, flavor, glycerin, propylene glycol, saccharin sodium,

sodium benzoate, sodium citrate, sorbitol, sucralose, water, xanthan gum

Questions?

1-800-362-1683

**TAMPER EVIDENT: DO NOT USE IF PRINTED SHRINKBAND ON BOTTLE IS
BROKEN OR MISSING.**

**DIST. BY PROCTER & GAMBLE,
CINCINNATI OH 45202**

PRINCIPAL DISPLAY PANEL - 236 ml Bottle Label

VICKS[®] NyQuil[™]

Kids

BERRY

COLD & COUGH + RUNNY NOSE

Chlorpheniramine maleate,

Dextromethorphan HBr

Sneezing, Runny Nose

Cough

AGES 6+

Nighttime Relief

FREE OF ALCOHOL

& ACETAMINOPHEN

8 FL OZ (236 mL)

Drug Facts (continued)

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 6 hrs
children 6 to under 12 yrs	15 mL every 6 hrs
children 4 to under 6 yrs	do not use unless directed by a doctor
children under 4 yrs	do not use

Other information

- each 15 mL contains: sodium 13 mg
- store at no greater than 25°C

Inactive ingredients citric acid, FD&C Red No. 40, flavor, glycerin, propylene glycol, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, water, xanthan gum

Questions? 1-800-362-1683

Drug Facts (continued)

- pharmacist before taking this product.
- to make a child sleepy

Ask a doctor before use if you have

- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- cough persists for more than 1 week, tends to recur, or is accompanied by a fever, rash, or persistent headache.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

TAMPER EVIDENT. DO NOT USE IF PRINTED SHRINKBAND SEAL AROUND THE NECK IS BROKEN OR MISSING. PEEL BACK FOR DRUG FACTS

Drug Facts

Active ingredients (in each 15 mL) Purpose
 Chlorpheniramine maleate 2 mg.....Antihistamine
 Dextromethorphan HBr 15 mg.....Cough suppressant

Uses temporarily relieves cold symptoms:
 • cough due to minor throat & bronchial irritation
 • sneezing • runny nose

Warnings
 Do not use
 • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist

PARENTS:
 Learn about tamper-evident seals at www.StopMedicineAbuse.org

DIST BY: PROCTER & GAMBLE, CINCINNATI, OH 45202
 Patents: www.pg.com/patents
 vicks.com

P&G
www.pg.com
 90451995

PLASTIC BOTTLE

23900 01433 6

VICKS

NyQuil Kids

BERRY

COLD & COUGH + RUNNY NOSE

Chlorpheniramine maleate, Dextromethorphan HBr

Sneezing, Runny Nose, Cough

AGES 6+

8 FL OZ (236 mL)

90472357
 Nighttime Relief
 FREE OF ALCOHOL & ACETAMINOPHEN

VICKS NYQUIL KIDS BERRY COLD AND COUGH PLUS RUNNY NOSE

chlorpheniramine maleate, dextromethorphan hbr liquid

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69423-779
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg in 15 mL	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg in 15 mL	

Inactive Ingredients	
Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
GLYCERIN (UNII: PDC6A3C0OX)	
SORBITOL (UNII: 506T60A25R)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	red	Score	
Shape		Size	
Flavor	BERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69423-779-08	236 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/06/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/06/2023	

Labeler - The Procter & Gamble Manufacturing Company (004238200)

Revised: 10/2023

The Procter & Gamble Manufacturing Company