SENNA-S- sennosides and docusate sodium tablet STRATEGIC SOURCING SERVICES LLC

ft 455b

Active ingredient (in each tablet)

Docusate Sodium 50 mg Sennosides 8.6 mg

Purpose

Stool softener

Laxative

Uses

- relieves occasional constipation (irregularity)
- this product generally produces a bowel movement in 6 to 12 hours

Warnings

Do not use

- if you are now taking mineral oil, unless directed by a doctor
- laxative products for longer than 1 week unless directed by a doctor

Ask a doctor before use if you have

- stomach pain, nausea or vomiting
- noticed a sudden change in bowel habits that lasts over two weeks

Stop use and ask a doctor if you have rectal bleeding or fail to have a bowel movement after use of a laxative. These may indicate a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (800-222-1222).

Directions

- If taking the starting dosage, take preferably at bedtime; or as directed by a doctor.
- If taking the maximum dosage, take in the morning and at bedtime.

Age	Starting Dose	Maximum Dose
adults and children 12 years of age and over	2 tablets once a day	4 tablets twice a day

children 6 to under 12 years	1 tablet once a day	2 tablets twice a day
children 2 to under 6 years	1/2 tablet once a day	2 tablets twice a day
children under 2 years	ask a doctor	ask a doctor

Other information

- each tablet contains: calcium 20 mg, sodium 3 mg
- store at 20° to 25°C (68° to 77°F); excursions permitted between 15° to 30°C (59° to 86°F).

Inactive ingredients

cellulose, croscarmellose sodium, dicalcium phosphate, FD&C yellow #6 lake, hypromellose, magnesium stearate, PEG, silica, talc, titanium dioxide.

Questions or Comments?

Call 833-358-6431 Monday to Friday 9:00am to 7:00pm EST

Package Label

		_
	NDC 70677-1069-1 COMPARE TO ACTIVE INGREDIENTS IN SENOKOT-S* DUAL ACTION*	
	Foster & Thrive™	
-	Senna-S	
	Sennosides 8.6 mg Docusate Sodium 50 mg NATURAL VEGETABLE LAXATIVE PLUS STOOL SOFTENER • Gentle, Dependable, Overnight Relief	
Foster € Thrive*	NDC 70677-1069-1 COMPARE TO ACTIVE INGREDIENTS IN SENOKOT-S* DUAL ACTION*	
Senna-S	Foster & Thrive™	
	Senna-S	
TAMPER EVIDENT: DO NOT USE IF IMPRINTED SEAL UNDER CAP IS MISSING OR BROKEN.	Sennosides 8.6 mg Docusate Sodium 50 mg NATURAL VEGETABLE LAXATIVE	
60 TABLETS	PLUS STOOL SOFTENER ACTUAL SIZE • Gentle, Dependable, Overnight Relief 60 TABLETS	
	Drug Facts SAVE CARTON FOR COMPLETE DRUG FACTS Active ingredients (in each tablet) Purpose Docusate Sodium 50 mg. Stool Softener Sennosides 8.6 mg. Stool Softener	
	Active ingredients (in each tablet) Purpose Docusate Sodium 50 mg	
	Drag Pacts Purpose Active ingredients (in each tablet) Purpose Docusate Sodium 50 mg. Stool Softener Senosides 8.6 mg. Stool Softener Laxative Laxative Uses • relieves occasional constipation (irregularity) • this product generally produces a bowel movement in 6 to 12 hours Warnings	3 4558
	Drug Facts Purpose Active ingredients (in each tablet) Purpose Docusate Sodium 50 mg. Stool Softener Sennosides 8.6 mg. Stool Softener Laxative Laxative Uses • relieves occasional constipation (irregularity) • this product generally produces a bowel movement in 6 to 12 hours Warnings Do not use • if you are now taking mineral oil, unless directed by a doctor • laxative products for longer than 1 week unless directed by a doctor Ask a doctor before use if you have • stomach pain, nausea or vomiting	M 02/23 456B
	Drug racis Purpose Active ingredients (in each tablet) Purpose Docusate Sodium 50 mg. Stool Softener Sennosides 8.6 mg. Stool Softener Uses • relieves occasional constipation (irregularity) • this product generally produces a bowel movement in 6 to 12 hours Warnings Do not use • if you are now taking mineral oil, unless directed by a doctor • laxative products for longer than 1 week unless directed by a doctor	REV 02/23 456B
	Drag Factor Purpose Active ingredients (in each tablet) Purpose Docusate Sodium 50 mg. Stool Softener Sennosides 8.6 mg. Stool Softener Laxative Laxative Uses • relieves occasional constipation (irregularity) • this product generally produces a bowel movement in 6 to 12 hours Warnings Do not use • if you are now taking mineral oil, unless directed by a doctor • laxative products for longer than 1 week unless directed by a doctor • laxative products for longer than 1 week unless directed by a doctor • ska a doctor before use if you have • stomach pain, nausea or vomiting • noticed a sudden change in bowel habits that lasts over two weeks Stop use and ask a doctor if you have rectal bleeding or fail to have a bowel movement after use	REV 02/23 45.68
	Drug Fuers Purpose Active ingredients (in each tablet) Purpose Docusate Sodium 50 mg. Stool Softener Sennosides 8.6 mg. Stool Softener Uses • relieves occasional constipation (irregularity) • this product generally produces a bowel movement in 6 to 12 hours Warnings Do not use • if you are now taking mineral oil, unless directed by a doctor • laxative products for longer than 1 week unless directed by a doctor • laxative products for longer than 1 week unless directed by a doctor • laxative products for longer than 1 week unless directed by a doctor • laxative. Stop use and ask a doctor if you have • stomach pain, nausea or vomiting • noticed a sudden change in bowel habits that lasts over two weeks Stop use and ask a doctor if you have rectal bleeding or fail to have a bowel movement after use of a laxative. These may indicate a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (800-222-1222).	
	Drug Facts Purpose Active ingredients (in each tablet) Purpose Docusate Sodium 50 mg. Stool Softener Sennosides 8.6 mg. Stool Softener Sennosides 8.6 mg. Laxative Uses • relieves occasional constipation (irregularity) • this product generally produces a bowel movement in 6 to 12 hours Warnings Do not use • if you are now taking mineral oil, unless directed by a doctor • laxative products for longer than 1 week unless directed by a doctor • laxative products for longer than 1 week unless directed by a doctor • laxative products for longer than 1 week unless directed by a doctor • laxative. These may indicate a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (800-222-1222). Drug Facts (continued) Directions • If taking the starting dosage, take preferably at bedtime; or as directed by a doctor.	
	Drug Facts Purpose Active ingredients (in each tablet) Purpose Docusate Sodium 50 mg. Stool Softener Sennosides 8.6 mg. Stool Softener Uses • relieves occasional constipation (irregularity) • this product generally produces a bowel movement in 6 to 12 hours Warnings Do not use • if you are now taking mineral oil, unless directed by a doctor • laxative products for longer than 1 week unless directed by a doctor • laxative products for longer than 1 week unless directed by a doctor • laxative products for longer than 1 week unless directed by a doctor • laxative products for longer than 1 week unless directed by a doctor • laxative products for longer than 1 week unless directed by a doctor • laxative. Out use • if you have • stomach pain, nausea or vomiting • noticed a sudden change in bowel habits that lasts over two weeks Stop use and ask a doctor if you have rectal bleeding or fail to have a bowel movement after use of a laxative. These may indicate a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (800-222-1222). Drug Facts (continued) Directions • If taking the starting dosage, take preferably at bedtime; or as directed by a doctor. <td></td>	
	Drug Facts Purpose Active ingredients (in each tablet) Purpose Docusate Sodium 50 mg. Stool Softener Sennosides 8.6 mg. Stool Softener Uses • relieves occasional constipation (irregularity) • this product generally produces a bowel movement in 6 to 12 hours Warnings Do not use • if you are now taking mineral oil, unless directed by a doctor • laxative products for longer than 1 week unless directed by a doctor • laxative products for longer than 1 week unless directed by a doctor • laxative products for longer than 1 week unless directed by a doctor • laxative products for longer than 1 week unless directed by a doctor • laxative products for longer than 1 week unless directed by a doctor • laxative products for longer than 1 week unless directed by a doctor • laxative products for longer than 1 week unless directed by a doctor • laxative products for longer than 1 week unless directed by a doctor • laxative the products for longer than 1 week unless directed by a doctor • noticed a sudden change in bowel habits that lasts over two weeks Stop use and ask a doctor if you have rectal bleeding or fail to have a bowel movement after use of a laxative. These may indicate a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (800-222-1222). Drug F	
	Drug Facts Purpose Docusate Sodium 50 mg. Stool Softener Sennosides 8.6 mg. Stool Softener Sennosides 8.6 mg. Stool Softener Uses • relieves occasional constipation (irregularity) • this product generally produces a bowel movement in 6 to 12 hours Warnings Do not use • if you are now taking mineral oil, unless directed by a doctor • laxative products for longer than 1 week unless directed by a doctor • laxative products for longer than 1 week unless directed by a doctor • laxative products for longer than 1 week unless directed by a doctor • laxative products for longer than 1 week unless directed by a doctor • laxative products for longer than 1 week unless directed by a doctor • laxative. These may indicate a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (800-222-1222). Drug Facts (continued) Directions • If taking the starting dosage, take preferably at bedtime; or as directed by a doctor. • If taking the maximum dosage, take in the morning and at bedtime. age starting dosage adults and children 12 years of age and over 2 tablets once a day 4 tablets twice a day	er distributed by Avio Health LP., EROKOT-SP
	Drug Facts Purpose Active ingredients (in each tablet) Purpose Docusate Sodium 50 mg. Stool Softener Sennosides 8.6 mg. Stool Softener Sennosides 8.6 mg. Laxative Uses • relieves occasional constipation (irregularity) • this product generally produces a bowel movement in 6 to 12 hours Warnings Do not use • if you are now taking mineral oil, unless directed by a doctor • laxative products for longer than 1 week unless directed by a doctor • laxative products for longer than 1 week unless directed by a doctor • laxative products for longer than 1 week unless directed by a doctor • laxative products for longer than 1 week unless directed by a doctor • laxative. These may indicate a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (800-222-1222). Drug Facts (continued) Directions • If taking the starting dosage, take preferably at bedtime; or as directed by a doctor. • If taking the maximum dosage, take in the morning and at bedtime. age starting dosage adults and children 12 years of age and over 2 tablets once a day 4 tablets twice a day children 6 to	er distributed by Avio Health LP., EROKOT-SP
	Drug Facts Purpose Active ingredients (in each tablet) Purpose Docusate Sodium 50 mg	er distributed by Avio Health LP., EROKOT-99 55597555
	Drug Facts Purpose Docusate Sodium 50 mg. Stool Softener Sennosides 8.6 mg. Stool Softener Uses • relieves occasional constipation (irregularity) • this product generally produces a bowel movement in 6 to 12 hours Warnings Do not use • if you are now taking mineral oil, unless directed by a doctor • laxative products for longer than 1 week unless directed by a doctor • laxative products for longer than 1 week unless directed by a doctor • laxative products for longer than 1 week unless directed by a doctor • laxative products for longer than 1 week unless directed by a doctor • laxative products for longer than 1 week unless directed by a doctor • laxative products for longer than 1 week unless directed by a doctor • laxative products a sudden change in bowel habits that lasts over two weeks Stop use and ask a doctor if you have estomach pain, nausea or vomiting • noticed a sudden change in bowel habits that lasts over two weeks Stop use and ask a doctor if you have rectal bleeding or fail to have a bowel movement after use of a laxative. These may indicate a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. n case of overdose, get medical help or contact a Poison Control Center right away (800-222-1222). Directions • If taking the starting dosage, take preferably at bedtime; or as directed by a doctor. • If taking the maximum dosage, take in the morning	androchured or dis inibuted by Anio Health LP., ad molecond: SR (2015) 3 9 1 9 5 5 9 7 5 5

SENNA-S

sennosides and docusate sodium tablet

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70677-1069

Active Ingred	ient/Acti	ive Moiety					
	Ir	ngredient Name	•	В	asis of S	trength	Strengt
SENNOSIDES (UNI	I: 3FYP5M0IJ	X) (SENNOSIDES - U	JNII:3FYP5M0IJX)	SE	NNOSIDES		8.6 mg
DOCUSATE SODIL	JM (UNII: F0	5Q2T2JA0) (DOCUSA	ATE - UNII:M7P2719	95AG) DO	CUSATE S	ODIUM	50 mg
Inactive Ingre	dients						
		Ingredie	nt Name			9	Strength
CELLULOSE, MICH	ROCRYSTA	LLINE (UNII: OP1R32	2D61U)				
CROSCARMELLOS	E SODIUM	(UNII: M28OL1HH48	3)				
CALCIUM PHOSPH	IATE, DIBA	SIC, ANHYDROUS	(UNII: L11K75P92J)				
MAGNESIUM STE	ARATE (UNI	: 70097M6I30)					
POLYETHYLENE G	LYCOL, UN	ISPECIFIED (UNII: 3	3WJQ0SDW1A)				
HYPROMELLOSES	(UNII: 3NX)	N29V3WO)					
FD&C YELLOW N	D. 6 (UNII: H	177VEI93A8)					
TITANIUM DIOXID	E (UNII: 15F	IX9V2JP)					
TALC (UNII: 7SEV7J	4R1U)						
SILICON DIOXIDE	(UNII: ETJ7Z	26XBU4)					
Product Char	acteristi						
	acteristi		Score			no score	
Color	acteristi	cs	Score Size			no score 9mm	
Color Shape	acteristi	CS orange	Size				
Product Char Color Shape Flavor Contains	acteristi	CS orange				9mm	
Color Shape Flavor	acteristi	CS orange	Size			9mm	
Color Shape Flavor	acteristi	CS orange	Size			9mm	
Color Shape Flavor Contains	acteristi	CS orange	Size			9mm	
Color Shape Flavor Contains Packaging	acteristi	CS orange	Size Imprint Code	Marketing Date	9 Start	9mm PSD22 Marke	ting End
Color Shape Flavor Contains Packaging	acteristi 1 in 1 CAF	CS orange ROUND	Size Imprint Code		9 Start	9mm PSD22 Marke	
Color Shape Flavor Contains Packaging # Item Code 1 NDC:70677- 1069-1	1 in 1 CAF	CS orange ROUND	Size Imprint Code	Date	9 Start	9mm PSD22 Marke	
Color Shape Flavor Contains Packaging # Item Code 1 NDC:70677- 1069-1	1 in 1 CAF 60 in 1 B0	CS orange ROUND Package Descr	Size Imprint Code	Date	9 Start	9mm PSD22 Marke	
Color Shape Flavor Contains Packaging # Item Code 1 NDC:70677- 1069-1 1	1 in 1 CAF 60 in 1 BC Product	CS orange ROUND Package Descr RTON DTTLE; Type 0: Not a	Size Imprint Code	Date	9 Start	9mm PSD22 Marke	
Color Shape Flavor Contains Packaging # Item Code 1 NDC:70677- 1069-1	1 in 1 CAF 60 in 1 BC Product Inform	CS orange ROUND Package Descr RTON DTTLE; Type 0: Not a	Size Imprint Code iption a Combination	Date 05/01/2023 Marketi	9 Start	9mm PSD22 Marke	
Color Shape Flavor Contains PE-Kaging Item Code NDC:70677- 1069-1 1 NDC:70677- 1069-1	1 in 1 CAF 60 in 1 BC Product Inform App	CS orange ROUND Package Descr RTON DTTLE; Type 0: Not a nation lication Number Citatio	Size Imprint Code iption a Combination	Date 05/01/2023 Marketi	g Start	9mm PSD22 Marke	Date eting End

Labeler - STRATEGIC SOURCING SERVICES LLC (116956644)

Registrant - Geri-Care Pharmaceutical Corp (611196254)