

**GFA FIRST AID 75 PERSON METAL CASE- benzalkonium chloride, benzalkonium chloride, lidocaine, bacitracin zinc, neomycin sulfate, polymyxin b sulfate, povidone-iodine, water, alcohol, aspirin, ibuprofen, acetaminophen, benzocaine, alcohol
Genuine First Aid, LLC**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

GFA First Aid Kit

Active Ingredient

Active Ingredient:

Benzalkonium Chloride 0.40%

Purpose

Antiseptic

Use

For Professional and Hospital use. Helps prevent infection. Antiseptic cleansing of face, hands and body without soap and water.

Warnings

Warning: For external use only.

Keep out of reach of children: If swallowed, get medical help or contact a Poison Control Center right away.

Stop use if unusual redness, swelling or other symptoms occur. Consult a physician immediately.

Do not use in the eyes or over large areas of the body.

Directions

Tear open packet, unfold towelette and use to cleanse desired skin area. Discard towelette appropriately after single use.

Inactive Ingredients

Inactive Ingredient: Purified water

LOT/EXP: Made in CHINA

20130301

Antiseptic Towelette

Genuine First Aid LLC, Clearwater FL 33755

www.GenuineFirstAid.com

1/pouch

GENUINE FIRST AID

Active Ingredients

Active Ingredient:Bacitracin Zinc 400 units

Neomycin Sulfate 5mg (equivalent to 3.5 mg Neomycin base)

Polymyxin B Sulfate 5000 units

Purpose

Triple Antibiotic

Uses: To help prevent infection in:
minor cuts; scrapes; burns

Warnings

For external use only.

Do not use: in eyes; over large areas of the body;

If allergic to any of the ingredients; for more than one week unless directed by a physician.

Stop use and consult a doctor:

if the condition persists or gets worse; a rash or other allergic reaction develops

Keep out of reach of children.

If ingested, contact a Poison

Control Center right away.

Directions

Directions: clean affected area; apply small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily; may be covered with a sterile bandage

Other information:

Store at room temperature.

Inactive Ingredient

water

Genuine Triple Antibiotic

First Aid Ointment

To Help Prevent Infection

Each Gram Contains:

Bacitracin Zinc 400 units

Neomycin Sulfate 5 mg

(equivalent to 3.5 mg
Neomycin base)
Polymyxin B Sulfate 5000 units
Net Wt. 0.5g ; (1/64 oz)
Manufactured in CHINA for
GENUINE FIRST AID.

Triple Antibiotic Ointment 10pcs
Net wt. 0.9g (1/32oz)

100
Triple Antibiotic

Active Ingredient

One Prep pad saturated with a 10% Povidone-Iodine solution equivalent to 1% available iodine

Purpose

antiseptic, germicidal

Uses

Provides, antiseptic germicidal skin preparation for minor invasive procedures

Keep out of reach of children

For external use only.

Caution

In case of deep or puncture wounds or serious burns, and if pain, irritation, redness, swelling or infection occurs, discontinue use and contact a physician.

Inactive Ingredients

Purified water.

Directions

Clean intended area thoroughly with pad. Discard after single use.

Antiseptic
Povidone-Iodine
Prep Pads
medium

Saturated with a 10% Povidone-Iodine
Solution equivalent to 1% available iodine

GFA Production Xiamen Co., Ltd.
www.gfaproduction.com

GFA Production Xiamen Co., Ltd.
No. 20 Huli Industrial Park, Meixi Road,
Tong'an, Xiamen, Fujian, China 361100

Wellkang Ltd t/a Wellkang Tech Consulting
Suite B, 29 Harley Street
LONDON W1G 9QR, England, United Kingdom

Active Ingredient

Ethyl Alcohol 62%

Purpose

Antimicrobial

Uses

For handwashing to decrease bacteria on the skin.

Warnings

For external use only.

Flammable

Do not use

in the eyes. In case of contact, rinse eyes thoroughly with water.

Stop use and consult a doctor

if irritation and redness develop and persist for more than 72 hours

Directions

Empty contents into palm. Rub hands until gel dissipates. Recommended for repeated use.

Inactive Ingredients

Propylene glycol, carbomer, titanium dioxide, purified water

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center right away.

Genuine Hand Sanitizer

Instant Gel

Antiseptic Handwash with Vitamin E and Aloe

Kills 99.9% of germs without water

Active Ingredient:

Ethyl alcohol 62%

Net Wt 0.9g (1/32oz)

Active Ingredients

ACTIVE INGREDIENTS:

Benzalkonium Chloride 0.13%

Lidocaine HCL 0.5%

Purpose

Purpose: First aid antiseptic, external analgesic

Uses

First aid to help prevent infection and for the temporary relief of pain and itching associated with:

Minor Cuts

Scrapes

Burns

Warnings

For external use only

Do not use: In eyes, in large quantities, over raw blistered areas, or on deep puncture wounds, animal bites or serious burns, for more than one week

Do not use:

in the eyes or apply over large areas of the body.

longer than 1 week unless directed by a doctor.

in large quantities, particularly over raw surfaces or blistered areas.

Ask a doctor before use if you have deep puncture wounds, animal bites or serious burns.

When using this product, avoid contact with the eyes.

Stop use and ask a doctor if
condition worsens

symptoms persist for more than 7 days

condition clears up and occurs again within a few days

Purpose

Pain reliever/fever reducer

Uses

temporarily relieves minor aches and pains due to:

the common cold

headache

toothache

muscular aches

backache

minor pain of arthritis

menstrual cramps temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: shock, facial swelling, asthma (wheezing) rash, skin reddening, blisters, hives. If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older, have had stomach ulcers or bleeding problems, take a blood thinner (anticoagulant) or steroid drug, take other drugs containing NSAIDs (aspirin, ibuprofen, naproxen, or others), have 3 or more alcoholic drinks every day while using this product, take more or for a longer time than directed

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer, right before or after heart surgery.

Ask a doctor before use if stomach bleeding warning applies to you; you have a history of stomach problems such as heartburn; you have a high blood pressure, heart disease, liver cirrhosis, or kidney disease; you are taking a diuretic

Ask a doctor before use if you are:

taking any other drug containing NSAID (prescription or nonprescription); taking aspirin for heart attack or stroke, because Ibuprofen may decrease this benefit of aspirin; taking any other drug

When using this product

take with food or milk if stomach upset occurs

Stop use and ask a doctor if

you experience any of the following signs of stomach bleeding; feel faint; vomit blood; have bloody or

black stools; have stomach pain that does get better; pain gets worse or lasts more than 10 days; fever gets worse or lasts more than 3 days; redness or swelling is present in the painful area; any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

do not use more than directed; the smallest effective dose should be used; do not take longer than 10 days, unless directed by a doctor.

Adults and Children (12 years and older): Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours, unless directed by a doctor.

Children under 12 years: Do not give to children under 12 years of age.

Other information

Store at controlled room temperature; avoid excessive heat 40 degree Celsius (104 degree Fahrenheit); tamper evident sealed packets; do not use any opened or torn packets

Inactive Ingredients

cellulose, corn starch, fumed silica gel, hypromellose, lactose, magnesium stearate, polydextrose, polyethylene glycol, povidone, silica, sodium starch glycolate, stearic acid, titanium dioxide, triacetin.

Distributed by GENUINE FIRST AID

600 Cleveland Str Suite 400, Clearwater, FL 33755

IBUPROFEN 2 Tablets

IBUPROFEN 2 Tablets

Active Ingredient (in each tablet)

Acetaminophen 325 mg

Purpose

Analgesic/antipyretic

Uses

temporary relief of minor aches and pains associated with:

common cold; headache; toothache; muscular aches; backache; arthritis; menstrual cramps; and reduction of fever

Warnings

Liver warning: This product contains acetaminophen.

Severe liver damage may occur if: adult takes more than 12 tablets in 24 hours, which is the maximum daily amount; child takes more than 5 doses in 24 hours; taken with other drugs containing acetaminophen; adult has 3 or more alcoholic drinks every day while using this product

Do not use

with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist; for more than 10 days for pain unless directed by a doctor; for more than 3 days for fever unless directed by a doctor

Ask a doctor before use if the user has liver disease

Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin

Stop use and ask a doctor if: symptoms do not improve; pain gets worse or lasts for more than 10 days; fever gets worse or lasts for more than 3 days; new symptoms occur; redness or swelling is present; a rare sensitivity reaction occurs

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of accidental overdose, contact a doctor or Poison Control Center immediately. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. Do not exceed recommended dosage

Inactive Ingredients

Cornstarch, polyethylene glycol, stearic acid, povidone

Directions

Adults and Children 12 years of age or older	Take 2 tablets every 4 to 6 hours as needed. Do not take more than 12 tablets in 24 hours.
--	--

Children 6-11 years of age	Take 1 tablet every 4 to 6 hours as needed. Do not take more than 5 tablets in 24 hours.
-------------------------------	--

Children under 6 years of age	Do not use this regular strength product. This will provide more than the recommended dose (overdose) and could cause serious health problems.
----------------------------------	---

Store at 59-86 degree F (15-30 degree C)

tamper evident sealed packets; do not use any open or torn packets

Distributed by GENUINE FIRST AID
600 Cleveland Str Suite 400, Clearwater, FL 33755

GENUINE FIRST AID 2 Tablets

NON-ASPIRIN

Active Ingredient (in each tablet)

Aspirin (NSAID*) 325 mg
*nonsteroidal anti-inflammatory drug

Purpose

Pain Reliever / fever reducer

Uses

Temporarily relieves minor aches and pains associated with:
headache ; muscular aches ; minor arthritis pain ; backache ; common cold ; toothache ; menstrual cramps
; Temporarily reduces fever

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: hives, skin reddening, facial swelling, rash, asthma (wheezing), blisters, shock, If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
are age 60 or older; have had stomach ulcers or bleeding problems; take a blood thinner (anticoagulant) or steroid drug; take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others); have 3 or more alcoholic drinks every day while using this product; take more or for a longer time than directed

Do not use

if you have ever had an allergic reaction to any other pain reliever/ fever reducer; right before or after heart surgery; if you are taking a prescription drug for gout, diabetes or arthritis

Ask a doctor before use if: stomach bleeding warning applies to you; you have a history of stomach problems such as heartburn; you have high blood pressure, heart disease, liver cirrhosis, or kidney disease; you are taking a diuretic

Ask a doctor or pharmacist before use if you are:

under a doctor's care for any serious condition; taking any other drug

When using this product: take with food or milk if stomach upset occurs

Stop use and ask a doctor if

Stop use and ask a doctor if

you experience any of the following signs of stomach bleeding:

feel faint; vomit blood; have bloody or black stools; have stomach pain that does not get better; pain gets worse or lasts more than 10 days; fever gets worse or lasts more than 3 days; you have difficulty swallowing; if ringing in the ears or loss of hearing occurs; redness or swelling is present in the painful areas; any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

do not use more than directed

the smallest effective dose should be used

drink a full glass of water with each dose

do not take longer than 10 days, unless directed by a doctor

Adults and children: (12 years and older) Take 1 or 2 tablets with water every 4 hours as needed. Do not take more than 12 tablets in 24 hours, or as directed by a doctor.

Children under 12 years: Do not give to children under 12 years of age.

Store at 59 - 86 degree Fahrenheit (15 - 30 degree Celsius); avoid excessive heat and humidity; tamper evident sealed packets;

Do not use any opened or torn packets

Inactive Ingredients

hypromellose, polyethylene glycol, propylene glycol, corn starch

Distributed by GENUINE FIRST AID

600 Cleveland Str Suite 400, Clearwater, FL 33755

ASPIRIN 2 Tablets

Active Ingredient:

Purified Water USP...q.s.

Inactive Ingredients:

Sodium Chloride USP 44mg

Monobasic Sodium Phosphate USP 18mg

Sodium Phosphate Dibasic USP 111mg

Edetate Disodium USP 10mg

Benzalkonium Chloride
NF (as preservative)

0.5mg

Store in a cool place. For irrigation only.

Discard unused portion of the solution.

Not for injection.

Warning:

If you experience eye pain, changes in vision, continued redness or irritation of the eye, or if the condition worsens or persists for more than 72 hours, discontinue use and consult a doctor. Do not use if solution changes color or becomes cloudy.

Directions

Remove contacts before using.

Twist top to remove.

Flush the affected area as needed. Control

Rate of flow by pressure on the bottle. Do not touch tip of the container to any surface. Do not reuse.

If necessary continue flushing with emergency eyewash or shower.

Discard bottle after use.

Uses:

For flushing or irrigating the eyes to remove loose foreign material, air pollutants, or chlorinated water.

Code No.: GUJ/DRUG/G/1080

Batch No.:

Mfg Date:

Exp: Date:

10ml

Sterile Isotonic Buffered Genuine

Eyewash

For single use only

Active Ingredient

Active Ingredient:

Isopropyl Alcohol, 70% v/v

Purpose

Antiseptic

Use

For preparation of skin before injection.

Warnings

For external use only.

Flammable - keep away from fire or flame

Store at room temperature 15-30 degree Celsius (59-86 degree Fahrenheit)

Do not use: with electrocautery, in the eyes.

Stop use if irritation and redness develop. If condition persists for more than 72 hours, consult your doctor.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

Wipe Injection site vigorously and discard.

Inactive Ingredient

Purified water.

LOT/EXP: Made in CHINA

20140301

Alcohol Cleansing Pad

Genuine First Aid LLC, Clearwater FL 33755

www.GenuineFirstAid.com

1/pouch

GENUINE FIRST AID

100 Alcohol cleansing pads

Reorder GFAP-39-02

Active Ingredient

Active Ingredient:

Benzocaine, 6% w/v

SD alcohol, 60% w/v

Purpose:

Purpose

Topical Anesthetic

Antiseptic

Uses

Use: For the temporary relief of pain and itching associated with minor burns, scrapes and insect bites.

Directions

Clean intended area thoroughly with pad. Discard after single use.

Warnings

Warnings: For external use only.

Avoid contact with eyes. If this happens, rinse thoroughly with water.

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center right away.

Inactive Ingredients

Purified water

Flammable - keep away from fire or flame.

Do not use: In eyes, on broken skin, deep puncture wounds. If unusual redness, swelling, irritation or other symptoms occur, consult a physician immediately.

Made in CHINA

LOT/EXP:

Insect Sting Relief Pad

Genuine First Aid LLC, Clearwater FL 33755

www.GenuineFirstAid.com

1/pouch

GENUINE FIRST AID



GENUINE FIRST AID®



ISO 9001:2010



Genuine First Aid LLC, Clearwater FL 33755
www.GenuineFirstAid.com

**Insect Sting
Relief Pad**

**Toallitas para
Picaduras de Insectos**

REORDER ISRP-001

DRUG FACTS - Insect Sting Relief Pad

Active Ingredient: Benzocaine, 6% w/v Topical Anesthetic

Purpose: SD alcohol, 60% w/v Antiseptic

Use: For the temporary relief of pain and itching associated with minor burns, scrapes and insect bites.

Warnings: For external use only. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. Flammable - keep away from fire or flame. Avoid contact with eyes. If this happens, rinse thoroughly with water.

Do not use: In eyes, on broken skin, deep puncture wounds. If unusual redness, swelling, irritation or other symptoms occur, consult a physician immediately.

--- TEAR HERE ---

Made in CHINA

REORDER ACP-001

**Alcohol Cleansing
Pad**

**Toallitas Humedas
con Alcohol**



ISO 9001:2010



Genuine First Aid LLC, Clearwater FL 33755
www.GenuineFirstAid.com



GENUINE FIRST AID®



DRUG FACTS - Alcohol Cleansing Pad

Active Ingredient: Isopropyl Alcohol, 70% v/v
Purpose: Antiseptic

Use: For preparation of the skin before injection.

Warnings: For external use only.

Flammable - keep away from fire or flame.

Do not use: with electrocautery, in the eyes

Stop use if irritation and redness develop.

If condition persists for more than 72 hours,
consult your doctor.

Keep out of reach of children. If swallowed,
get medical help or contact a Poison Control
Center right away.

Directions: Wipe injection site vigorously and
discard.

Other information: Store at room temperature
15°-30° C (59°-86° F)

Inactive ingredient: Purified water.

----- TEAR HERE -----

LOT/EXP:

Made in CHINA

20130301

10ml

Sterile Isotonic buffered
GENUINE
Eyewash

For single use only

Directions: Remove contact lens before using. Twist top to remove. Flush the affected area as needed. Control rate of flow by pressure on the bottle. Do not touch tip of the container to any surface. Do not reuse. If necessary, continue flushing with emergency eyewash or shower. Discard bottle after use.

Uses: For flushing or irrigating the eyes to remove loose foreign material, air pollutants, or chlorinated water.

Each 10 ml contains:

Active Ingredient:
Purified Water ...q.s.

Inactive Ingredients:
Sodium Chloride
Monobasic Sodium Phosphate
Dibasic Sodium Phosphate
Ectate Disodium
Benzalkonium Chloride
NF (as preservative)

Store in a cool place. For irrigation only. Discard unused portion of the solution. Not for injection.

Mfg. Lic. No.: KTK/28/332/2002
Batch No.:
Mfg. date:
Exp. date:



Manufactured in INDIA for
GENUINE FIRST AID

Warning:
If you experience eye pain, changes in vision, continued redness or irritation of the eye, or if the condition worsens or persists for more than 72 hours, discontinue use and consult a doctor. Do not use if solution changes color or becomes cloudy.



REORDER
EYEW-10S

FOR EXTERNAL USE ONLY

STERILE A

ASPIRIN

2 Tablets

Active ingredient (in each tablet) **Purpose**
Aspirin (NSAID)* 325 mg Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses Temporarily relieves minor aches and pains associated with
■ headache ■ muscular aches ■ minor arthritis pain ■ backache
■ common cold ■ toothache ■ menstrual cramps

Temporarily reduces fever

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ skin reddening ■ facial swelling ■ rash
■ asthma (wheezing) ■ blisters ■ shock
If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older ■ have had stomach ulcers or bleeding problems
■ take a blood thinner (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery ■ if you are taking a prescription drug for gout, diabetes or arthritis

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have a history of stomach problems such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic

Warnings (continued)

Ask a doctor or pharmacist before use if you are
■ under a doctor's care for any serious condition ■ taking any other drug
When using this product ■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if

■ you experience any of the following signs of stomach bleeding:
■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days ■ you have difficulty swallowing
■ if ringing in the ears or loss of hearing occurs ■ redness or swelling is present in the painful area ■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

■ do not use more than directed
■ the smallest effective dose should be used
■ drink a full glass of water with each dose
■ do not take longer than 10 days, unless directed by a doctor

Adults and children: (12 years and older) Take 1 or 2 tablets with water every 4 hours as needed. Do not take more than 12 tablets in 24 hours, or as directed by a doctor.

Children under 12 years: Do not give to children under 12 years of age.

Other information ■ store at 59°-86°F (15°-30°C) ■ avoid excessive heat and humidity ■ tamper evident sealed packets ■ do not use any opened or torn packets

Inactive ingredients hypromellose, polyethylene glycol, propylene glycol, corn starch

Distributed by GENUINE FIRST AID
600 Cleveland Str Suite 400, Clearwater, FL 33755



GENUINE FIRST AID. 2 Tablets

NON-ASPIRIN

Active ingredient (in each tablet) Purpose
Acetaminophen 325 mg Analgesic/antipyretic

Uses

temporary relief of minor aches and pains associated with

- common cold ■ headache ■ toothache
- muscular aches ■ backache ■ arthritis
- menstrual cramps ■ and reduction of fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if ■ adult takes more than 12 tablets in 24 hours, which is the maximum daily amount ■ child takes more than 5 doses in 24 hours ■ taken with other drugs containing acetaminophen ■ adult has 3 or more alcoholic drinks every day while using this product

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist ■ for more than 10 days for pain unless directed by a doctor ■ for more than 3 days for fever unless directed by a doctor

Ask a doctor before use if the user has liver disease

Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin ▶

Warnings (continued)

Stop use and ask a doctor if ■ symptoms do not improve ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts for more than 3 days ■ new symptoms occur ■ redness or swelling is present ■ a rare sensitivity reaction occurs

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of accidental overdose, contact a doctor or Poison Control Center immediately. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. **Do not exceed recommended dosage.**

Directions

Adults and children 12 years of age and older	Take 2 tablets every 4 to 6 hours as needed. Do not take more than 12 tablets in 24 hours.
Children 6-11 years of age	Take 1 tablet every 4 to 6 hours as needed. Do not take more than 5 tablets in 24 hours.
Children under 6 years of age	Do not use this regular strength product. This will provide more than the recommended dose (overdose) and could cause serious health problems.

Other information ■ store at 59°-86°F (15°-30°C) ■ tamper evident sealed packets ■ do not use any open or torn packets

Inactive ingredients corn starch, hydroxypropyl methylcellulose, polyethylene glycol, pregelatinized starch, stearic acid. May contain povidone and sodium starch glycolates.

Distributed by GENUINE FIRST AID
600 Cleveland Str Suite 400, Clearwater, FL 33755

IBUPROFEN 2 Tablets

Active ingredient (in each tablet) Purpose
Ibuprofen USP (NSAID)* 200mg Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses

temporarily relieves minor aches and pains due to:
■ the common cold ■ headache ■ toothache ■ muscular aches
■ backache ■ minor pain of arthritis ■ menstrual cramps
temporarily reduces fever

Warnings Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ shock ■ facial swelling ■ asthma (wheezing) ■ rash ■ skin reddening ■ blisters ■ hives If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinner (anticoagulant) or steroid drug ■ take other drugs containing NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have a history of stomach problems such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are ■ taking any other drug containing an NSAID (prescription or nonprescription) ▶

Warnings (continued)

■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug

When using this product

■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ do not use more than directed ■ the smallest effective dose should be used ■ do not take longer than 10 days, unless directed by a doctor

Adults and Children (12 years and older): Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours, unless directed by a doctor.

Children under 12 years: Do not give to children under 12 years of age.

Other information ■ store at controlled room temperature ■ avoid excessive heat 40°C (104°F) ■ tamper evident sealed packets ■ do not use any opened or torn packets

Inactive ingredients cellulose, corn starch, fumed silica gel, hypromellose, lactose, magnesium stearate, polydextrose, polyethylene glycol, povidone, silica, sodium starch glycolate, stearic acid, titanium dioxide, triacetin

Distributed by GENUINE FIRST AID
600 Cleveland Str Suite 400, Clearwater, FL 33755

REORDER GFAP-29-06

Genuine Hand Sanitizer

Instant Gel Antiseptic Handwash with Vitamin E & Aloe

Kills 99.9% of Germs Without Water
Active Ingredient: Ethyl alcohol 62%
Net Wt 0.9g - (1/32 oz)



GFA Production Xiamen Co., Ltd
No. 20 Huli Industrial Park, Meizi Road,
Tonglan, Xiamen, Fujian, China 361100
Wellkang Ltd is Wellkang Tech Consulting
Suite B, 29 Harley Street
LONDON W1G 9QR, England, United Kingdom

Drug Facts

Uses: For hand washing to decrease bacteria on the skin

Warnings:

For external use only

Flammable, keep away from fire or flame

Do not use in the eyes. In case of contact, rinse eyes thoroughly with water

Stop use and consult a doctor if irritation and redness develop and persist for more than 72 hours

Directions: Empty contents into palm. Rub hands until gel dissipates. Recommended for repeated use

Other information: Store at 15° to 25° C (59° to 77° F)

Inactive Ingredients: Propylene glycol, Carbomer, Titaniumdioxide, Purified water.

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center right away.

LOT/EXP:
20140115

REORDER POV-001

Antiseptic Povidone-Iodine Prep Pads

medium
Saturated with a 10% Povidone-Iodine Solution equivalent to 1% available iodine



GFA Production Xiamen Co., Ltd
www.gfaproduction.com



GFA Production Xiamen Co., Ltd
No. 20 Huli Industrial Park, Meizi Road,
Tonglan, Xiamen, Fujian, China 361100
Wellkang Ltd is Wellkang Tech Consulting
Suite B, 29 Harley Street
LONDON W1G 9QR, England, United Kingdom



For External Use Only

CONTENTS: One Prep Pad saturated with a 10% Povidone-Iodine solution equivalent to 1% available iodine.

Uses: Provides antiseptic, germicidal skin preparation for minor invasive procedures.

CAUTION: In case of deep or puncture wounds or serious burns, and if pain, irritation, redness, swelling or infection occurs, discontinue use and contact a physician.

Inactive Ingredients: Purified water.

DIRECTIONS: Clean intended area thoroughly with pad. Discard after single use.

Keep out of reach of children

LOT/EXP:
20140325

--- TEAR HERE ---

Reorder TAO-001

Genuine Triple Antibiotic

**First Aid Ointment
To Help Prevent Infection**

**Each Gram Contains:
Bacitracin Zinc 400 units
Neomycin Sulfate 5 mg
(equivalent to 3.5 mg
Neomycin base)
Polymyxin B Sulfate 5000 units
Net Wt 0.5g • (1/64 oz)**



Manufactured in CHINA for
GENUINE FIRST AID.

Drug Facts

Uses: To help prevent infection
in • minor cuts • scrapes • burns

Warnings:

For external use only

Do not use • in eyes • over large
areas of the body • if allergic to
any of the ingredients • for more
than one week unless directed
by a physician

Stop use and consult a doctor
• if the condition persists or gets
worse • a rash or other allergic
reaction develops

Keep out of reach of children.

If ingested, contact a Poison
Control Center right away.

Directions: • clean affected area
• apply small amount of this product
(an amount equal to the surface
area of the tip of a finger) on the
area 1 to 3 times daily • may be
covered with a sterile bandage

Other information:

Store at room temperature

LOT/EXP:

20130118

Reorder BC-001

Genuine First Aid Burn Cream

**Antiseptic Pain Relief
With Aloe**

ACTIVE INGREDIENTS:

Benzalkonium

Chloride 0.13%

Lidocaine HCL 0.5%

Net Wt 0.9g · (1/32 oz)



Manufactured in CHINA for
GENUINE FIRST AID.

Drug Facts

Purpose: First aid antiseptic,
external analgesic

Uses: First aid to help prevent
infection and for the temporary
relief of pain and itching

associated with: • minor cuts
• scrapes • burns

Warnings:

For external use only

Do not use • in eyes • in large
quantities • over raw blistered
areas, or on deep puncture
wounds, animal bites or serious
burns • for more than one week
Keep out of reach of children.

If ingested, contact a Poison
Control Center right away.

Directions: • clean affected area
• apply small amount not more
than 3 times daily • may be
covered with a sterile bandage

Other information:

Store at room temperature

LOT/EXP:

20130301

REORDER AST-001

**Antiseptic
Towelette**
**Toallitas
Antisepicas**



Genuine First Aid LLC, Clearwater FL 33755
www.GenuineFirstAid.com



GENUINE FIRST AID®



DRUG FACTS - Antiseptic Towelette

Active Ingredient: Benzalkonium Chloride 0.40% Purpose: First Aid Antiseptic

Use: For Professional and Hospital use. Helps prevent infection. Antiseptic cleansing of face, hands and body without soap and water.

Warnings: For external use only. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. If unusual redness, swelling or other symptoms occur, consult a physician immediately.

Do not use: In the eyes, or over large areas of the body.

Directions: Tear open packet, unfold towelette and use to cleanse desired skin area. Discard towelette appropriately after single use.

Inactive ingredient: Purified water

Made in CHINA

LOT/EXP:
20130301

--- TEAR HERE ---



**EXCEEDS
OSHA/ANSI/ISEA
Z308.1-2009.**

FIRST AID KIT BULK

ANSI/ISEA Z308.1-2009



**75
Person**

ANSI/ISEA Z308.1-2009 TYPE ■

Caution! This kit meets ANSI/ISEA Z308.1-2009 only when required minimum fill is maintained with first aid products marked "ANSI/ISEA Z308.1-2009."

Hard Case

1 Hardcase 36 unit	1	1 Sterile Eye Pad	2
1 GFA Easy Access Pocket System 50P	1	1 Sterile Eye Wash 10 ml	2
1 Hand Sanitizer 0.9 g	5	1 Butterfly Wound Cloures	1
1 Adhesive Plastic Bandages 3/4" x 3" Pocket 1	100	1 First Aid Tape Roll 1/2" x 2.5 yd	2
1 PP Adhesive Plastic Bandages 1" x 3"	200	1 First Aid Tape Roll 1/2" x 5 yd	2
1 PP Knuckle Fabric Bandage	18	1 Combine Pad 5" x 9"	2
1 PP Adhesive Spot Bandages 7/8" x 7/8"	125	1 Combine Pad 8" x 10"	2
1 PP Fingertip Fabric Bandage	18	1 Roller Gauze Bandage 3" x 4,1 yds	2
1 PP Elbow/Knee Adhesive Bandage	2	1 CPR Breathing Barrier	1
1 Ziplock Bag for Bandages, BZKs etc 50	1	1 Medical Grade Vinyl Gloves	8
1 Burn Cream	12	1 Alcohol Cleansing Pads	30
1 Antiseptic Towelettes	50	1 Insect Sting Relief Pads	12
1 Triple Antibiotic Ointment 0.9 g	10	1 Povidone Wipes	10
1 Cotton Tipped Applicators in wrapper	40	1 Triangular Bandage 42" x 42" x 59" nonwoven	3
1 Ibuprofen 200 mg 2/paper poly 2 years	2	1 Instant Cold Compress	3
1 Non Aspirin 325 mg 2/paper poly 3 years	2	1 Emergency First Aid Guide	1
1 Aspirin 325 mg 325 mg 2/paper poly 3 years	4	1 Plastic Tweezers	1
1 Pressure Bandages 2" x 2"	4	1 Assorted Safety Pins	10
1 Pressure Bandages 3" x 3"	2	1 Scissors	1
1 Blood Stopper Dressing	2	1 Wooden Finger Splints	2
1 Sterile Gauze Pad 2" x 2"	4		
1 Sterile Gauze Pad 3" x 3"	4		
1 Sterile Gauze Pad 4" x 4"	4		
1 Roller Gauze Bandage 2" x 4,1 yds	2		

Manufactured in China for:
Genuine First Aid LLC,
600 Cleveland Street,
Suite 400
Clearwater, FL 33755
www.GenuineFirstAid.com

GENUINE FIRST AID.



Copyright © 2010 Genuine First Aid LLC.
All rights reserved.

GFAP-21-04

GFA FIRST AID 75 PERSON METAL CASE

benzalkonium chloride, benzalkonium chloride, lidocaine, bacitracin zinc, neomycin sulfate, polymyxin b sulfate, povidone-iodine, water, alcohol, aspirin, ibuprofen, acetaminophen, benzocaine, alcohol kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:52124-1154
---------------------	----------------	---------------------------	----------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-1154-1	1 in 1 KIT		

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	50 PACKAGE	40.0 mL
Part 2	12 PACKET	10.8 g
Part 3	10 TUBE	5.0 g
Part 4	10 PACKET	5 mL
Part 5	2 BOTTLE	20 mL
Part 6	5 PACKET	4.5 g
Part 7	30 PACKAGE	15 mL
Part 8	12 PACKAGE	6 mL
Part 9	2 PACKET	4
Part 10	2 PACKAGE	4
Part 11	4 PACKAGE	8

Part 1 of 11**ANTISEPTIC TOWELETTE**

benzalkonium chloride swab

Product Information

Item Code (Source)	NDC:52124-0001
---------------------------	----------------

Route of Administration	TOPICAL
--------------------------------	---------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7) (BENZALKONIUM - UNII:7N6JUD5X6Y)	BENZALKONIUM CHLORIDE	0.4 mL in 100 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-0001-1	0.8 mL in 1 PACKAGE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part333E	05/03/2011	

Part 2 of 11

GENUINE FIRST AID BURN ANTISEPTIC PAIN RELIEF WITH ALOE

benzalkonium chloride, lidocaine cream

Product Information

Item Code (Source)	NDC:52124-0004
Route of Administration	TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7) (BENZALKONIUM - UNII:7N6JUD5X6Y)	BENZALKONIUM CHLORIDE	0.13 g in 100 g
LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	0.5 g in 100 g

Inactive Ingredients

Ingredient Name	Strength
ALOE VERA LEAF (UNII: ZY81Z83H0X)	
CETYL ALCOHOL (UNII: 936JST6JCN)	
DIAZOLIDINYL UREA (UNII: H5RIZ3MPW4)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
GLYCERIN (UNII: PDC6A3C0OX)	
GLYCERYL MONOSTEARATE (UNII: 230OU9XXE4)	
METHYL PARABEN (UNII: A2I8C7HI9T)	
MINERAL OIL (UNII: T5L8T28FGP)	
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYL PARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-0004-1	0.9 g in 1 PACKET		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part333A	04/13/2011	

Part 3 of 11

GENUINE TRIPLE ANTIBIOTIC

bacitracin zinc,neomycin sulfate,polymyxin b sulfate ointment

Product Information

Item Code (Source)	NDC:52124-0003
Route of Administration	TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BACITRACIN ZINC (UNII: 89Y4M234ES) (BACITRACIN - UNII:58H6RWO52I)	BACITRACIN ZINC	400 [iU] in 1 g
NEOMYCIN SULFATE (UNII: 057Y626693) (NEOMYCIN - UNII:I16QD7X297)	NEOMYCIN SULFATE	5 mg in 1 g
POLYMYXIN B SULFATE (UNII: 19371312D4) (POLYMYXIN B - UNII:J2VZ07J96K)	POLYMYXIN B SULFATE	5000 [iU] in 1 g

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-0003-1	0.5 g in 1 TUBE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part333B	04/13/2011	

Part 4 of 11

POVIDONE-IODINE PREP

povidone-iodine swab

Product Information

Item Code (Source)	NDC:52124-2901
Route of Administration	TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
POVIDONE-IODINE (UNII: 85H0HZU99M) (IODINE - UNII:9679TC07X4)	IODINE	1 mL in 100 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0K00R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-2901-0	0.5 mL in 1 PACKET		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part333E	05/03/2011	

Part 5 of 11

STERILE ISOTONIC BUFFERED GENUINE EYEWASH

water liquid

Product Information

Item Code (Source)	NDC:52124-0005
Route of Administration	OPHTHALMIC

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
WATER (UNII: 059QF0K00R) (WATER - UNII:059QF0K00R)	WATER	98.16 mL in 100 mL

Inactive Ingredients

Ingredient Name	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM PHOSPHATE, MONOBASIC (UNII: 3980JIH2SW)	
SODIUM PHOSPHATE, DIBASIC (UNII: GR686LBA74)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-0005-1	10 mL in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part349	05/04/2011	

Part 6 of 11

GENUINE HAND SANITIZER

alcohol gel

Product Information

Item Code (Source)	NDC:52124-2906
Route of Administration	TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALCOHOL (UNII: 3K9958V90M) (ALCOHOL - UNII:3K9958V90M)	ALCOHOL	62 g in 100 g

Inactive Ingredients

Ingredient Name	Strength
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
WATER (UNII: 059QF0K00R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-2906-1	0.9 g in 1 PACKET		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part333A	03/28/2011	

Part 7 of 11

ALCOHOL CLEANSING PAD

isopropyl alcohol liquid

Product Information

Item Code (Source)	NDC:52124-0002
Route of Administration	TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ISOPROPYL ALCOHOL (UNII: ND2M416302) (ISOPROPYL ALCOHOL - UNII:ND2M416302)	ISOPROPYL ALCOHOL	70 mL in 100 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-0002-1	0.5 mL in 1 PACKAGE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part333A	04/23/2010	

Part 8 of 11

INSECT STING RELIEF PAD

benzocaine, alcohol swab

Product Information

Item Code (Source)	NDC:52124-0008
Route of Administration	TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	6 mL in 100 mL
ALCOHOL (UNII: 3K9958V90M) (ALCOHOL - UNII:3K9958V90M)	ALCOHOL	60 mL in 100 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0K00R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-0008-1	0.5 mL in 1 PACKAGE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part348	04/23/2011	

Part 9 of 11

IBUPROFEN
ibuprofen tablet

Product Information

Item Code (Source)	NDC:52124-0009
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
POWDERED CELLULOSE (UNII: SMD1X3XO9M)	
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	

LACTOSE (UNII: J2B2A4N98G)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	

Product Characteristics

Color	white (White)	Score	no score
Shape	ROUND	Size	10 mm
Flavor		Imprint Code	44;352
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-0009-1	2 in 1 PACKET		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075010	02/17/2010	

Part 10 of 11

NON-ASPIRIN
acetaminophen tablet

Product Information

Item Code (Source)	NDC:52124-0010
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

Inactive Ingredients

Ingredient Name	Strength
-----------------	----------

STARCH, CORN (UNII: O8232NY3SJ)	
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
POVIDONE (UNII: FZ989GH94E)	

Product Characteristics

Color	white (White)	Score	no score
Shape	ROUND	Size	11mm
Flavor		Imprint Code	AZ;234
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-0010-1	2 in 1 PACKAGE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	04/23/2010	

Part 11 of 11

ASPIRIN

aspirin tablet

Product Information

Item Code (Source)	NDC:52124-0011
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSES (UNII: 3NXW29V3WO)	
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
STARCH, CORN (UNII: O8232NY3SJ)	

Product Characteristics

Color	white (White)	Score	no score
Shape	ROUND	Size	11mm
Flavor		Imprint Code	44;157;ASPIRIN
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-0011-1	2 in 1 PACKAGE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part343	04/23/2010	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part333E	05/03/2011	

Labeler - Genuine First Aid, LLC (619609857)