

**DIAZEPAM- diazepam injection**  
**Meridian Medical Technologies, Inc.**

-----  
**DIAZEPAM AUTOINJECTOR C-IV**

**Rx Only**

**WARNING: RISKS FROM CONCOMITANT USE WITH OPIOIDS; ABUSE, MISUSE, AND ADDICTION; and DEPENDENCE AND WITHDRAWAL REACTIONS**


- **Concomitant use of benzodiazepines and opioids may result in profound sedation, respiratory depression, coma, and death. Reserve concomitant prescribing of these drugs for patients for whom alternative treatment options are inadequate. Limit dosages and durations to the minimum required. Follow patients for signs and symptoms of respiratory depression and sedation (see WARNINGS and PRECAUTIONS).**
- **The use of benzodiazepines, including Diazepam injection, exposes users to risks of abuse, misuse, and addiction, which can lead to overdose or death. Abuse and misuse of benzodiazepines commonly involve concomitant use of other medications, alcohol, and/or illicit substances, which is associated with an increased frequency of serious adverse outcomes. Before prescribing Diazepam injection and throughout treatment, assess each patient's risk for abuse, misuse, and addiction (see WARNINGS).**
- **The continued use of benzodiazepines may lead to clinically significant physical dependence. The risks of dependence and withdrawal increase with longer treatment duration and higher daily dose. Although Diazepam injection is indicated only for intermittent use (see INDICATIONS AND USAGE and DOSAGE AND ADMINISTRATION), if used more frequently than recommended, abrupt discontinuation or rapid dosage reduction of DIAZEPAM INJECTION may precipitate acute withdrawal reactions, which can be life-threatening. For patients using DIAZEPAM INJECTION more frequently than recommended, to reduce the risk of withdrawal reactions, use a gradual taper to discontinue DIAZEPAM INJECTION (see WARNINGS).**

**DESCRIPTION**

Diazepam injection is a sterile solution packaged within a device that delivers its entire 2 mL contents automatically upon activation. Each mL contains 5 mg diazepam compounded with 40% propylene glycol, 10% ethyl alcohol, 5% sodium benzoate and benzoic acid as buffers, and 1.5% benzyl alcohol as preservative.

Diazepam is a benzodiazepine derivative. Chemically, diazepam is 7-chloro-1,3- dihydro-1-methyl-5-phenyl-2H-1, 4-benzodiazepin-2-one. It is a colorless crystalline compound, insoluble in water and has a molecular weight of 284.74. Its structural formula is as

follows:



## CLINICAL PHARMACOLOGY

In animals, diazepam appears to act on parts of the limbic system, the thalamus and hypothalamus, and induces calming effects. Diazepam, unlike chlorpromazine and reserpine, has no demonstrable peripheral autonomic blocking action, nor does it produce extrapyramidal side effects. However, animals treated with diazepam do have a transient ataxia at higher doses. Diazepam was found to have transient cardiovascular depressor effects in dogs. Long-term experiments in rats revealed no disturbances of endocrine function. Injections into animals have produced localized irritation of tissue surrounding injection sites and some thickening of veins after intravenous use.

### Pharmacokinetics (autoinjector)

A study performed in 24 healthy male subjects comparing the I.M. Injection of 10 mg of diazepam in the mid-anterior/lateral thigh by the autoinjector versus 10 mg I.M. by a syringe (operated manually) indicates that the mean percent availability of the drug from the autoinjector is 100% of that obtained from the syringe.

In addition, the mean  $C_{max}$  value from the autoinjector was 314 ng/mL (c.v. = 18.7%, range 185 to 439 ng/mL) and 48.6 ng/mL (c.v. = 19.8%, range 29.4 to 69.7 ng/mL) for diazepam and desmethyldiazepam, respectively, while the syringe gave corresponding values of 287 ng/mL (c.v. = 18.9%, range 174 to 378 ng/mL) and 47.2 ng/mL (c.v. = 19.4%, range 33.1 to 61.2 ng/mL) for diazepam and desmethyldiazepam, respectively. The corresponding mean  $T_{max}$  values were 1.47 hours (c.v. = 69.9%, range 0.8 to 6 hours) and 61.0 hours (c.v. = 56.8%, range 24 to 144 hours) for diazepam and desmethyldiazepam for the autoinjector whereas the syringe gave values of 1.31 hours (c.v. = 32%, range 0.7 to 2.0 hours) and 54.5 hours (c.v. = 47.3%, range 12 to 96 hours) for diazepam and desmethyldiazepam, respectively.

## INDICATIONS

Diazepam is indicated for the management of anxiety disorders for the short-term relief of the symptoms of anxiety. Anxiety or tension associated with the stress of everyday life usually does not require treatment with an anxiolytic.

Note: Because the autoinjector provides a minimum dose of 10 mg diazepam, it should not be used to treat individuals with mild and moderate degrees of anxiety and anxiety related disorders that would ordinarily be managed with intramuscular doses of less than 10 mg.

In acute alcohol withdrawal, diazepam may be useful in the symptomatic relief of acute agitation, tremor, impending or acute delirium tremens and hallucinosis.

As an adjunct prior to endoscopic procedures if apprehension, anxiety or acute stress

reactions are present, and to diminish the patient's recall of the procedures (See WARNINGS).

Diazepam is a useful adjunct for the relief of skeletal muscle spasm due to reflex spasm to local pathology (such as inflammation of the muscles or joints, or secondary to trauma), spasticity caused by upper motor neuron disorders (such as cerebral palsy and paraplegia); athetosis; stiff man syndrome; and tetanus.

Diazepam injection is a useful adjunct in status epilepticus and severe recurrent convulsive seizures.

Diazepam is useful premedication for relief of anxiety and tension in patients who are to undergo surgical procedures.

## **CONTRAINDICATIONS**

### **INTRAVENOUS ADMINISTRATION OF DIAZEPAM WITH THE AUTOINJECTOR IS CONTRAINDICATED.**

Diazepam injection is contraindicated in patients with a known hypersensitivity to this drug; acute narrow angle glaucoma; and open angle glaucoma unless patients are receiving appropriate therapy.

## **WARNINGS**

*Diazepam Autoinjector is to be administered only by the intramuscular (I.M.) route.*

### **Risks from Concomitant Use With Opioids**

Concomitant use of benzodiazepines, including diazepam, and opioids may result in profound sedation, respiratory depression, coma, and death. Because of these risks, reserve concomitant prescribing of benzodiazepines and opioids for patients for whom alternative treatment options are inadequate.

Observational studies have demonstrated that concomitant use of opioid analgesics and benzodiazepines increases the risk of drug-related mortality compared to use of opioids alone. If a decision is made to prescribe diazepam concomitantly with opioids, prescribe the lowest effective dosages and minimum durations of concomitant use, and follow patients closely for signs and symptoms of respiratory depression and sedation. Advise both patients and caregivers about the risks of respiratory depression and sedation when diazepam is used with opioids (*see PRECAUTIONS*).

### **Abuse, Misuse, and Addiction**

The use of benzodiazepines, including Diazepam injection, exposes users to the risks of abuse, misuse, and addiction, which can lead to overdose or death. Abuse and misuse of benzodiazepines often (but not always) involve the use of doses greater than the maximum recommended dosage and commonly involve concomitant use of other medications, alcohol, and/or illicit substances, which is associated with an increased frequency of serious adverse outcomes, including respiratory depression, overdose, or death (*see DRUG ABUSE AND DEPENDENCE: Abuse*).

Before prescribing Diazepam injection and throughout treatment, assess each patient's

risk for abuse, misuse, and addiction. Use of Diazepam injection, particularly in patients at elevated risk, necessitates counseling about the risks and proper use of Diazepam injection along with monitoring for signs and symptoms of abuse, misuse, and addiction. Do not exceed the recommended dosing frequency; avoid or minimize concomitant use of CNS depressants and other substances associated with abuse, misuse, and addiction (e.g., opioid analgesics, stimulants); and advise patients on the proper disposal of unused drug. If a substance use disorder is suspected, evaluate the patient and institute (or refer them for) early treatment, as appropriate.

### **Dependence and Withdrawal Reactions After Use of Diazepam injection More Frequently Than Recommended**

For patients using Diazepam injection more frequently than recommended, to reduce the risk of withdrawal reactions, use a gradual taper to discontinue Diazepam injection (a patient-specific plan should be used to taper the dose).

Patients at an increased risk of withdrawal adverse reactions after benzodiazepine discontinuation or rapid dosage reduction include those who take higher dosages, and those who have had longer durations of use.

#### **Acute Withdrawal Reactions**

The continued use of benzodiazepines may lead to clinically significant physical dependence. Although Diazepam injection is indicated only for intermittent use (see INDICATIONS AND USAGE and DOSAGE AND ADMINISTRATION), if used more frequently than recommended, abrupt discontinuation or rapid dosage reduction of Diazepam injection, or administration of flumazenil (a benzodiazepine antagonist) may precipitate acute withdrawal reactions, which can be life-threatening (e.g., seizures) (see DRUG ABUSE AND DEPENDENCE: Dependence).

#### **Protracted Withdrawal Syndrome**

In some cases, benzodiazepine users have developed a protracted withdrawal syndrome with withdrawal symptoms lasting weeks to more than 12 months (see DRUG ABUSE AND DEPENDENCE: Dependence).

### **Respiratory and Central Nervous System Depression**

Extreme care must be used in administering injectable diazepam to the elderly, to very ill patients and to those with limited pulmonary reserve because of the possibility that apnea and/or cardiac arrest may occur. Concomitant use of barbiturates, alcohol, or other central nervous system depressants increases depression with increased risk of apnea. Resuscitative equipment including that necessary to support respiration should be readily available.

When diazepam is used with a narcotic analgesic, the dosage of the narcotic should be reduced by at least one-third and administered in small increments. In some cases the use of a narcotic may not be necessary (*see PRECAUTIONS*).

Diazepam injection should not be administered to patients in shock, coma, or in acute alcoholic intoxication with depression of vital signs. As is true of most CNS-acting drugs, patients receiving diazepam should be cautioned against engaging in hazardous occupations requiring complete mental alertness, such as operating machinery or

driving a motor vehicle.

### **Use in Patients with Petit Mal Status**

Tonic status epilepticus has been precipitated in patients treated with I.V. diazepam for petit mal status or petit mal variant status.

### **Usage in Pregnancy**

An increased risk of congenital malformations associated with the use of minor tranquilizers (diazepam, meprobamate and chlordiazepoxide) during the first trimester of pregnancy has been suggested in several studies. Because use of these drugs is rarely a matter of urgency, their use during this period should almost always be avoided. The possibility that a woman of childbearing potential may be pregnant at the time of institution of therapy should be considered. Patients should be advised that if they become pregnant during therapy or intend to become pregnant they should communicate with their physicians about the desirability of discontinuing the drug.

In humans, measurable amounts of diazepam were found in maternal and cord blood, indicating placental transfer of the drug. Until additional information is available, diazepam injection is not recommended for obstetrical use.

### **Pediatric Use**

Efficacy and safety of parenteral diazepam has not been established in the neonate (30 days or less of age).

Prolonged central nervous system depression has been observed in neonates, apparently due to inability to biotransform diazepam into inactive metabolites.

## **PRECAUTIONS**

### **Drug Interactions**

#### **Effect of Concomitant Use of Benzodiazepines and Opioids**

The concomitant use of benzodiazepines and opioids increases the risk of respiratory depression because of actions at different receptor sites in the CNS that control respiration. Benzodiazepines interact at GABA<sub>A</sub> sites, and opioids interact primarily at mu receptors. When benzodiazepines and opioids are combined, the potential for benzodiazepines to significantly worsen opioid-related respiratory depression exists. Limit dosage and duration of concomitant use of benzodiazepines and opioids, and follow patients closely for respiratory depression and sedation.

#### **Concomitant Use with Other Psychotropic Agents or Anticonvulsant Drugs**

If diazepam is to be combined with other psychotropic agents or anticonvulsant drugs, careful consideration should be given to the pharmacology of the agents to be employed - particularly with known compounds which may potentiate the action of diazepam, such as phenothiazines, narcotics, barbiturates, MAO inhibitors and other antidepressants. In highly anxious patients with evidence of accompanying depression, particularly those who may have suicidal tendencies, protective measures may be necessary. The usual

precautions in treating patients with impaired hepatic function should be observed. Metabolites of diazepam are excreted by the kidney; to avoid their excess accumulation, caution should be exercised in the administration to patients with compromised kidney function.

#### Concomitant Use with Cimetidine

The clearance of diazepam and certain other benzodiazepines can be delayed in association with Tagamet (cimetidine) administration. The clinical significance of this is unclear.

### **Administration Considerations**

#### Long Term Control of Seizures

Although seizures may be brought under control promptly with a single 10 mg intramuscular dose, a significant proportion of patients may experience a return to seizure activity. Consequently, it may become necessary to re-administer the drug. The cumulative maximum dose of diazepam should not exceed 30 mg; the interval between doses should be no less than 10 minutes. Diazepam is not recommended for maintenance, and once seizures are brought under control, consideration should be given to the administration of agents useful in longer term control of seizures.

Since an increase in cough reflex and laryngospasm may occur with peroral endoscopic procedures, the use of a topical anesthetic agent and the availability of necessary countermeasures are recommended.

#### Use During Pregnancy

Until additional information is available, injectable diazepam is not recommended for obstetrical use.

#### Hypotension and Muscular Weakness

Diazepam injection has produced hypotension or muscular weakness in some patients particularly when used with narcotics, barbiturates or alcohol.

### **ADVERSE REACTIONS**

Side effects most commonly reported were drowsiness, fatigue and ataxia; venous thrombosis and phlebitis at the site of injection. Other adverse reactions less frequently reported include: *CNS*: confusion, depression, dysarthria, headache, hypoactivity, slurred speech, syncope, tremor, vertigo. *G.I.*: constipation, nausea. *G.U.*: incontinence, changes in libido, urinary retention. *Cardiovascular*: bradycardia, cardiovascular collapse, hypotension. *EENT*: blurred vision, diplopia, nystagmus. *Skin*: urticaria, skin rash. *Other*: hiccups, changes in salivation, neutropenia, jaundice. Paradoxical reactions such as acute hyperexcited states, anxiety, hallucinations, increased muscle spasticity, insomnia, rage, sleep disturbances and stimulation have been reported; should these occur, use of the drug should be discontinued.

Minor changes in EEG patterns, usually low-voltage fast activity, have been observed in patients during and after diazepam therapy and are of no known significance.

In peroral endoscopic procedures, coughing, depressed respiration, dyspnea, hyperventilation, laryngospasm and pain in throat or chest have been reported.

Because of isolated reports of neutropenia and jaundice, periodic blood counts and liver function tests are advisable during long-term therapy.

## **DRUG ABUSE AND DEPENDENCE**

### **Controlled Substance**

Diazepam is a schedule IV controlled substance.

### **Abuse**

Diazepam injection is a benzodiazepine and a CNS depressant with a potential for abuse and addiction. Abuse is the intentional, non-therapeutic use of a drug, even once, for its desirable psychological or physiological effects. Misuse is the intentional use, for therapeutic purposes, of a drug by an individual in a way other than prescribed by a health care provider or for whom it was not prescribed. Drug addiction is a cluster of behavioral, cognitive, and physiological phenomena that may include a strong desire to take the drug, difficulties in controlling drug use (e.g., continuing drug use despite harmful consequences, giving a higher priority to drug use than other activities and obligations), and possible tolerance or physical dependence. Even taking benzodiazepines as prescribed may put patients at risk for abuse and misuse of their medication. Abuse and misuse of benzodiazepines may lead to addiction.

Abuse and misuse of benzodiazepines often (but not always) involve the use of doses greater than the maximum recommended dosage and commonly involve concomitant use of other medications, alcohol, and/or illicit substances, which is associated with an increased frequency of serious adverse outcomes, including respiratory depression, overdose, or death. Benzodiazepines are often sought by individuals who abuse drugs and other substances, and by individuals with addictive disorders (see WARNINGS: Abuse, Misuse, and Addiction).

The following adverse reactions have occurred with benzodiazepine abuse and/or misuse: abdominal pain, amnesia, anorexia, anxiety, aggression, ataxia, blurred vision, confusion, depression, disinhibition, disorientation, dizziness, euphoria, impaired concentration and memory, indigestion, irritability, muscle pain, slurred speech, tremors, and vertigo.

The following severe adverse reactions have occurred with benzodiazepine abuse and/or misuse: delirium, paranoia, suicidal ideation and behavior, seizures, coma, breathing difficulty, and death. Death is more often associated with polysubstance use (especially benzodiazepines with other CNS depressants such as opioids and alcohol).

### **Dependence**

Physical Dependence After Use of Diazepam Injection More Frequently Than Recommended

Diazepam injection may produce physical dependence if used more frequently than recommended. Physical dependence is a state that develops as a result of physiological

adaptation in response to repeated drug use, manifested by withdrawal signs and symptoms after abrupt discontinuation or a significant dose reduction of a drug. Although Diazepam injection is indicated only for intermittent use (see INDICATIONS AND USAGE and DOSAGE AND ADMINISTRATION), if used more frequently than recommended, abrupt discontinuation or rapid dosage reduction or administration of flumazenil, a benzodiazepine antagonist, may precipitate acute withdrawal reactions, including seizures, which can be life-threatening. Patients at an increased risk of withdrawal adverse reactions after benzodiazepine discontinuation or rapid dosage reduction include those who take higher dosages (i.e., higher and/or more frequent doses) and those who have had longer durations of use (see WARNINGS: Dependence and Withdrawal Reactions).

For patients using Diazepam injection more frequently than recommended, to reduce the risk of withdrawal reactions, use a gradual taper to discontinue Diazepam injection (see WARNINGS: Dependence and Withdrawal Reactions).

### *Acute Withdrawal Signs and Symptoms*

Acute withdrawal signs and symptoms associated with benzodiazepines have included abnormal involuntary movements, anxiety, blurred vision, depersonalization, depression, derealization, dizziness, fatigue, gastrointestinal adverse reactions (e.g., nausea, vomiting, diarrhea, weight loss, decreased appetite), headache, hyperacusis, hypertension, irritability, insomnia, memory impairment, muscle pain and stiffness, panic attacks, photophobia, restlessness, tachycardia, and tremor. More severe acute withdrawal signs and symptoms, including life-threatening reactions, have included catatonia, convulsions, delirium tremens, depression, hallucinations, mania, psychosis, seizures, and suicidality.

### *Protracted Withdrawal Syndrome*

Protracted withdrawal syndrome associated with benzodiazepines is characterized by anxiety, cognitive impairment, depression, insomnia, formication, motor symptoms (e.g., weakness, tremor, muscle twitches), paresthesia, and tinnitus that persists beyond 4 to 6 weeks after initial benzodiazepine withdrawal. Protracted withdrawal symptoms may last weeks to more than 12 months. As a result, there may be difficulty in differentiating withdrawal symptoms from potential re-emergence or continuation of symptoms for which the benzodiazepine was being used.

### Tolerance

Tolerance to Diazepam injection may develop after use more frequently than recommended. Tolerance is a physiological state characterized by a reduced response to a drug after repeated administration (i.e., a higher dose of a drug is required to produce the same effect that was once obtained at a lower dose). Tolerance to the therapeutic effect of benzodiazepines may develop; however, little tolerance develops to the amnestic reactions and other cognitive impairments caused by benzodiazepines.

## **DOSAGE AND ADMINISTRATION (INTRAMUSCULAR ONLY)**

### **Administration of the Diazepam Autoinjector**



1. *Pull off gray safety cap*
2. *Place Black end on mid outer thigh*
3. *Push hard until injector functions*
4. *Withdraw after 10 seconds*

## **Dosage**

**Caution: Because the autoinjector automatically delivers a fixed dose of 10 mg of diazepam, it cannot be used in situations requiring lower total doses or those in which small incremental increases of diazepam are required.**

The usual recommended dose in older children and adults ranges from 10 mg to 20 mg I.M. depending on the indication and its severity. The cumulative total dose and individual maximum dose for intramuscular administration will vary with the specific indication (See dosage for specific indications).

### **Intramuscular**

When instructions are followed properly the Diazepam Autoinjector injects deeply into the muscle.

### **Intravenous Use**

The Diazepam Autoinjector is not designed or intended for intravenous use.

### **Severe Anxiety Disorders and Symptoms of Anxiety**

Usual Adult Dosage: 10 mg, I. M. Repeat in 3 to 4 hours, if necessary.

### **Acute Alcohol Withdrawal**

As an aid in symptomatic relief of acute agitation, tremor, impending or acute delirium tremens and hallucinosis.

Usual Adult Dosage: 10 mg, I. M. initially, then 10 mg in 3 to 4 hours, if necessary.

### **Endoscopic procedures**

Adjunctively, if apprehension, anxiety or acute stress reactions are present prior to endoscopic procedures. Dosage of narcotics should be reduced by at least a third and in some cases may be omitted. See Precautions for peroral procedures.

Usual Adult Dosage: 10 mg I. M., approximately 30 minutes prior to the procedure.

### **Muscle Spasm**

Associated with local pathology, cerebral palsy, athetosis, stiff-man syndrome or tetanus.

Usual Adult Dosage: 10 mg, I. M., initially, then 10 mg in 3 to 4 hours, if necessary. For tetanus, larger doses may be required.

### **Status Epilepticus and Severe Recurrent Convulsive Seizures**

In the convulsing patient, the I.V. route is preferred. However, if conditions preclude intravenous administration, the I.M. route may be used.

Usual Adult Dosage: 10 mg, initially. This injection may be repeated if necessary at 10 to 15 minute intervals up to a maximum dose of 30 mg. If necessary, therapy with diazepam may be repeated in 2 to 4 hours; however, residual active metabolites may persist, and re-administration should be made with this consideration.

Extreme caution must be exercised with individuals with chronic lung disease or unstable cardiovascular status.

#### Preoperative Medication

To relieve anxiety and tension. (if atropine, scopolamine or other premedications are desired, they must be administered in separate syringes).

Usual Adult Dosage: 10 mg, I.M. (preferred route), before surgery.

#### General

Once the acute symptomatology has been properly controlled with diazepam injection, the patient may be placed on oral therapy with diazepam if further treatment is required.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever the solution and container permit.

#### Management of Overdosage

**Manifestations of diazepam overdose include somnolence, confusion, coma and diminished reflexes. Respiration, pulse and blood pressure should be monitored as in all cases of drug overdose, although, in general, these effects have been minimal. General supportive measures should be employed, along with intravenous fluids, and an adequate airway maintained. Hypotension may be combated by the use of norepinephrine or metaraminol. Dialysis is of limited value.**

**Flumazenil, a specific benzodiazepine receptor antagonist, is indicated for the complete or partial reversal of the sedative effects of benzodiazepine and may be used in situations when an overdose with a benzodiazepine is known or suspected. Prior to the administration of flumazenil, necessary measures should be instituted to secure airways, ventilation, and intravenous access. Flumazenil is intended as an adjunct to, not as a substitute for, proper management of benzodiazepine overdose. Patients treated with flumazenil should be monitored for re sedation, respiratory depression, and other residual benzodiazepine effects for an appropriate period after treatment. The prescriber should be aware of a risk of seizure in association with flumazenil treatment, particularly in long term benzodiazepine users and in cyclic antidepressant overdose. The complete flumazenil package insert including CONTRAINDICATIONS, WARNINGS, and PRECAUTIONS should be consulted prior to use.**

#### HOW SUPPLIED

Diazepam injection, 5 mg per mL, is available in 2 mL Disposable Autoinjectors.

## **ANIMAL PHARMACOLOGY**

Oral LD<sub>50</sub> of diazepam is 720 mg/kg in mice and 1240 mg/kg in rats. Intraperitoneal administration of 400 mg/kg to a monkey resulted in death on the sixth day.

## **Reproduction Studies**

A series of rat reproduction studies was performed with diazepam in oral doses of 1, 10, 80 and 100 mg/kg given for periods ranging from 60-228 days prior to mating. At 100 mg/kg there was a decrease in the number of pregnancies and surviving offspring in these rats. These effects may be attributable to prolonged sedative activity, resulting in lack of interest in mating and lessened maternal nursing and care of the young. Neonatal survival of rats at doses lower than 100 mg/kg was within normal limits. Several neonates, both controls and experimentals, in these rat reproduction studies showed skeletal or other defects. Further studies in rats at doses up to and including 80 mg/kg/day did not reveal significant teratological effects on the offspring. Rabbits were maintained on doses of 1, 2, 5 and 8 mg/kg from day 6 through day 18 of gestation. No adverse effects on reproduction and no teratological changes were noted.

## **STORAGE**

Store at 25°C (77°F); Excursions permitted to 15-30°C (59-86°F). [See USP Controlled Room Temperature]. Do not refrigerate.

Office of the Surgeon General  
U.S. Army Medical Research and Development Command (MCMR-RCQ-HR)  
Fort Detrick  
Frederick, Maryland 21702-5012

Manufactured by:  
Revision: 2/2021  
Meridian Medical Technologies, Inc.  
Columbia, MD 21046  
A Pfizer company  
0002147

## **Principal Display Panel - 2 mL Injector Label**

**MERIDIAN**

**MEDICAL TECHNOLOGIES™**

**Columbia, MD 21046**

**A subsidiary of King**

**Pharmaceuticals®, Inc.**

**AUTOINJECTOR  
FOR BUDDY USE**

**Store at 25°C (77°F); Excursions permitted to 15-30°C (59-86°F). [See USP Controlled Room Temperature]. Do not refrigerate.**

**AUTOINJECTOR  
FOR BUDDY USE**

**Rx Only**

- 1. Pull off gray safety cap**
- 2. Place black end on mid outer thigh and**
- 3. Push hard until injector functions**
- 4. Withdraw after 10 seconds**

**MERIDIAN**

**MEDICAL TECHNOLOGIES™**

**3 NDC 11704 60001 5**

**NSN 6505-01-274-0951 C-IV**

**DIAZEPAM INJECTION, USP, 10 mg AUTOMATIC**

**MERIDIAN**  
MEDICAL TECHNOLOGIES™  
Columbia, MD 21046  
A subsidiary of King  
Pharmaceuticals\*, Inc.

# AUTOINJECTOR FOR BUDDY USE



Store at 25° C (77° F); Excursions permitted  
to 15 - 30° C (59 - 86° F). [See USP Controlled  
Room Temperature]. Do not refrigerate.

Rx Only

# AUTOINJECTOR FOR BUDDY USE

0001612

1. Pull off gray safety cap
2. Place black end on mid outer thigh and
3. Push hard until injector functions
4. Withdraw after 10 seconds

**MERIDIAN**  
MEDICAL TECHNOLOGIES™

3 NDC 11704 60001 5



NSN 6505-01-274-0951

# C-IV

## DIAZEPAM INJECTION, USP, 10 mg AUTOMATIC

### DIAZEPAM

diazepam injection

#### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:11704-600
Route of Administration	INTRAMUSCULAR	DEA Schedule	CIV

#### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
diazepam (UNII: Q3JTX2Q7TU) (diazepam - UNII:Q3JTX2Q7TU)	diazepam	10 mg in 2 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>Sodium Benzoate</b> (UNII: OJ245FE5EU)	
<b>Benzoic Acid</b> (UNII: 8SKN0B0MIM)	
<b>Propylene Glycol</b> (UNII: 6DC9Q167V3)	
<b>Alcohol</b> (UNII: 3K9958V90M)	
<b>Benzyl Alcohol</b> (UNII: LKG8494WBH)	
<b>Water</b> (UNII: 059QF0KO0R)	
<b>Nitrogen</b> (UNII: N762921K75)	
<b>Sodium Hydroxide</b> (UNII: 55X04QC32I)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11704-600-01	1 in 1 POUCH	12/05/1990	
1		2 mL in 1 SYRINGE, GLASS; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020124	12/05/1990	

**Labeler** - Meridian Medical Technologies, Inc. (167671341)

## Establishment

Name	Address	ID/FEI	Business Operations
Meridian Medical Technologies, Inc.		038889234	MANUFACTURE(11704-600) , LABEL(11704-600) , PACK(11704-600) , ANALYSIS(11704-600) , STERILIZE(11704-600)

Revised: 11/2021

Meridian Medical Technologies, Inc.