

**DAYTIME NIGHTTIME COLD AND FLU RELIEF MULTI-SYMP TOM-  
acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine  
hcl  
WinCo Foods, LLC**

-----

**Drug Facts**

**Active ingredients in Daytime (in each softgel)**

**Acetaminophen 325 mg**

Dextromethorphan HBr 10 mg

Phenylephrine HCL 5 mg

**Active ingredients in Nighttime (in each softgel)**

**Acetaminophen 325 mg**

Dextromethorphan HBr 15 mg

Doxylamine succinate 6.25 mg

**Purpose for Daytime**

**Pain reliever/fever reducer**

Cough suppressant

Nasal decongestant

**Purpose for Nighttime**

**Pain reliever/fever reducer**

Cough suppressant

Antihistamine

**Uses**

**DAYTIME**

- temporarily relieves common cold and flu symptoms
  - cough due to minor throat and bronchial irritation
  - nasal congestion
  - headache
  - minor aches and pains
  - fever
  - sore throat

## **NIGHTTIME**

- temporarily relieves common cold and flu symptoms
  - cough due to minor throat and bronchial irritation
  - sore throat
  - headache
  - minor aches and pain
  - fever
  - runny nose and sneezing

## **Warnings**

### **DAYTIME and NIGHTTIME**

**Liver warning:** These products contain acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

## **Do not use**

### **DAYTIME and NIGHTTIME**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

## **Ask a doctor before use if you have**

### DAYTIME

- liver disease
- heart disease
- diabetes
- thyroid disease
- high blood pressure
- cough that occurs with too much phlegm (mucus)

- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

#### **NIGHTTIME**

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts such as occur with smoking, asthma, chronic bronchitis or emphysema
- trouble urinating due to an enlarged prostate gland

#### **Ask a doctor or pharmacist before use if you are**

#### **DAYTIME**

- taking the blood thinning drug warfarin

#### **NIGHTTIME**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

#### **When using this product,**

#### **DAYTIME**

#### **do not exceed recommended dosage.**

#### **NIGHTTIME**

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

#### **Stop use and ask a doctor if**

#### **DAYTIME**

- nervousness, dizziness or sleeplessness occur
- pain, cough, and nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

#### **NIGHTTIME**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**DAYTIME and NIGHTTIME**

**Overdose warning:** Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

**DAYTIME**

- **do not take more than directed (see Overdose warning )**
- do not take more than 4 doses in 24 hours
- swallow whole; do not crush, chew, or dissolve
- adults and children 12 years and over: take 2 softgels with water every 4 hours
- children under 12 years : do not use

**Nighttime:**

- **do not take more than directed (see Overdose warning)**
- do not take more than 4 doses in 24 hours
- swallow whole; do not crush, chew, or dissolve
- adults and children 12 years and over: take 2 softgels with water every 6 hours
- children under 12 years: do not use

**Other information**

**DAYTIME and NIGHTTIME**

- store between 15-30°C (59-86°F)
- avoid excessive heat

**Inactive ingredients**

**Daytime** butylated hydroxyanisole, butylated hydroxytoluene, FD&C yellow #6, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan\*, sorbitol, white ink

**Nighttime** D&C yellow #10, FD&C blue #1, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

**Questions or comments?**

Call **1-800-824-1706** Monday-Friday 9AM-4PM MST

## **Principal Display Panel**

†Compare to the active ingredients in VICKS® DAYQUIL® & NYQUIL® Cold & Flu LiquiCaps®

### **MULTI-SYMP TOM COLD & Flu Relief**

NON-DROWSY

#### **Daytime**

ACETAMINOPHEN / PAIN RELIEVER - FEVER REDUCER

DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT

PHENYLEPHRINE HCL / NASAL DECONGESTANT

Relieves: Aches, Fever, Sore Throat, Cough & Nasal Congestion

Non-drowsy, Alcohol-free, Antihistamine-free

softgels\*\*

\*\*Liquid-Filled Capsules

#### **Nighttime**

ACETAMINOPHEN / PAIN RELIEVER - FEVER REDUCER

DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT

DOXYLAMINE SUCCINATE / ANTIHISTAMINE

Relieves: Aches, Fever, Sore Throat, Cough, Sneezing & Runny Nose

Alcohol-free

Softgels

\*\*Liquid-Filled Capsules

**When using Daytime and Nighttime products, carefully read the labeling to ensure correct dosing.**

†This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, DayQuil®, NyQuil®, and LiquiCap® are registered trademarks of The Procter and Gamble Company.

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

DISTRIBUTED BY:

WINCO FOODS, LLC,

BOISE, ID 83704

[www.wincofoods.com](http://www.wincofoods.com)

## Product Label

**Daytime Cold & Flu Softgel**

**Drug Facts (continued)**

**Ask a doctor before use if you have** ■ liver disease ■ heart disease ■ high blood pressure ■ diabetes ■ thyroid disease ■ cough that occurs with too much phlegm (mucus) ■ trouble urinating due to an enlarged prostate gland ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.**

**When using this product, do not exceed recommended dosage.**

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleepiness occur
- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- redness or swelling is present ■ new symptoms occur
- fever gets worse or lasts more than 3 days
- cough comes back or occurs with rash or headache that lasts

**If pregnant or breast-feeding, ask a health professional before use.** These could be signs of a serious condition.

**Keep out of reach of children. Overdose warning:** Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in 24 hours
- swallow whole; do not crush, chew, or dissolve
- adults and children 12 years and over: take 2 softgels with water every 4 hours
- children under 12 years: do not use

**Nighttime Cold & Flu Softgel**

**Drug Facts (continued)**

**Ask a doctor before use if you have** ■ liver disease ■ gout ■ cough that occurs with too much phlegm (mucus) ■ asthma, chronic bronchitis, or emphysema

**Ask a doctor or pharmacist before use if you are** ■ taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers ■ taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers

**When using this product** ■ do not exceed recommended dosage

**marked drowsiness may occur** ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present ■ new symptoms occur
- cough comes back or occurs with rash or headache that lasts

**If pregnant or breast-feeding, ask a health professional before use.** These could be signs of a serious condition.

**Keep out of reach of children. Overdose warning:** Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in 24 hours
- swallow whole; do not crush, chew, or dissolve
- adults and children 12 years and over: take 2 softgels with water every 6 hours
- children under 12 years: do not use

**Daytime Cold & Flu Softgel**

**Drug Facts**

**Purposes**

- Acetaminophen 325 mg.....Pain reliever/reducer
- Dextromethorphan HBr 10 mg.....Cough suppressant
- Phenylephrine HCl 5 mg.....Nasal decongestant

**Uses** ■ temporarily relieves common cold and flu symptoms: ■ cough due to minor throat and bronchial irritation ■ nasal congestion ■ headache ■ minor aches and pains ■ fever ■ sore throat

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Nighttime Cold & Flu Softgel**

**Drug Facts**

**Purposes**

- Acetaminophen 325 mg.....Pain reliever/reducer
- Dextromethorphan HBr 15 mg.....Cough suppressant
- Doxylamine succinate 6.25 mg.....Antihistamine

**Uses** ■ temporarily relieves common cold and flu symptoms: ■ cough due to minor throat and bronchial irritation ■ sore throat ■ headache ■ minor aches and pains ■ fever ■ runny nose & sneezing

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.



\*Compare to the active ingredients in VICKS® DAYQUIL® & NYQUIL® Cold & Flu LiquiCaps®

**NON-DROWSY**  
**Daytime**  
**ACETAMINOPHEN/**  
**PAIN RELIEVER - FEVER REDUCER**  
**DEXTROMETHORPHAN HBr/**  
**COUGH SUPPRESSANT**  
**PHENYLEPHRINE HCl/**  
**NASAL DECONGESTANT**  
 Relieves: Aches, Fever, Sore Throat,  
 Cough & Nasal Congestion  
 Non-drowsy, Alcohol-free, Antihistamine-free



\*\*Liquid-Filled Capsules



PLD-05704 FC006976 W17806

Comments & Questions?  
800-824-1706  
www.wincofoods.com

**100% SATISFACTION GUARANTEE**

The Supermarket  
Low Price Leader®

DISTRIBUTED BY:  
WINCO FOODS, LLC,  
BOISE, ID 83704

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BUSTERS UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® DayQuil®, NyQuil®, and LiquiCaps® are registered trademarks of The Procter and Gamble Company.

<p><b>Daytime Cold &amp; Flu Softgel</b></p> <p><b>Drug Facts (continued)</b></p> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>■ store between 15-30°C (59-86°F)</li> <li>■ avoid excessive heat</li> </ul> <p><b>Inactive ingredients</b> butylated hydroxytoluene, butylated hydroxytoluene, FD&amp;C yellow #6, gelatin, glycerin, mannitol, polyethylene glycol, polyethylene glycol, purified water, sorbitol, white ink</p> <p><b>Questions or comments?</b></p> <p>Call 800-824-1706 Monday - Friday 9am - 4pm MST</p>	<p><b>Nighttime Cold &amp; Flu Softgel</b></p> <p><b>Drug Facts (continued)</b></p> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>■ store between 15-30°C (59-86°F)</li> <li>■ avoid excessive heat</li> </ul> <p><b>Inactive ingredients</b> D&amp;C yellow #10, FD&amp;C blue #1, gelatin, glycerin, mannitol, polyethylene glycol, polyethylene glycol, purified water, sorbitol, white ink</p> <p><b>Questions or comments?</b></p> <p>Call 800-824-1706 Monday - Friday 9am - 4pm MST</p>
---	---

### WINCO FOODS Daytime Nighttime Cold & Flu Relief

## DAYTIME NIGHTTIME COLD AND FLU RELIEF MULTI-SYMPOM

acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl kit

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:67091-336
---------------------	----------------	---------------------------	---------------

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:67091-336-48	1 in 1 KIT; Type 0: Not a Combination Product	06/01/2019	

### Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	16 BLISTER PACK	16



<b>Part 2</b>	32 BLISTER PACK	32
---------------	-----------------	----

## Part 1 of 2

### NIGHT TIME COLD AND FLU RELIEF MULTI SYMPTOM

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate capsule, liquid filled

#### Product Information

<b>Item Code (Source)</b>	NDC:67091-338
<b>Route of Administration</b>	ORAL

#### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9BI9B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

#### Inactive Ingredients

Ingredient Name	Strength
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SORBITAN</b> (UNII: 6O92ICV9RU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>MANNITOL</b> (UNII: 3OWL53L36A)	

#### Product Characteristics

<b>Color</b>	green (Clear)	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	20mm
<b>Flavor</b>		<b>Imprint Code</b>	P30
<b>Contains</b>			

#### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date

1	16 in 1 CARTON		
1	1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/01/2019	

## Part 2 of 2

### DAYTIME COLD AND FLU NON DROWSY

acetaminophen, dextromethorphan hbr, phenylephrine hcl capsule, liquid filled

## Product Information

Item Code (Source)	NDC:67091-337
Route of Administration	ORAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>BUTYLATED HYDROXYANISOLE</b> (UNII: REK4960K2U)	
<b>BUTYLATED HYDROXYTOLUENE</b> (UNII: 1P9D0Z171K)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SORBITAN</b> (UNII: 6O921CV9RU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>MANNITOL</b> (UNII: 3OWL53L36A)	

## Product Characteristics

<b>Color</b>	orange (Clear)	<b>Score</b>	no score
<b>Shape</b>	CAPSULE	<b>Size</b>	20mm
<b>Flavor</b>		<b>Imprint Code</b>	P19
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		32 in 1 CARTON		
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/01/2019	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/01/2019	

**Labeler** - WinCo Foods, LLC (056098817)

Revised: 5/2024

WinCo Foods, LLC