

**ADVIL COLD AND SINUS- ibuprofen and pseudoephedrine
hydrochloride tablet, coated
Haleon US Holdings LLC**

Drug Facts

ACTIVE INGREDIENTS (IN EACH CAPLET)

Ibuprofen 200 mg (NSAID)*

Pseudoephedrine HCl 30 mg

*nonsteroidal anti-inflammatory drug

PURPOSES

Pain reliever/fever reducer

Nasal decongestant

USES

temporarily relieves these symptoms associated with the common cold or flu:

- headache
- fever
- sinus pressure
- nasal congestion
- minor body aches and pains

WARNINGS

Allergy alert:

Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- in children under 12 years of age
- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, thyroid disease, diabetes, have trouble urinating due to an enlarged prostate gland, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other product that contains pseudoephedrine or any other nasal decongestant
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- fever gets worse or lasts more than 3 days
- nasal congestion lasts for more than 7 days
- symptoms continue or get worse
- redness or swelling is present in the painful area
- you get nervous, dizzy, or sleepless
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

DIRECTIONS

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years of age and over:
 - take 1 caplet every 4 to 6 hours while symptoms persist. If symptoms do not respond to 1 caplet, 2 caplets may be used.
 - do not use more than 6 caplets in any 24-hour period unless directed by a doctor
- children under 12 years of age: do not use

OTHER INFORMATION

- store at 20-25°C (68-77°F). Avoid excessive heat above 40°C (104°F).
- read all warnings and directions before use. Keep carton.

INACTIVE INGREDIENTS

acetylated monoglycerides, carnauba wax, colloidal silicon dioxide, corn starch, croscarmellose sodium, methylparaben, microcrystalline cellulose, pharmaceutical glaze, pharmaceutical ink, povidone, pregelatinized starch, propylparaben, sodium benzoate, sodium lauryl sulfate, stearic acid, sucrose, synthetic iron oxides, titanium dioxide

QUESTIONS OR COMMENTS?

Call toll free **1-800-88-ADVIL**

PRINCIPAL DISPLAY PANEL

Advil®

COLD & SINUS Non-Drowsy

Ibuprofen 200 mg ▶ Pain Reliever/Fever Reducer **(NSAID)**

Pseudoephedrine HCl 30 mg ▶ Nasal Decongestant

**Relieves Sinus Pressure,
Nasal Congestion and Fever**

**40
COATED
CAPLETS***


***Oval-Shaped Tablets**

Advil®

COLD & SINUS Non-Drowsy

**Ibuprofen 200 mg > Pain Reliever/Fever Reducer (NSAID)
Pseudoephedrine HCl 30 mg > Nasal Decongestant**

**Relieves Sinus Pressure,
Nasal Congestion and Fever**



**40
COATED
CAPLETS***

**Oval-Shaped Tablets*

FPO 128 Code

FPO 128 Code

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Drug Facts (continued)

Warnings: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include hives, facial swelling, asthma (wheezing), or difficulty breathing. Stop use and seek medical help if an allergic reaction occurs, stop use and seek medical help if an allergic reaction occurs, stop use and seek medical help if an allergic reaction occurs.

Other Facts: (continued)

- **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.
- **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.
- **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Warnings: (continued)

- **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.
- **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.
- **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Warnings: (continued)

- **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.
- **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.
- **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Warnings: (continued)

- **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.
- **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.
- **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Warnings: (continued)

- **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.
- **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.
- **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Advil® COLD & SINUS

Product inside sealed in plastic blister with foil backing. Do Not Use if plastic blister or foil barrier is broken.

FPO

3 0573 0180 214

Advil® COLD & SINUS

© 2016 Pfizer Inc.
For most recent product information,
visit www.advil.com

PAA 08/02/46

ADVIL COLD AND SINUS

ibuprofen and pseudoephedrine hydrochloride tablet, coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0573-0180
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg
PSEUDOEPHEDRINE HYDROCHLORIDE (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)	PSEUDOEPHEDRINE HYDROCHLORIDE	30 mg

Inactive Ingredients	
Ingredient Name	Strength
DIACETYLATED MONOGLYCERIDES (UNII: 5Z17386USF)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SHELLAC (UNII: 46N107B71O)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCROSE (UNII: C151H8M554)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics			
Color	BROWN (butterscotch)	Score	no score
Shape	OVAL (oblong)	Size	14mm
Flavor		Imprint Code	Advil;Cold;Sinus
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	NDC:0573			

1	NDC:0573-0180-10	2 in 1 CARTON	09/19/1989	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:0573-0180-21	4 in 1 CARTON	09/19/1989	
2		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA019771	09/19/1989	

Labeler - Haleon US Holdings LLC (079944263)

Establishment

Name	Address	ID/FEI	Business Operations
PF Consumer Healthcare B.V. (Puerto Rico Operations) LLC		118346012	ANALYSIS(0573-0180) , MANUFACTURE(0573-0180)

Revised: 1/2024

Haleon US Holdings LLC