

**FLU HBP MAXIMUM STRENGTH- acetaminophen, chlorpheniramine maleate, dextromethorphan hbr tablet, film coated**  
**Walgreen Company**

-----  
**Walgreens 44-738**

***Active ingredients (in each caplet)***

Acetaminophen 325 mg  
Chlorpheniramine maleate 2 mg  
Dextromethorphan HBr 10 mg

***Purpose***

Pain reliever/fever reducer  
Antihistamine  
Cough suppressant

***Uses***

- temporarily relieves these common cold and flu symptoms:
  - minor aches and pains
  - sore throat
  - sneezing
  - headache
  - runny nose
  - cough
- temporarily reduces fever

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- blisters
- rash
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a

doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if you have**

- difficulty in urination due to enlargement of the prostate gland
- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema

### **Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

### **When using this product**

- avoid alcoholic beverages
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- excitability may occur, especially in children
- use caution when driving a motor vehicle or operating machinery

### **Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### ***Directions***

- **do not take more than directed**
- adults and children 12 years and over: take 2 caplets every 4 hours while symptoms persist. Do not take more than 10 caplets in 24 hours.
- children under 12 years: ask a doctor

### ***Other information***

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number
- protect from excessive moisture

### ***Inactive ingredients***

corn starch, crospovidone, FD&C red #40 aluminum lake, lecithin, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

### ***Questions or comments?***

**1-800-426-9391**

### **Principal Display Panel**

NDC 0363-7380-08

### ***Walgreens***

### **WALGREENS PHARMACIST RECOMMENDED†**

Compare to the active ingredients  
in Coricidin® HBP Maximum  
Strength Multi-Symptom Flu††

### **Flu HBP**

**ACETAMINOPHEN** / PAIN RELIEVER / FEVER REDUCER  
**CHLORPHENIRAMINE MALEATE** / ANTIHISTAMINE  
**DEXTROMETHORPHAN HBr** / COUGH SUPPRESSANT

### **Maximum Strength Decongestant Free**

- Relieves body aches, pains, headache, runny nose, fever, cough, sore throat & sneezing
- For people with high blood pressure

**24**

CAPLETS

ACTUAL SIZE

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER**

# UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

†Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.

DISTRIBUTED BY: **WALGREEN CO.**  
**200 WILMOT RD., DEERFIELD, IL 60015**  
**100% SATISFACTION GUARANTEED**  
**walgreens.com ©2022 Walgreen Co.**

††This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Coricidin® HBP Maximum Strength Multi-Symptom Flu.

50844 ORG101973808

TAMPER EVIDENT - DO NOT USE IF PACKAGE IS OPENED OR IF BUSTERS  
UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.  
DISTRIBUTED BY: WALGREEN CO.  
200 WILMOT RD., DEERFIELD, IL 60015  
100% SATISFACTION GUARANTEED  
walgreens.com ©2022 Walgreen Co.

PARENTS:  
Learn about child-resistant packaging  
www.bayer.com/childresist  
how2recycle.info

NDC 0363-7380-08



Compare to the active ingredients in Coricidin® HBP Maximum Strength Multi-Symptom Flu\*

## Flu HBP

**ACETAMINOPHEN / PAIN RELIEVER / FEVER REDUCER  
CHLORPHENIRAMINE MALEATE / ANTIHISTAMINE  
DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT**

**Maximum Strength    Decongestant Free**

- Relieves body aches, pains, headache, runny nose, fever, cough, sore throat & sneezing
- For people with high blood pressure

**24**

CAPLETS



ACTUAL SIZE

**Drug Facts (continued)**

**Warnings**  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take  
 ■ more than 4,000 mg of acetaminophen in 24 hours  
 ■ with other drugs containing acetaminophen  
 ■ 3 or more alcoholic drinks every day while using this product  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:  
 ■ blisters    ■ rash    ■ skin reddening  
 If a skin reaction occurs, stop use and seek medical help right away.  
**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**  
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.  
 ■ if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have**  
 ■ difficulty in urination due to enlargement of the prostate gland  
 ■ liver disease    ■ glaucoma  
 ■ cough that occurs with too much phlegm (mucus)  
 ■ a breathing problem or persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema

**Drug Facts (continued)**

Ask a doctor or pharmacist before use if you are  
 ■ taking sedatives or tranquilizers  
 ■ taking the blood thinning drug warfarin

**When using this product**  
 ■ avoid alcoholic beverages  
 ■ marked drowsiness may occur  
 ■ alcohol, sedatives, and tranquilizers may increase drowsiness  
 ■ excitability may occur, especially in children  
 ■ use caution when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**  
 ■ pain or cough gets worse or lasts more than 7 days    ■ new symptoms occur  
 ■ fever gets worse or lasts more than 3 days  
 ■ redness or swelling is present  
 ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.  
**Keep out of reach of children.** In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**  
 ■ do not take more than directed  
 ■ adults and children 12 years and over: take 2 caplets every 4 hours while symptoms persist. Do not take more than 10 caplets in 24 hours.  
 ■ children under 12 years: ask a doctor

**Other information**  
 ■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN

**Drug Facts (continued)**

FDCC red #40 aluminum lake, lecithin, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

**Questions or comments? 1-800-426-9391**

**Drug Facts (continued)**

store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)  
 ■ see end flap for expiration date and lot number  
 ■ protect from excessive moisture

**Inactive ingredients**    com starch, croscarmellose

W3ORG0422-F  
REV0422

This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Coricidin® HBP Maximum Strength Multi-Symptom Flu.

ITEM 168432    W00000-0000-0



3 11917 00948 3

50844 ORG101973808

B-2201-738-08-HR  
ORG101973808

**No Print or Varnish Lot and Exp. Date ONLY**

**Walgreens 44-738**

## FLU HBP MAXIMUM STRENGTH

acetaminophen, chlorpheniramine maleate, dextromethorphan hbr tablet, film coated

---

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0363-7380
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>CHLORPHENIRAMINE MALEATE</b> (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>CROSPVIDONE, UNSPECIFIED</b> (UNII: 2S7830E561)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>LECITHIN, SOYBEAN</b> (UNII: 1DI56QDM62)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	red	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	16mm
<b>Flavor</b>		<b>Imprint Code</b>	44;738
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-7380-08	2 in 1 CARTON	03/19/2020	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:0363-7380-09	2 in 1 CARTON	03/19/2020	01/13/2024
2		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/19/2020	

**Labeler** - Walgreen Company (008965063)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(0363-7380) , pack(0363-7380)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(0363-7380)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(0363-7380)

Revised: 5/2024

Walgreen Company