

**ALLERGY MULTI-SYMPTOM- acetaminophen, chlorpheniramine maleate and phenylephrine hcl tablet, film coated  
DOLGENCORP, LLC**

-----  
**Allergy Multi-Symptom**

***Active ingredients (in each caplet)***

Acetaminophen 325 mg  
Chlorpheniramine maleate 2 mg  
Phenylephrine HCl 5 mg

***Purpose***

Pain reliever  
Antihistamine  
Nasal decongestant

***Uses***

- temporarily relieves these symptoms of hay fever or other upper respiratory allergies:
  - sinus congestion and pressure
  - nasal congestion
  - runny nose and sneezing
  - headache
  - minor aches and pains
- temporarily relieves these additional symptoms of hay fever:
  - itching of the nose or throat
  - itchy, watery eyes
  - helps clear nasal passages
  - helps decongest sinus openings and passages

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- blisters

- rash
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if you have**

- glaucoma
- difficulty in urination due to enlargement of the prostate gland
- high blood pressure
- a breathing problem such as emphysema or chronic bronchitis
- heart disease
- thyroid disease
- diabetes
- liver disease

### **Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

### **When using this product**

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic beverages
- use caution when driving a motor vehicle or operating machinery
- drowsiness may occur

### **Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain or nasal congestion gets worse or lasts more than 7 days
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present

These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use.

## **Keep out of reach of children.**

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### ***Directions***

- **do not take more than directed**
- adults and children 12 years and over
  - take 2 caplets every 4 hours
  - swallow whole - do not crush, chew, or dissolve
  - do not take more than 10 caplets in 24 hours
- children under 12 years: ask a doctor

### ***Other information***

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

### ***Inactive ingredients***

corn starch, crospovidone, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, sucralose, talc, titanium dioxide

### ***Questions or comments?***

**1-888-309-9030**

### ***Principal Display Panel***

**DG™ | health**

#### **Allergy**

#### **Multi-Symptom**

**Acetaminophen, 325 mg**  
Chlorpheniramine Maleate, 2 mg  
Phenylephrine HCl, 5 mg

**Pain Reliever, Antihistamine,**  
Nasal Decongestant

Relieves:

- Headache, sinus pressure
- Nasal congestion, runny nose
- Sneezing & itchy, watery eyes

Pseudoephedrine Free

**24** Caplets

Actual Caplet Size

Cool Blast Flavor

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS  
TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING**

50844 REV0818C45508

DISTRIBUTED BY DOLGENCORP, LLC

100 MISSION RIDGE

GOODLETTSVILLE, TN 37072

**100%**

**Satisfaction**

**Guaranteed!**

(888) 309-9030

**100%**  
Satisfaction  
Guaranteed!  
(888) 309-9030

IF PACKAGE IS  
TORN, BROKEN  
OR TAMPERED

155508

DOLGENCORP, LLC  
GE  
E, TN 37072



# Allergy Multi-Symptom

Acetaminophen, 325 mg  
Chlorpheniramine Maleate, 2 mg  
Phenylephrine HCl, 5 mg  
Pain Reliever, Antihistamine,  
Nasal Decongestant

Relieves:



**Drug Facts**  
KEEP OUTER PACKAGE FOR  
COMPLETE PRODUCT INFORMATION

**Active ingredients**  
(in each caplet)  
Acetaminophen 325 mg.....Pain reliever  
Chlorpheniramine maleate 2 mg.....Antihistamine  
Phenylephrine HCl 5 mg.....Nasal decongestant

**Drug Facts (continued)**

temporarily relieves these additional symptoms of  
hay fever: ■ itching of the nose or throat  
■ itchy, watery eyes  
■ helps clear nasal passages  
■ helps decongest sinus openings and passages

**Warnings**  
Liver warning: This product contains  
acetaminophen. Severe liver damage may occur if  
you take  
■ more than 4,000 mg of acetaminophen in 24 hours  
■ with other drugs containing acetaminophen  
■ 3 or more alcoholic drinks every day while using  
this product  
Allergy alert: Acetaminophen may cause severe skin  
reactions. Symptoms may include:  
■ rash ■ skin redness  
■ blisters ■ hives  
If a skin reaction occurs, stop use and seek medical  
help right away.

Do not use ■ with any other drug containing  
acetaminophen (prescription or nonprescription).  
If you are not sure whether a drug contains  
acetaminophen, ask a doctor or pharmacist.  
■ if you are now taking a prescription monoamine  
oxidase inhibitor (MAOI) (certain drugs for  
depression, psychiatric or emotional conditions, or  
Parkinson's disease), or for 2 weeks after stopping  
the MAOI drug. If you do not know if your  
prescription drug contains an MAOI, ask a doctor  
or pharmacist before taking this product.  
■ if you have ever had an allergic reaction to this  
product or any of its ingredients

Ask a doctor before use if you have  
■ difficulty in urination due to enlargement of the  
prostate gland ■ high blood pressure

**Drug Facts (continued)**

**Uses**  
temporarily relieves these symptoms of hay fever  
or other upper respiratory allergies:  
■ sinus congestion and pressure  
■ runny nose and sneezing  
■ headache

**Drug Facts**

■ a breathing problem such as emphysema or chronic  
bronchitis ■ heart disease ■ thyroid disease  
■ diabetes ■ liver disease ■ glaucoma

Ask a doctor or pharmacist before use if you are  
■ taking sedatives or tranquilizers  
■ taking the blood thinning drug warfarin

When using this product  
■ do not exceed recommended dosage  
■ excitability may occur, especially in children  
■ alcohol, sedatives, and tranquilizers may increase  
drowsiness  
■ avoid alcoholic beverages  
use caution when driving a motor vehicle or  
operating machinery ■ drowsiness may occur

Stop use and ask a doctor if  
■ nervousness, dizziness, or sleeplessness occur  
■ pain or nasal congestion gets worse or lasts more  
than 7 days ■ new symptoms occur  
■ fever gets worse or lasts more than 3 days  
■ redness or swelling is present  
These could be signs of a serious condition.  
If pregnant or breast-feeding, ask a health  
professional before use.  
Keep out of reach of children. In case of accidental  
overdose, get medical help or contact a Poison  
Control Center (1-800-222-1222) right away. Prompt  
medical attention is critical for adults as well as for  
children even if you do not notice any signs or  
symptoms.

**Directions** ■ do not take more than directed  
■ adults and children 12 years and over  
■ take 2 caplets every 4 hours  
■ swallow whole - do not crush, chew, or dissolve  
■ do not take more than 10 caplets in 24 hours  
■ children under 12 years: ask a doctor

**Drug Facts (continued)**  
Questions or comments? 1-888-309-9030

No Print / No Varnish  
Lot no. & Exp. date



8-0315-455C-08AMSR  
REV0818C45508



44-455C

## ALLERGY MULTI-SYMP TOM

acetaminophen, chlorpheniramine maleate and phenylephrine hcl tablet, film coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:55910-455
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>CHLORPHENIRAMINE MALEATE</b> (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>CROSPVIDONE, UNSPECIFIED</b> (UNII: 2S7830E561)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	17mm
<b>Flavor</b>	MENTHOL	<b>Imprint Code</b>	44;455
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55910-455-08	2 in 1 CARTON	06/28/2005	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/28/2005	

**Labeler** - DOLGENCORP, LLC (068331990)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(55910-455)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(55910-455) , pack(55910-455)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(55910-455)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(55910-455)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(55910-455)