

**LEVOCETIRIZINE DIHYDROCHLORIDE- levocetirizine dihydrochloride tablet, coated**

**Rite Aid Corporation**

-----

**Levocetirizine Dihydrochloride Tablets USP, 5 mg**

**Active ingredient (in each tablet)**

Levocetirizine dihydrochloride USP, 5 mg

**Purpose**

Antihistamine

**Uses**

temporarily relieves these symptoms due to hay fever or other respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

**Warnings**

**Do not use**

- if you have kidney disease
- if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing cetirizine

**Ask a doctor before use if you have**

- ever had trouble urinating or emptying your bladder

**When using this product**

- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

**Stop use and ask doctor if**

- you have trouble urinating or emptying your bladder
- an allergic reaction to this product occurs. Seek medical help right away.

**If pregnant or breast-feeding:**

- if breast-feeding: not recommended

- if pregnant: ask a health professional before use

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

adults 65 years of age and older	<ul style="list-style-type: none"> <li>• ask a doctor</li> </ul>
adults and children 12-64 years of age	<ul style="list-style-type: none"> <li>• take 1 tablet (5 mg) once daily in the evening</li> <li>• do not take more than 1 tablet (5 mg) in 24 hours</li> <li>• 1/2 tablet (2.5 mg) once daily in the evening may be appropriate for less severe symptoms</li> </ul>
children 6-11 years of age	<ul style="list-style-type: none"> <li>• take 1/2 tablet (2.5 mg) once daily in the evening</li> <li>• do not take more than 1/2 tablet (2.5 mg) in 24 hours</li> </ul>
children under 6 years of age	<ul style="list-style-type: none"> <li>• do not use</li> </ul>
consumers with kidney disease	<ul style="list-style-type: none"> <li>• do not use</li> </ul>

**Other information**

- store between 20° and 25°C (68° and 77°F)
- (bottles only) safety sealed: do not use if carton was opened or if printed foil inner seal on bottle is torn or missing
- (blister only) safety sealed: do not use if seal is broken or if individual blister unit is open or torn

**Inactive ingredients**

colloidal silicon dioxide, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide

**Questions or comments?**

Call **1-888-375-3784**

**PRINCIPAL DISPLAY PANEL**



ORIGINAL  
PRESCRIPTION  
STRENGTH

**allergy relief**

levocetirizine dihydrochloride  
tablets, 5 mg

antihistamine

24 hour

120 TABLETS  
5 mg EACH

**Active ingredient (in each tablet)**

Levocetirizine dihydrochloride USP, 5 mg.....Antihistamine

**Uses** temporarily relieves these symptoms due to hay fever or other respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat **Warnings Do not use** ■ if you have kidney disease ■ if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing cetirizine **Ask a doctor before use if you have** ■ ever had trouble urinating or emptying your bladder **When using this product** ■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery **Stop use and ask a doctor if** ■ you have trouble urinating or emptying your bladder ■ an allergic reaction to this product occurs. Seek medical help right away. **If pregnant or breast-feeding:** ■ if breast-feeding: not recommended ■ if pregnant: ask a health professional before use

**Purpose**

DISTRIBUTED BY: RITE AID  
30 HUNTER LANE  
CAMP HILL, PA 17011

MADE IN INDIA

LOT / EXP

150072302

Peel Here



**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). **Directions** **adults 65 years of age and older** ■ ask a doctor **adults and children 12-64 years of age** ■ take 1 tablet (5 mg) once daily in the evening ■ do not take more than 1 tablet (5 mg) in 24 hours ■ 1/2 tablet (2.5 mg) once daily in the evening may be appropriate for less severe symptoms **children 6-11 years of age** ■ take 1/2 tablet (2.5 mg) once daily in the evening ■ do not take more than 1/2 tablet (2.5 mg) in 24 hours **children under 6 years of age** ■ do not use **consumers with kidney disease** ■ do not use **Other information** ■ store between 20° and 25°C (68° and 77°F) ■ safety sealed: do not use if carton was opened or if printed foil inner seal on bottle is torn or missing **Inactive ingredients** colloidal silicon dioxide, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide **Questions or comments?** call 1-888-375-3784



## LEVOCETIRIZINE DIHYDROCHLORIDE

levocetirizine dihydrochloride tablet, coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:11822-5252(NDC:43598-735)
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>levocetirizine dihydrochloride</b> (UNII: SOD6A38AGA) (levocetirizine - UNII:6U5EA9RT2O)	levocetirizine dihydrochloride	5 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XB4)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6130)	

<b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>Polyethylene Glycol, Unspecified</b> (UNII: 3WJQ0SDW1A)	

Product Characteristics			
<b>Color</b>	white	<b>Score</b>	2 pieces
<b>Shape</b>	OVAL	<b>Size</b>	9mm
<b>Flavor</b>		<b>Imprint Code</b>	L
<b>Contains</b>			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-5252-7	1 in 1 CARTON	02/15/2020	
1		35 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:11822-5252-1	1 in 1 CARTON	02/15/2020	
2		120 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA210375	02/15/2020	

**Labeler** - Rite Aid Corporation (014578892)

Establishment			
Name	Address	ID/FEI	Business Operations
Dr.Reddy's Laboratories Limited-FTO 3		918608162	analysis(11822-5252) , manufacture(11822-5252)

Establishment			
Name	Address	ID/FEI	Business Operations
Reed-Lane, Inc.		001819879	repack(11822-5252)