

SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 430 CEDAR- octinoxate and titanium dioxide emulsion
 SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 440 AMBER- octinoxate and titanium dioxide emulsion
 SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 450 COPPER- octinoxate and titanium dioxide emulsion
 SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 460 TOPAZ- octinoxate and titanium dioxide emulsion
 SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 510 SUEDE- octinoxate and titanium dioxide emulsion
 SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 520 ROSEWOOD- octinoxate and titanium dioxide emulsion
 SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 530 HENNA- octinoxate and titanium dioxide emulsion
 SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 540 MAHOGANY- octinoxate and titanium dioxide emulsion
 SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 550 JASPER- octinoxate and titanium dioxide emulsion
 SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 560 OBSIDIAN- octinoxate and titanium dioxide emulsion
 SHISEIDO AMERICAS CORPORATION

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION

Drug Facts

<i>Active ingredients</i>	<i>Purpose</i>
Octinoxate 4.9%	Sunscreen
Titanium Dioxide 1%	Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see ***Directions***), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs.

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: ask a doctor
- **Sun Protection Measures**. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
 - limit time in the sun, especially from 10 a.m. – 2 p.m.
 - wear long-sleeve shirts, pants, hats, and sunglasses

Inactive Ingredients

DIMETHICONE • WATER • GLYCERIN •
 TRIFLUOROPROPYLDIMETHYL/TRIMETHYLSILOXYSILICATE • PHENYL TRIMETHICONE •
 PEG-10 DIMETHICONE • BUTYLENE GLYCOL • DIMETHICONE CROSSPOLYMER • BIS-
 BUTYLDIMETHICONE POLYGLYCERYL-3 • SORBITAN SESQUIISOSTEARATE • ISOPROPYL
 MYRISTATE • DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER • NACRE POWDER •
 MAGNESIUM ALUMINOMETASILICATE • POLYQUATERNIUM-51 • THYMUS SERPYLLUM
 EXTRACT • SODIUM ACETYLATED HYALURONATE • CYCLOPENTASILOXANE •
 DISTEARDIMONIUM HECTORITE • POLYSILICONE-2 • POLYMETHYLSILSESQUIOXANE •
 TRIMETHYLSILOXYSILYL CARBAMOYL PULLULAN • ALUMINUM HYDROXIDE • STEARIC
 ACID • LITHIUM MAGNESIUM SODIUM SILICATE • PEG/PPG-19/19 DIMETHICONE • CITRIC
 ACID • SODIUM CITRATE • BHT • SILICA • TOCOPHEROL • PHENOXYETHANOL • IRON
 OXIDES • TITANIUM DIOXIDE • MICA •

Other information

Protect this product in this container from excessive heat and direct sun.

Questions or comments?

Call toll free 1-800-906-7503

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 430 Cedar

SHISEIDO
 GINZA TOKYO

SYNCHRO SKIN
 SELF-REFRESHING

Foundation
 Oil-Free

BROAD SPECTRUM
 SPF 30

Sunscreen

30mL 1 FL. OZ.

SELF-REFRESHING
Foundation



SHISEIDO
GINZA TOKYO

東京
銀座

A SHISEIDO first in foundation that refreshes non-stop. ActiveForce technology synchronizes with skin and helps to resist heat, oil, humidity and motion for a fresh, natural finish that lasts all day, 24-hour wear. Flawless. Weightless comfort. Medium, buildable coverage.

DERMATOLOGIST-TESTED.
NON-COMEDOGENIC.

Drug Facts

Active ingredients

Purpose

OCTINOXATE 4.9% -----
Sunscreen
TITANIUM DIOXIDE 1% -----
Sunscreen

Uses

- helps prevent sunburn
- If used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

SHISEIDO AMERICAS CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.
www.shiseido.com



Drug Facts
(continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor

Sun Protection Measures.

Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m. - 2 p.m.
- wear long-sleeve shirts, pants, hats, and sunglasses

SYNCHRO SKIN
SELF-REFRESHING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL 1 FL. OZ.

<http://s1872.com/16092AA>
See our website for more information.

Drug Facts
(continued)

Inactive ingredients


DIMETHICONE - WATER - GLYCERIN - TRIFLUOROPROPYLDIMETHYL/TRIMETHYLSILOXYMETHYL/TRIMETHYLSILOXYSILICATE - PHENYL TRIMETHICONE - PEG-10 DIMETHICONE - BUTYLENE GLYCOL - DIMETHICONE CROSSPOLYMER - BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 - SORBITAN SESQUISOSEARATE - ISOPROPYL MYRISTATE - DIMETHICONE/METHYL DIMETHICONE CROSSPOLYMER - NACRE POWDER - MAGNESIUM ALUMINOMETASILICATE - POLYQUATERNIUM-51 - THYMUS SERPYLLUM EXTRACT - SODIUM ACETYLATED HYALURONATE - CYCLOPENTASILOXANE - DISTEARDIMONIUM HECTORITE - POLYSILOXANE-2 - POLYMETHYLSILSESQUOXANE - TRIMETHYLSILOXYMETHYL CARBAMOYL PULLULAN - ALUMINUM HYDROXIDE - STEARIC ACID - LITHIUM MAGNESIUM SODIUM SILICATE - PEG/PPG-19/19 DIMETHICONE - CITRIC ACID - SODIUM CITRATE - BHT - SILICA - TOCOPHEROL - PHENOXYETHANOL - IRON OXIDES - TITANIUM DIOXIDE - MICA

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503



PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 440 Amber

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
SELF-REFRESHING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL 1 FL. OZ.

SELF-REFRESHING
Foundation



SHISEIDO
GINZA TOKYO

東京
銀座

A SHISEIDO first in foundation that refreshes non-stop. ActiveForce technology synchronizes with skin and helps to resist heat, oil, humidity and motion for a fresh, natural finish that lasts all day, 24-hour wear. Flawless. Weightless comfort. Medium, buildable coverage.

DERMATOLOGIST-TESTED.
NON-COMEDOGENIC.

Drug Facts

Active ingredients

Purpose
OCTINOXATE 4.9% -----
Sunscreen
TITANIUM DIOXIDE 1% -----
Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove. ▶

SHISEIDO AMERICAS CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.
www.shiseido.com



Drug Facts
(continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- **Sun Protection Measures.** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
 - limit time in the sun, especially from 10 a.m. - 2 p.m.
 - wear long-sleeve shirts, pants, hats, and sunglasses ▶

<http://s1872.com/16092AA>
See our website for more information.

SYNCHRO SKIN
SELF-REFRESHING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL 1 FL. OZ.

Drug Facts
(continued)

Inactive ingredients


DIMETHICONE - WATER - GLYCERIN - TRIFLUOROPROPYLDIMETHYLTRIMETHYLSILOXYSULFATE - PHENYL TRIMETHICONE - PEG-10 DIMETHICONE - BUTYLENE GLYCOL - DIMETHICONE CROSSPOLYMER - BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 - SORBITAN SESQUISOSEARATE - ISOPROPYL MYRISTATE - DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER - NACRE POWDER - MAGNESIUM ALUMINOMETASILICATE - POLYQUATERNIUM-61 - THYMUS SERPYLLUM EXTRACT - SODIUM ACETYLATED HYALURONATE - CYCLOPENTASILOXANE - DISTEARDIMONIUM HECTORITE - POLYSILOXANE-2 - POLYMETHYLSILOXANOL - TRIMETHYLSILOXYSILYL CARBAMOYL PULLULAN - ALUMINIUM HYDROXIDE - STEARIC ACID - LITHIUM MAGNESIUM SODIUM SILICATE - PEG/PPG-18/18 DIMETHICONE - CITRIC ACID - SODIUM CITRATE - BHT - SILICA - TOCOPHEROL - PHENOXYETHANOL - IRON OXIDES - TITANIUM DIOXIDE - MICA

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503



PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 450 Copper

SHISEIDO
GINZA TOKYO
SYNCHRO SKIN
SELF-REFRESHING
Foundation
Oil-Free
BROAD SPECTRUM
SPF 30
Sunscreen
30mL 1 FL. OZ.

SELF-REFRESHING
Foundation



SHISEIDO
GINZA TOKYO

東京銀座

A SHISEIDO first in foundation that refreshes non-stop. ActiveForce technology synchronizes with skin and helps to resist heat, oil, humidity and motion for a fresh, natural finish that lasts all day, 24-hour wear. Flawless. Weightless comfort. Medium, buildable coverage.

DERMATOLOGIST-TESTED.
NON-COMEDOGENIC.

Drug Facts

Active ingredients

Purpose
OCTINOXATE 4.9% -----
Sunscreen
TITANIUM DIOXIDE 1% -----
Sunscreen

Uses

■ helps prevent sunburn
■ If used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

SHISEIDO AMERICAS CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.

www.shiseido.com



Drug Facts
(continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor

■ **Sun Protection Measures.**

Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m. - 2 p.m.
- wear long-sleeve shirts, pants, hats, and sunglasses

<http://s1872.com/16092AA>
See our website for more information.

Drug Facts
(continued)

Inactive ingredients

DIMETHICONE - WATER - GLYCERIN - TRIFLUOROPROPYLDIMETHYL/ TRIMETHYLSILOXYSULFATE - PHENYL TRIMETHICONE - PEG-10 DIMETHICONE - BUTYLENE GLYCOL - DIMETHICONE CROSSPOLYMER - BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 - SORBITAN SESQUISOOSTEARATE - ISOPROPYL MYRISTATE - DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER - NACRE POWDER - MAGNESIUM ALUMINOMETASILICATE - POLYQUATERNIUM-51 - THYMUS SERPYLLUM EXTRACT - SODIUM ACETYLATED HYALURONATE - CYCLOPENTASILOXANE - DISTEARDIMONIUM HECTORITE - POLYSILICONE-2 - POLYMETHYLSILSESQUOXANE - TRIMETHYLSILOXYSILYL CARBAMOYL PULLULAN - ALUMINUM HYDROXIDE - STEARIC ACID - LITHIUM MAGNESIUM SODIUM SILICATE - PEG/PPG-19/19 DIMETHICONE - CITRIC ACID - SODIUM CITRATE - BHT - SILICA - TOCOPHEROL - PHENOXYETHANOL - IRON OXIDES - TITANIUM DIOXIDE - MICA

Other Information

■ protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503


SYNCHRO SKIN
SELF-REFRESHING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL 1 FL. OZ.



PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 460 Topaz

SHISEIDO
GINZA TOKYO
SYNCHRO SKIN
SELF-REFRESHING
Foundation
Oil-Free
BROAD SPECTRUM
SPF 30
Sunscreen
30mL 1 FL. OZ.

SELF-REFRESHING
Foundation



SHISEIDO
GINZA TOKYO

東京
銀座

A SHISEIDO first in foundation that refreshes non-stop. ActiveForce technology synchronizes with skin and helps to resist heat, oil, humidity and motion for a fresh, natural finish that lasts all day, 24-hour wear. Flawless. Weightless comfort. Medium, buildable coverage.

DERMATOLOGIST-TESTED.
NON-COMEDOGENIC.

Drug Facts

Active ingredients

Purpose

OCTINOXATE 4.9% -----
Sunscreen
TITANIUM DIOXIDE 1% -----
Sunscreen

Uses

- helps prevent sunburn
- If used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

SHISEIDO AMERICAS CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.
www.shiseido.com



Drug Facts (continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor

Sun Protection Measures.

Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m. - 2 p.m.
- wear long-sleeve shirts, pants, hats, and sunglasses

<http://s1872.com/16092AA>
See our website for more information.

Drug Facts (continued)

Inactive ingredients

DIMETHICONE - WATER - GLYCERIN - TRIFLUOROPROPYLDIMETHYL/ TRIMETHYLSILOXYSILICATE - PHENYL TRIMETHICONE - PEG-10 DIMETHICONE - BUTYLENE GLYCOL - DIMETHICONE CROSSPOLYMER - BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 - SORBITAN SESQUISOESTEARATE - ISOPROPYL MYRISTATE - DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER - NACRE POWDER - MAGNESIUM ALUMINOMETASILICATE - POLYQUATERNIUM-51 - THYMUS SERPYLLUM EXTRACT - SODIUM ACETYLATED HYALURONATE - CYCLOPENTASILOXANE - DISTEARDIMONIUM HECTORITE - POLYSILICONE-2 - POLYMETHYLSILSESQUOXANE - TRIMETHYLSILOXYSILYL CARBAMOYL PULLULAN - ALUMINUM HYDROXIDE - STEARIC ACID - LITHIUM MAGNESIUM SODIUM SILICATE - PEG/PPG-19/19 DIMETHICONE - CITRIC ACID - SODIUM CITRATE - BHT - SILICA - TOCOPHEROL - PHENOXYETHANOL - IRON OXIDES - TITANIUM DIOXIDE - MICA -

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-905-7503



SYNCHRO SKIN
SELF-REFRESHING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL 1 FL. OZ.



PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 510 Suede

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
SELF-REFRESHING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL 1 FL. OZ.

SELF-REFRESHING
Foundation



SHISEIDO
GINZA TOKYO

東京銀座

A SHISEIDO first in foundation that refreshes non-stop. ActiveForce technology synchronizes with skin and helps to resist heat, oil, humidity and motion for a fresh, natural finish that lasts all day, 24-hour wear. Flawless. Weightless comfort. Medium, buildable coverage.

DERMATOLOGIST-TESTED.
NON-COMEDOGENIC.

Drug Facts

Active ingredients

Purpose

OCTINOXATE 4.9% -----
Sunscreen
TITANIUM DIOXIDE 1% -----
Sunscreen

Uses

- helps prevent sunburn
- If used as directed with other sun protection measures (see Directions), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

SHISEIDO AMERICAS CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.
www.shiseido.com



Drug Facts
(continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor

■ Sun Protection Measures.

Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m. - 2 p.m.
- wear long-sleeve shirts, pants, hats, and sunglasses

SYNCHRO SKIN
SELF-REFRESHING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL 1 FL. OZ.

<http://s1872.com/16092AA>
See our website for more information.

Drug Facts
(continued)

Inactive ingredients


DIMETHICONE - WATER - GLYCERIN - TRIFLUOROPROPYLDIMETHYL/TRIMETHYLSILOXYSILICATE - PHENYL TRIMETHICONE - PEG-10 DIMETHICONE - BUTYLENE GLYCOL - DIMETHICONE CROSSPOLYMER - BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 - SORBITAN SESQUISOOSTEARATE - ISOPROPYL MYRISTATE - DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER - NACRE POWDER - MAGNESIUM ALUMINOMETASILICATE - POLYQUATERNIUM-51 - THYMUS SERPYLLUM EXTRACT - SODIUM ACETYLATED HYALURONATE - CYCLOPENTASILOXANE - DISTEARDIMONIUM HECTORITE - POLYSILICONE-2 - POLYMETHYLSILSESQUOXANE - TRIMETHYLSILOXYSILYL CARBAMOYL PULLULAN - ALUMINUM HYDROXIDE - STEARIC ACID - LITHIUM MAGNESIUM SODIUM SILICATE - PEG/PPG-19/19 DIMETHICONE - CITRIC ACID - SODIUM CITRATE - BHT - SILICA - TOCOPHEROL - PHENOXYETHANOL - IRON OXIDES - TITANIUM DIOXIDE - MICA

Other Information

■ protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503



PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 520 Rosewood

SHISEIDO
GINZA TOKYO
SYNCHRO SKIN
SELF-REFRESHING
Foundation
Oil-Free
BROAD SPECTRUM
SPF 30
Sunscreen
30mL 1 FL. OZ.

SELF-REFRESHING
Foundation



SHISEIDO
GINZA TOKYO

東京銀座

A SHISEIDO first in foundation that refreshes non-stop. ActiveForce technology synchronizes with skin and helps to resist heat, oil, humidity and motion for a fresh, natural finish that lasts all day, 24-hour wear. Flawless. Weightless comfort. Medium, buildable coverage.

DERMATOLOGIST-TESTED.
NON-COMEDOGENIC.

Drug Facts

Active ingredients

Purpose
OCTINOXATE 4.9% -----
Sunscreen
TITANIUM DIOXIDE 1% -----
Sunscreen

Uses

- helps prevent sunburn
- If used as directed with other sun protection measures (see Directions), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

SHISEIDO AMERICAS CORPORATION DIST. NEW YORK, NY 10017 MADE IN U.S.A.

www.shiseido.com



Drug Facts (continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor

Sun Protection Measures.

Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m. - 2 p.m.
- wear long-sleeve shirts, pants, hats, and sunglasses

**SYNCHRO SKIN
SELF-REFRESHING**

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL 1 FL. OZ.

Drug Facts (continued)

Inactive ingredients

DIMETHICONE - WATER - GLYCERIN - TRIFLUOROPROPYLDIMETHYL/ TRIMETHYLSILOXYSILICATE - PHENYL TRIMETHICONE - PEG-10 DIMETHICONE - BUTYLENE GLYCOL - DIMETHICONE CROSSPOLYMER - BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 - SORBITAN SESQUISOESTEARATE - ISOPROPYL MYRISTATE - DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER - NACRE POWDER - MAGNESIUM ALUMINOMETASILICATE - POLYQUATERNIUM-51 - THYMUS SERPYLLUM EXTRACT - SODIUM ACETYLATED HYALURONATE - CYCLOPENTASILOXANE - DISTEARDIMONIUM HECTORITE - POLYSILICONE-2 - POLYMETHYLSILSESQUOXANE - TRIMETHYLSILOXYSILYL CARBAMOYL PULLULAN - ALUMINUM HYDROXIDE - STEARIC ACID - LITHIUM MAGNESIUM SODIUM SILICATE - PEG/PPG-19/19 DIMETHICONE - CITRIC ACID - SODIUM CITRATE - BHT - SILICA - TOCOPHEROL - PHENOXYETHANOL - IRON OXIDES - TITANIUM DIOXIDE - MICA


Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

<http://s1872.com/16092AA>
See our website for more information.



PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 530 Henna

SHISEIDO
GINZA TOKYO
SYNCHRO SKIN
SELF-REFRESHING
Foundation
Oil-Free
BROAD SPECTRUM
SPF 30
Sunscreen
30mL 1 FL. OZ.

SELF-REFRESHING
Foundation



SHISEIDO
GINZA TOKYO

東京銀座

A SHISEIDO first in foundation that refreshes non-stop. ActiveForce technology synchronizes with skin and helps to resist heat, oil, humidity and motion for a fresh, natural finish that lasts all day, 24-hour wear. Flawless. Weightless comfort. Medium, buildable coverage.

DERMATOLOGIST-TESTED.
NON-COMEDOGENIC.

Drug Facts

Active ingredients

Purpose

OCTINOXATE 4.9% -----
Sunscreen
TITANIUM DIOXIDE 1% -----
Sunscreen

Uses

- helps prevent sunburn
- If used as directed with other sun protection measures (see Directions), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

**Drug Facts
(continued)**

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor

■ Sun Protection Measures.

Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m. - 2 p.m.
- wear long-sleeve shirts, pants, hats, and sunglasses

**Drug Facts
(continued)**

Inactive ingredients

DIMETHICONE - WATER - GLYCERIN - TRIFLUOROPROPYLDIMETHYL/ TRIMETHYLSILOXY/SILICATE - PHENYL TRIMETHICONE - PEG-10 DIMETHICONE - BUTYLENE GLYCOL - DIMETHICONE CROSSPOLYMER - BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 - SORBITAN SESQUISOOSTEARATE - ISOPROPYL MYRISTATE - DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER - NACRE POWDER - MAGNESIUM ALUMINOMETASILICATE - POLYQUATERNIUM-51 - THYMUS SERPYLLUM EXTRACT - SODIUM ACETYLATED HYALURONATE - CYCLOPENTASILOXANE - DISTEARDIMONIUM HECTORITE - POLYSILICONE-2 - POLYMETHYLSILSESQUOXANE - TRIMETHYLSILOXY/SILYL CARBAMOYL PULLULAN - ALUMINUM HYDROXIDE - STEARIC ACID - LITHIUM MAGNESIUM SODIUM SILICATE - PEG/PPG-19/19 DIMETHICONE - CITRIC ACID - SODIUM CITRATE - BHT - SILICA - TOCOPHEROL - PHENOXYETHANOL - IRON OXIDES - TITANIUM DIOXIDE - MICA

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

SYNCHRO SKIN
SELF-REFRESHING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL 1 FL. OZ.

SHISEIDO AMERICAS
CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.
www.shiseido.com



<http://s1872.com/16092AA>
See our website for more information.



PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 540 Mahogany

SHISEIDO
GINZA TOKYO
SYNCHRO SKIN
SELF-REFRESHING
Foundation
Oil-Free
BROAD SPECTRUM
SPF 30
Sunscreen
30mL 1 FL. OZ.

SELF-REFRESHING
Foundation



SHISEIDO
GINZA TOKYO

東京銀座

A SHISEIDO first in foundation that refreshes non-stop. ActiveForce technology synchronizes with skin and helps to resist heat, oil, humidity and motion for a fresh, natural finish that lasts all day, 24-hour wear. Flawless. Weightless comfort. Medium, buildable coverage.

DERMATOLOGIST-TESTED.
NON-COMEDOGENIC.

Drug Facts

Active ingredients

Purpose
OCTINOXATE 4.9% -----
Sunscreen
TITANIUM DIOXIDE 1% -----
Sunscreen

Uses

- helps prevent sunburn
- If used as directed with other sun protection measures (see Directions), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

SHISEIDO AMERICAS CORPORATION DIST. NEW YORK, NY 10017 MADE IN U.S.A.

www.shiseido.com



Drug Facts (continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor

Sun Protection Measures.

Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m. - 2 p.m.
- wear long-sleeve shirts, pants, hats, and sunglasses

**SYNCHRO SKIN
SELF-REFRESHING**

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL 1 FL. OZ.

Drug Facts (continued)

Inactive ingredients

DIMETHICONE - WATER - GLYCERIN - TRIFLUOROPROPYLDIMETHYL/ TRIMETHYLSILOXYSILICATE - PHENYL TRIMETHICONE - PEG-10 DIMETHICONE - BUTYLENE GLYCOL - DIMETHICONE CROSSPOLYMER - BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 - SORBITAN SESQUISOESTEARATE - ISOPROPYL MYRISTATE - DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER - NACRE POWDER - MAGNESIUM ALUMINOMETASILICATE - POLYQUATERNIUM-51 - THYMUS SERPYLLUM EXTRACT - SODIUM ACETYLATED HYALURONATE - CYCLOPENTASILOXANE - DISTEARDIMONIUM HECTORITE - POLYSILICONE-2 - POLYMETHYLSILSESQUOXANE - TRIMETHYLSILOXYSILYL CARBAMOYL PULLULAN - ALUMINUM HYDROXIDE - STEARIC ACID - LITHIUM MAGNESIUM SODIUM SILICATE - PEG/PPG-19/19 DIMETHICONE - CITRIC ACID - SODIUM CITRATE - BHT - SILICA - TOCOPHEROL - PHENOXYETHANOL - IRON OXIDES - TITANIUM DIOXIDE - MICA


Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

<http://s1872.com/16092AA>
See our website for more information.



PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 550 Jasper

SHISEIDO
GINZA TOKYO
SYNCHRO SKIN
SELF-REFRESHING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL 1 FL. OZ.

SELF-REFRESHING
Foundation



SHISEIDO
GINZA TOKYO

東京銀座

A SHISEIDO first in foundation that refreshes non-stop. ActiveForce technology synchronizes with skin and helps to resist heat, oil, humidity and motion for a fresh, natural finish that lasts all day, 24-hour wear. Flawless. Weightless comfort. Medium, buildable coverage.

DERMATOLOGIST-TESTED.
NON-COMEDOGENIC.

Drug Facts

Active ingredients

Purpose

OCTINOXATE 4.9% -----
Sunscreen
TITANIUM DIOXIDE 1% -----
Sunscreen

Uses

- helps prevent sunburn
- If used as directed with other sun protection measures (see Directions), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

**Drug Facts
(continued)**

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor

■ Sun Protection Measures.

Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m. - 2 p.m.
- wear long-sleeve shirts, pants, hats, and sunglasses

**Drug Facts
(continued)**

Inactive ingredients

DIMETHICONE - WATER - GLYCERIN - TRIFLUOROPROPYLDIMETHYL/ TRIMETHYLSILOXYSILICATE - PHENYL TRIMETHICONE - PEG-10 DIMETHICONE - BUTYLENE GLYCOL - DIMETHICONE CROSSPOLYMER - BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 - SORBITAN SESQUISOOSTEARATE - ISOPROPYL MYRISTATE - DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER - NACRE POWDER - MAGNESIUM ALUMINOMETASILICATE - POLYQUATERNIUM-51 - THYMUS SERPYLLUM EXTRACT - SODIUM ACETYLATED HYALURONATE - CYCLOPENTASILOXANE - DISTEARDIMONIUM HECTORITE - POLYSILICONE-2 - POLYMETHYLSILSESQUOXANE - TRIMETHYLSILOXYSILYL CARBAMOYL PULLULAN - ALUMINUM HYDROXIDE - STEARIC ACID - LITHIUM MAGNESIUM SODIUM SILICATE - PEG/PPG-19/19 DIMETHICONE - CITRIC ACID - SODIUM CITRATE - BHT - SILICA - TOCOPHEROL - PHENOXYETHANOL - IRON OXIDES - TITANIUM DIOXIDE - MICA

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

SYNCHRO SKIN
SELF-REFRESHING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30


Sunscreen

30mL 1 FL. OZ.

SHISEIDO AMERICAS CORPORATION DIST. NEW YORK, NY 10017 MADE IN U.S.A. www.shiseido.com



<http://s1872.com/16092AA>
See our website for more information.



PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 560 Obsidian

SHISEIDO
GINZA TOKYO
SYNCHRO SKIN
SELF-REFRESHING
Foundation
Oil-Free
BROAD SPECTRUM
SPF 30
Sunscreen
30mL 1 FL. OZ.

SELF-REFRESHING
Foundation



SHISEIDO
GINZA TOKYO

東京銀座

A SHISEIDO first in foundation that refreshes non-stop. ActiveForce technology synchronizes with skin and helps to resist heat, oil, humidity and motion for a fresh, natural finish that lasts all day, 24-hour wear. Flawless. Weightless comfort. Medium, buildable coverage.

DERMATOLOGIST-TESTED.
NON-COMEDOGENIC.

Drug Facts

Active ingredients

Purpose

OCTINOXATE 4.9% -----
Sunscreen
TITANIUM DIOXIDE 1% -----
Sunscreen

Uses

- helps prevent sunburn
- If used as directed with other sun protection measures (see Directions), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

SHISEIDO AMERICAS CORPORATION DIST. NEW YORK, NY 10017 MADE IN U.S.A.

www.shiseido.com



Drug Facts (continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor

■ Sun Protection Measures.

Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m. - 2 p.m.
- wear long-sleeve shirts, pants, hats, and sunglasses

<http://s1872.com/16092AA>
See our website for more information.

Drug Facts (continued)

Inactive ingredients

DIMETHICONE - WATER - GLYCERIN - TRIFLUOROPROPYLDIMETHYL/ TRIMETHYLSILOXYSILICATE - PHENYL TRIMETHICONE - PEG-10 DIMETHICONE - BUTYLENE GLYCOL - DIMETHICONE CROSSPOLYMER - BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 - SORBITAN SESQUISOESTEARATE - ISOPROPYL MYRISTATE - DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER - NACRE POWDER - MAGNESIUM ALUMINOMETASILICATE - POLYQUATERNIUM-51 - THYMUS SERPYLLUM EXTRACT - SODIUM ACETYLATED HYALURONATE - CYCLOPENTASILOXANE - DISTEARDIMONIUM HECTORITE - POLYSILICONE-2 - POLYMETHYLSILSESQUOXANE - TRIMETHYLSILOXYSILYL CARBAMOYL PULLULAN - ALUMINUM HYDROXIDE - STEARIC ACID - LITHIUM MAGNESIUM SODIUM SILICATE - PEG/PPG-19/19 DIMETHICONE - CITRIC ACID - SODIUM CITRATE - BHT - SILICA - TOCOPHEROL - PHENOXYETHANOL - IRON OXIDES - TITANIUM DIOXIDE - MICA

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503


SYNCHRO SKIN
SELF-REFRESHING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL 1 FL. OZ.



SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 430 CEDAR

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-504
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	1.69 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	0.345 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
GLYCERIN (UNII: PDC6A3C0OX)	
PHENYL TRIMETHICONE (UNII: DR0K5NOJ4R)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
DIMETHICONE CROSSPOLYMER (45000 MPAS AT 12% IN CYCLOPENTASILOXANE) (UNII: UF7620L1W6)	
SORBITAN SESQUISO STEARATE (UNII: VU97D01BF9)	
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
SILODRATE (UNII: 9T3UU8T0QK)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
THYMUS SERPYLLUM (UNII: 86H4S6K51N)	
SODIUM ACETYLATED HYALURONATE (UNII: WN66R7GL93)	
CYCLOMETHICONE 5 (UNII: 0THT5PC10R)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
POLYMETHYLSILSESQUOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	

STEARIC ACID (UNII: 4ELV7Z65AP)	
LAPONITE (UNII: D703131383)	
PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
PHENOXYETHANOL (UNII: HE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-504-10	1 in 1 CARTON	09/01/2019	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC MONOGRAPH FINAL	part352	09/01/2019		

SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 440 AMBER
 octinoxate and titanium dioxide emulsion

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-505	
Route of Administration	TOPICAL			

Active Ingredient/Active Moiety			
	Ingredient Name	Basis of Strength	Strength
	OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	1.69 g in 30 mL
	TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	0.345 g in 30 mL

Inactive Ingredients		
	Ingredient Name	Strength
	DIMETHICONE (UNII: 92RU3N3Y1O)	
	WATER (UNII: 059QF0KO0R)	
	GLYCERIN (UNII: PDC6A3C0OX)	
	PHENYL TRIMETHICONE (UNII: DR0K5NOJ4R)	
	PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	

BUTYLENE GLYCOL (UNII: 3XUS85K0RA)
DIMETHICONE CROSSPOLYMER (450000 MPAS AT 12% IN CYCLOPENTASILOXANE) (UNII: UF7620L1W6)
SORBITAN SESQUISO STEARATE (UNII: VU97D01BF9)
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)
SILODRATE (UNII: 9T3UU8T0QK)
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)
THYMUS SERPYLLUM (UNII: 86H4S6K51N)
SODIUM ACETYLATED HYALURONATE (UNII: WN66R7GL93)
CYCLOMETHICONE 5 (UNII: 0TH5PC10R)
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)
POLYMETHYLSILSESQUOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)
STEARIC ACID (UNII: 4ELV7Z65AP)
LAPONITE (UNII: D703131383)
PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL)
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)
PHENOXYETHANOL (UNII: HIE492ZZ3T)
FERRIC OXIDE RED (UNII: 1K09F3G675)
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)
FERROSFERRIC OXIDE (UNII: XM0M87F357)
MICA (UNII: V8A1AW0880)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-505-10	1 in 1 CARTON	09/01/2019	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part352	09/01/2019	

SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 450 COPPER

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-506
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	1.69 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	0.345 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
GLYCERIN (UNII: PDC6A3C0OX)	
PHENYL TRIMETHICONE (UNII: DR0K5NOJ4R)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
DIMETHICONE CROSSPOLYMER (450000 MPAS AT 12% IN CYCLOPENTASILOXANE) (UNII: UF7620L1W6)	
SORBITAN SESQUISO STEARATE (UNII: VU97D01BF9)	
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
SILODRATE (UNII: 9T3UU8T0QK)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
THYMUS SERPYLLUM (UNII: 86H4S6K51N)	
SODIUM ACETYLATED HYALURONATE (UNII: WN66R7GL93)	
CYCLOMETHICONE 5 (UNII: 0THT5PC10R)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
POLYMETHYLSILSESQUOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
LAPONITE (UNII: D703131383)	
PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-506-10	1 in 1 CARTON	09/01/2019	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part352	09/01/2019	

SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 460 TOPAZ

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-507
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	1.69 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	0.345 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
GLYCERIN (UNII: PDC6A3C0OX)	
PHENYL TRIMETHICONE (UNII: DR0K5NOJ4R)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
DIMETHICONE CROSSPOLYMER (45000 MPAS AT 12% IN CYCLOPENTASILOXANE) (UNII: UF7620L1W6)	
SORBITAN SESQUISO STEARATE (UNII: VU97D01BF9)	
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
SILODRATE (UNII: 9T3UU8T0QK)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
THYMUS SERPYLLUM (UNII: 86H4S6K51N)	
SODIUM ACETYLATED HYALURONATE (UNII: WN66R7GL93)	
CYCLOMETHICONE 5 (UNII: 0THT5PC10R)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
POLYMETHYLSILSESQUOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
LAPONITE (UNII: D703131383)	
PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	

PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSO FERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-507-10	1 in 1 CARTON	09/01/2019	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part352	09/01/2019	

SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 510 SUEDE

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-508
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	1.69 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	0.345 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
GLYCERIN (UNII: PDC6A3C0OX)	
PHENYL TRIMETHICONE (UNII: DR0K5NOJ4R)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
DIMETHICONE CROSSPOLYMER (450000 MPAS AT 12% IN CYCLOPENTASILOXANE) (UNII: UF7620L1W6)	
SORBITAN SESQUISO STEARATE (UNII: VU97D01BF9)	
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
SILODRATE (UNII: 9T3UU8T0QK)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	

THYMUS SERPYLLUM (UNII: 86H4S6K51N)
SODIUM ACETYLATED HYALURONATE (UNII: WN66R7GL93)
CYCLOMETHICONE 5 (UNII: 0TH5PC10R)
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)
POLYMETHYLSILSESQUOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)
STEARIC ACID (UNII: 4ELV7Z65AP)
LAPONITE (UNII: D703131383)
PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL)
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)
PHENOXYETHANOL (UNII: HIE492ZZ3T)
FERRIC OXIDE RED (UNII: 1K09F3G675)
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)
FERROSFERRIC OXIDE (UNII: XM0M87F357)
MICA (UNII: V8A1AW0880)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-508-10	1 in 1 CARTON	09/01/2019	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part352	09/01/2019	

SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 520 ROSEWOOD

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-509
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	1.69 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	0.345 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
GLYCERIN (UNII: PDC6A3C0OX)	
PHENYL TRIMETHICONE (UNII: DR0K5NOJ4R)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
DIMETHICONE CROSSPOLYMER (450000 MPAS AT 12% IN CYCLOPENTASILOXANE) (UNII: UF7620L1W6)	
SORBITAN SESQUISO STEARATE (UNII: VU97D01BF9)	
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
SILODRATE (UNII: 9T3UU8T0QK)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
THYMUS SERPYLLUM (UNII: 86H4S6K51N)	
SODIUM ACETYLATED HYALURONATE (UNII: WN66R7GL93)	
CYCLOMETHICONE 5 (UNII: 0THT5PC10R)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
LAPONITE (UNII: D703131383)	
PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-509-10	1 in 1 CARTON	09/01/2019	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part352	09/01/2019	

SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 530 HENNA

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-510
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	1.69 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	0.345 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
GLYCERIN (UNII: PDC6A3C0OX)	
PHENYL TRIMETHICONE (UNII: DR0K5NOJ4R)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
DIMETHICONE CROSSPOLYMER (450000 MP.A.S AT 12% IN CYCLOPENTASILOXANE) (UNII: UF7620L1W6)	
SORBITAN SESQUISO STEARATE (UNII: VU97D01BF9)	
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
SILODRATE (UNII: 9T3UU8T0QK)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
THYMUS SERPYLLUM (UNII: 86H4S6K51N)	
SODIUM ACETYLATED HYALURONATE (UNII: WN66R7GL93)	
CYCLOMETHICONE 5 (UNII: 0THT5PCI0R)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
POLYMETHYLSILSESQUOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
LAPONITE (UNII: D703131383)	
PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSO FERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-510-10	1 in 1 CARTON	09/01/2019	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part352	09/01/2019	

SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 540 MAHOGANY

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-511
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	1.69 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	0.345 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
GLYCERIN (UNII: PDC6A3C0OX)	
PHENYL TRIMETHICONE (UNII: DR0K5NOJ4R)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
DIMETHICONE CROSSPOLYMER (45000 MPAS AT 12% IN CYCLOPENTASILOXANE) (UNII: UF7620L1W6)	
SORBITAN SESQUISO STEARATE (UNII: VU97D01BF9)	
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
SILODRATE (UNII: 9T3UU8T0QK)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
THYMUS SERPYLLUM (UNII: 86H4S6K51N)	
SODIUM ACETYLATED HYALURONATE (UNII: WN66R7GL93)	
CYCLOMETHICONE 5 (UNII: 0THT5PC10R)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	

STEARIC ACID (UNII: 4ELV7Z65AP)
LAPONITE (UNII: D703131383)
PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL)
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)
PHENOXYETHANOL (UNII: HIE492ZZ3T)
FERRIC OXIDE RED (UNII: 1K09F3G675)
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)
FERROSFERRIC OXIDE (UNII: XM0M87F357)
MICA (UNII: V8A1AW0880)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-511-10	1 in 1 CARTON	09/01/2019	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part352	09/01/2019	

SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 550 JASPER

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-512
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	1.69 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	0.345 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
GLYCERIN (UNII: PDC6A3C0OX)	
PHENYL TRIMETHICONE (UNII: DR0K5NOJ4R)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	

BUTYLENE GLYCOL (UNII: 3XUS85K0RA)
DIMETHICONE CROSSPOLYMER (450000 MPA.S AT 12 % IN CYCLOPENTASILOXANE) (UNII: UF7620L1W6)
SORBITAN SESQUISO STEARATE (UNII: VU97D01BF9)
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)
SILODRATE (UNII: 9T3UU8T0QK)
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)
THYMUS SERPYLLUM (UNII: 86H4S6K51N)
SODIUM ACETYLATED HYALURONATE (UNII: WN66R7GL93)
CYCLOMETHICONE 5 (UNII: 0THT5PC10R)
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)
POLYMETHYLSILSESQUOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)
STEARIC ACID (UNII: 4ELV7Z65AP)
LAPONITE (UNII: D703131383)
PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL)
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)
PHENOXYETHANOL (UNII: HIE492ZZ3T)
FERRIC OXIDE RED (UNII: 1K09F3G675)
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)
FERROSFERRIC OXIDE (UNII: XM0M87F357)
MICA (UNII: V8A1AW0880)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-512-10	1 in 1 CARTON	09/01/2019	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part352	09/01/2019	

SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 560 OBSIDIAN

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-513
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	1.69 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	0.345 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0K00R)	
GLYCERIN (UNII: PDC6A3C0OX)	
PHENYL TRIMETHICONE (UNII: DR0K5NOJ4R)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
DIMETHICONE CROSSPOLYMER (45000 MPAS AT 12% IN CYCLOPENTASILOXANE) (UNII: UF7620L1W6)	
SORBITAN SESQUISO STEARATE (UNII: VU97D01BF9)	
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
SILODRATE (UNII: 9T3UU8T0QK)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
THYMUS SERPYLLUM (UNII: 86H4S6K51N)	
SODIUM ACETYLATED HYALURONATE (UNII: WN66R7GL93)	
CYCLOMETHICONE 5 (UNII: 0THT5PC10R)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
POLYMETHYLSILSESQUOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
LAPONITE (UNII: D703131383)	
PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-513-10	1 in 1 CARTON	09/01/2019	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part352	09/01/2019	

Labeler - SHISEIDO AMERICAS CORPORATION (193691821)

Establishment

Name	Address	ID/FEI	Business Operations
SHISEIDO AMERICAS INC.		782677132	MANUFACTURE(58411-504, 58411-505, 58411-506, 58411-507, 58411-508, 58411-509, 58411-510, 58411-511, 58411-512, 58411-513) , ANALYSIS(58411-504, 58411-505, 58411-506, 58411-507, 58411-508, 58411-509, 58411-510, 58411-511, 58411-512, 58411-513)

Revised: 2/2020

SHISEIDO AMERICAS CORPORATION