

LACTATED RINGERS- lactated ringers injection, solution

Aspen Veterinary Resources

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, [click here](#).

Lactated Ringer's Injection, USP

For Animal Use Only

Sterile Nonpyrogenic Solution

DESCRIPTION

Lactated Ringer's Injection is a sterile, nonpyrogenic solution for fluid and electrolyte replenishment in single dose containers for parenteral administration. It contains no antimicrobial agents.

COMPOSITION:

Each 100 mL of intravenous infusion contains:

Sodium chloride 600 mg

Sodium lactate 310 mg

Potassium chloride 30 mg

Calcium chloride dihydrate 20 mg

Water for injection q.s.

Electrolytes: The calcium, potassium and sodium contents are approximately 2.7, 4 and 130 milliequivalents per liter, respectively. Osmolarity: 273 mOsm/l

CLINICAL PHARMACOLOGY

A multiple electrolyte intravenous solution is intended to restore the electrolyte balance and water for hydration. A combination of multiple electrolytes and sodium lactate, an alkalinizing agent, will provide electrolyte balance and normalize the pH of the acid-base of the physiological system.

INDICATIONS

Lactated Ringer's Injection is indicated for the correction of electrolyte depletion, metabolic acidosis and dehydration of cattle, calves, horses, sheep and swine.

CONTRAINDICATIONS

Lactated Ringer's Injection is contraindicated in patients with a known hypersensitivity to sodium lactate; congestive heart failure or severe impairment of renal function; clinical states in which the administration of sodium and chloride is detrimental.

WARNINGS

The introduction of additives to any solution, regardless of type of container, requires special attention to ensure that no incompatibilities result. While some incompatibilities are readily absorbed, one must be aware that subtle physical, chemical and pharmacological incompatibilities can occur. The medical literature, the package insert and other available sources of information should be reviewed for thorough understanding of possible incompatibilities.

Lactated Ringer's Injection should be used with great care, if at all, in patients with congestive heart failure, severe renal insufficiency, and in clinical states in which there exists edema and sodium retention.

Lactated Ringer's Injection should be used with great care, if at all, in patients with hyperkalemia, severe renal failure, and in conditions in which potassium retention is present.

Lactated Ringer's Injection should be used with great care in patients with metabolic or respiratory alkalosis. The administration of lactate ions should be done with great care in those conditions in which there is an increased level or an impaired utilization of these ions, such as severe hepatic insufficiency.

Lactated Ringer's Injection should not be administered simultaneously with blood through the same administration set because of likelihood of coagulation.

The intravenous administration of Lactated Ringer's Injection can cause fluid and/or solute overloading resulting in dilution of serum electrolyte concentrations, over-hydration, congested states, or pulmonary edema. The risk of dilutional states is inversely proportional to the electrolyte concentrations of the injections. The risk of solute overloading causing congested states with peripheral and pulmonary edema is directly proportional to the electrolyte concentrations of the injections.

In patients with diminished renal function, administration of Lactated Ringer's Injection may result in sodium or potassium retention.

Do not administer to horses by intraperitoneal injection. Do not administer to animals with inadequate renal function. Not for use in lactic acidosis.

ADVERSE REACTIONS

Adverse reactions may occur due to the solution or the technique of administration including fever response, infection at the site of injection or allergic reactions. Prolonged intravenous infusion of this type of product may cause venous thrombosis or phlebitis extending from the site of injection, extravasation, and hypervolemia.

If an adverse reaction does occur, discontinue the infusion and evaluate the patient, institute appropriate therapeutic countermeasures, and save the remainder of the fluid for examination if deemed necessary.

PRECAUTIONS

This is a single dose unit. It contains no preservatives. Use entire contents when first opened.

Clinical evaluation and periodic laboratory determinations are necessary to monitor changes in fluid balance, electrolyte concentrations, and acid base balance during prolonged therapy or whenever the condition of the patient warrants such evaluation.

Lactated Ringer's Injection should be used with caution. Excess administration may result in metabolic alkalosis.

Do not administer unless solution is clear and both seal and container are intact.

Solution must be warmed to body temperature prior to administration and administered at a slow rate. Use solution promptly following initial entry.

Reactions which may occur because of the solution or the technique of administration, include febrile response, infection at the site of injection, extravasation, and hypervolemia.

If an adverse reaction does occur, discontinue the infusion and evaluate the patient, institute appropriate therapeutic countermeasures, and save the remainder of the fluid for examination if deemed necessary.

DOSAGE AND ADMINISTRATION

To be used as directed by a licensed veterinarian. Dosage is dependent upon the age, weight and clinical conditions of the patient as well as laboratory determinations.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration.

For use in one patient on one occasion only. Discard any unused portion. Care should be taken with administration technique to avoid administration site reactions and infection.

Additives may be incompatible. Complete information is not available. Those additives known to be incompatible should not be used. Consult with pharmacist, if available. If, in the informed judgement of the doctor, it is deemed advisable to introduce additives, use aseptic technique. Mix thoroughly when additives have been introduced. Do not store solutions containing additives.

OVERDOSAGE:

In an event of over-hydration or solute overload, re-evaluate the patient and institute appropriate corrective measures. See Warnings, Precautions and Adverse Events.

PACKAGING

1000 mL bag with clear overwrap: NDC 46066-350-06

3000 mL bag with clear overwrap: NDC 46066-350-08

5000 mL bag with clear overwrap: NDC 46066-350-09

STORAGE:

Store below 25°C (77°F).

Keep out of reach of children.

Directions for Use of Plastic Container:


To Open

Tear overwrap at slit and remove solution container. Some opacity of the plastic due to moisture absorption during the sterilization process may be observed. This is normal and does not affect the solution quality or safety. The opacity will diminish gradually. Check for minute leaks by squeezing solution container firmly. If leaks are found, discard solution as sterility may be impaired. If supplemental medication is desired, follow directions below:

Preparation for Administration

(Use Aseptic Technique)

1. Close flow control clamp of administration set.
2. Twist off the protector cap from the administration port of the bags.



3. Insert the spike, a slight resistance should be felt as the port membrane is broken.

Use a twisting motion during insertion. Insert the spike until the shoulder of the spike is level with the port.

NOTE: See full directions on administration set carton.

4. Suspend container from hanger.

5. Squeeze and release drip chamber to establish proper fluid level in chamber.

6. Open flow control clamp and clear air from set. Close clamp.

7. Attach set to venipuncture device. If device is not indwelling, prime and make venipuncture.

8. Regulate rate of administration with flow control clamp.

WARNING: Do not use flexible container in series connections.

To Add Medication

WARNING: Additives may be incompatible.

To add medication before solution administration

1. Prepare medication site.
2. Using syringe with 0.63 mm to 0.80 mm needle, puncture medication port and inject.
3. Mix solution and medication thoroughly. For high density medication such as potassium chloride, squeeze ports while ports are upright and mix thoroughly.

To add medication during solution administration

1. Close the clamp on the administration set.
2. Prepare medication site.
3. Using syringe with 0.63 mm to 0.80 mm needle, puncture medication port and inject.
4. Remove container from IV pole and/or turn to an upright position.
5. Evacuate both ports by squeezing them while container is in the upright position.
6. Mix solution and medication thoroughly.
7. Return container to in use position and continue administration.

CAUTION:

Federal law restricts this drug to use by or on the order of a licensed veterinarian.

Manufactured for:

Aspen Veterinary Resources, Ltd.

Liberty, MO 64068

www.aspenveterinaryresources.com

Manufactured by:

INFOMED FLUIDS S.R.L., Bucharest, Romania

Revised July 2016

PACKAGE/LABEL PRINCIPAL DISPLAY PANEL

NDC 46066-350-06

Lactated Ringer's Injection, USP

1000 mL

NDC 46066-350-06
Lactated Ringer's Injection, USP

1000 mL

**Sterile Nonpyrogenic Solution
For Animal Use Only**

COMPOSITION

Each 100 mL of intravenous infusion contains:

Sodium chloride	600 mg
Sodium lactate	310 mg
Potassium chloride	30 mg
Calcium chloride dihydrate	20 mg
Water for injection	q.s.

Electrolytes: The calcium, potassium and sodium contents are approximately 2.7, 4 and 130 milliequivalents per liter, respectively.

Osmolarity: 273 mOsm/l

INDICATIONS: For the correction of electrolyte depletion, metabolic acidosis and dehydration of cattle, calves, horses, sheep and swine.


DOSAGE AND ADMINISTRATION: As directed by a veterinarian. Dosage is dependent upon the age, weight and clinical condition of the patient, as well as laboratory determinations. Administer intravenously, subcutaneously, or intraperitoneally (except in horses) using strict aseptic technique. See package insert.

CAUTION: Solution must be warmed to body temperature prior to administration and administered at a slow rate. This is a single dose unit. It contains no preservatives. Use solution promptly following initial entry. Use entire contents when first opened. Squeeze and inspect inner bag which maintains product sterility. Discard if leaks are found or if the solution contains visible solid particles. Do not administer simultaneously with blood. Do not use unless solution is clear and seal is intact.

WARNING: Do not administer to horses by intraperitoneal injection. Do not administer to animals with inadequate renal function. Not for use in lactic acidosis. Additives may be incompatible. Consult with pharmacist if available. When introducing additives use aseptic technique. Mix thoroughly. If entire contents are not used, discard the unused portion. Store below 25°C (77°F).


CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

Keep out of reach of children.



Manufactured for:
Aspen Veterinary Resources®, Ltd.
Liberty, MO 64068
www.aspenveterinaryresources.com

Manufactured by:
Infomed Fluids S.R.L., Bucharest, Romania
Rev 07/16. A5261CH



LOT NUMBER: 0000000 EXP. DATE: MM YYYY

LACTATED RINGERS

lactated ringers injection, solution

Product Information

Product Type	PRESCRIPTION ANIMAL DRUG	Item Code (Source)	NDC:46066-350
Route of Administration	INTRAVENOUS, SUBCUTANEOUS, INTRAPERITONEAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X) (CHLORIDE ION - UNII:Q32ZN48698)	SODIUM CHLORIDE	600 mg in 100 mL
SODIUM LACTATE (UNII: TU7HW0W0QT) (LACTIC ACID, UNSPECIFIED FORM - UNII:33X04XA5AT)	SODIUM LACTATE	310 mg in 100 mL
POTASSIUM CHLORIDE (UNII: 660YQ98I10) (POTASSIUM CATION - UNII:295O53K152)	POTASSIUM CHLORIDE	30 mg in 100 mL
CALCIUM CHLORIDE (UNII: M4I0D6VV5M) (CALCIUM CATION - UNII:2M83C4R6ZB)	CALCIUM CHLORIDE	20 mg in 100 mL

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:46066-350-06	1000 mL in 1 BAG		
2	NDC:46066-350-08	3000 mL in 1 BAG		
3	NDC:46066-350-09	5000 mL in 1 BAG		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
Unapproved drug other		01/31/2017	

Labeler - Aspen Veterinary Resources (627265361)

Revised: 1/2017

Aspen Veterinary Resources