

**ANTACID MINT- aluminum hydroxide, magnesium hydroxide,  
demethicone suspension  
Walgreens**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Drug Facts**

**Active Ingredients**

Aluminum hydroxide (equiv. to dried gel, USP) 200 mg  
Magnesium hydroxide 200 mg  
Simethicone 20 mg

**Purpose**

Antacid

Antacid

Antigas

**Uses**

- For the relief of
  - acid indigestion
  - heartburn
  - sour stomach
  - upset stomach associated with these symptoms
  - pressure and bloating commonly referred to as gas

**Warnings**

**Do not take**

more than 16 teaspoonfuls in a 24-hour period or use the maximum dosage for more than 2 weeks except under the advice and supervision of a physician

**Ask a doctor before use if you have**

- kidney disease • a magnesium restricted diet

**Ask a doctor or pharmacist before use if you are**

presently taking a prescription drug. Antacids may interact with certain prescription drugs.

**Stop use and ask a doctor if**

symptoms last for more than 2 weeks

**If pregnant or breast-feeding**

ask a health professional before use.

**Keep out of the reach of children.**

In case of overdose, get medical help or contact a Poison Controls Center (1-800-222-1222) right away.

**Directions**

- shake well before using
- do not take more than 16 teaspoonfuls in 24 hours or use the maximum dosage for more than 2 week
- adults and children 12 years of age and older: take 2 to 4 teaspoonfuls four times a day or as directed by a physican
- children under 12 years of age: do not use

**Other Information**

- each teaspoon (5 mL) contains: magnesium 80 mg, potassium 5 mg, and sodium 2 mg
- store between 20-25°C (68-77°F). Avoid freezing

**Inactive Ingredients**

benzyl alcohol, carboxymethylcellulose sodium, flavors, glycerin, microcrystalline cellulose, potassium citrate, propylene glycol, purified water, sorbitol, saccharin sodium xanthan gum

**Principal Display Panel**

Compare to Maalox® Advanced Regular Strength Active ingredient<sup>††</sup>

Antacid +

Anti-Gas Liquid

ALUMINUM HYDROXIDE 200 mg / ANTACID

MAGNESIUM HYDROXIDE 200 mg / ANTACID

SIMETHICONE 20 mg /ANTI-GAS

REGULAR STRENGTH

- Fast relief of heartburn, pressure, bloating & acid indigestion

FL OZ (mL)

MINT FLAVOR

††This product is not manufactured or distributed by Novartis Consumer Health, Inc., distributor of Maalox® Advanced Regular Strength.

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.**

**DISTRIBUTED BY: WALGREENS CO.,  
200 WILMOT RD., DEERFIELD, IL 60015**

walgreens.com

## Package Label



Compare to Maalox® Advanced Regular Strength Active Ingredients††

NDC 0363-0432-26

# Antacid + Anti-Gas Liquid

ALUMINUM HYDROXIDE 200 mg / ANTACID  
MAGNESIUM HYDROXIDE 200 mg / ANTACID  
SIMETHICONE 20mg / ANTI-GAS

**REGULAR STRENGTH**

- Fast relief of heartburn, pressure, bloating & acid indigestion

26 FL OZ (769 mL)



MINT FLAVOR

PLD-A457B LB005702

Drug Facts	
Active ingredients (in each 5 mL = 1 teaspoon)	Purposes
Aluminum hydroxide (equiv. to dried gel USP) 200 mg.....	Antacid
Magnesium hydroxide 200 mg.....	Antacid
Simethicone 20 mg.....	Antigas

**Uses** ■ for the relief of

- acid indigestion ■ heartburn ■ sour stomach
- upset stomach associated with these symptoms
- pressure and bloating commonly referred to as gas

**Warnings**

Do not take more than 16 teaspoonfuls in a 24-hour period or use the maximum dosage for more than 2 weeks except under the advice and supervision of a physician.

Ask a doctor before use if you have

- kidney disease ■ a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are presently taking a prescription drug. Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if symptoms last more than 2 weeks.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions** ■ shake well before using

- do not take more than 16 teaspoonfuls in 24 hours or use the maximum dosage for more than 2 weeks

**Drug Facts (continued)**

- adults and children 12 years of age and older: take 2 to 4 teaspoonfuls four times a day or as directed by a physician
- children under 12 years of age: do not use

**Other information**

- each teaspoon (5 mL) contains: magnesium 80 mg, potassium 5 mg, and sodium 2 mg
- store between 20-25°C (68-77°F). Avoid freezing.

**Inactive ingredients** benzyl alcohol, carboxymethylcellulose sodium, flavor, glycerin, microcrystalline cellulose, potassium citrate, propylene glycol, purified water, saccharin sodium, sorbitol, xanthan gum

**Questions or comments?** Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.**

Walgreens Pharmacist Recommended  
Walgreens Pharmacist Survey

††This product is not manufactured or distributed by Novartis Consumer Health, Inc., distributor of Maalox® Advanced Regular Strength.

DISTRIBUTED BY: WALGREEN CO.,  
200 WILMOT RD., DEERFIELD, IL 60015  
100% SATISFACTION GUARANTEED  
walgreens.com ©2018 Walgreen Co.  
ITEM 272286 W10023-0418-F



3 11917 20012 5  
PLD-A457B LB005703 ORG0418-F2

**WALGREENS Antacid and Anti-Gas**

## ANTACID MINT

aluminum hydroxide, magnesium hydroxide, demithicone suspension

## Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-0432	
Route of Administration	ORAL			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0) (ALUMINUM HYDROXIDE - UNII:5QB0T2IUN0)		ALUMINUM HYDROXIDE	200 mg in 5 mL	
MAGNESIUM HYDROXIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838)		MAGNESIUM HYDROXIDE	200 mg in 5 mL	
DIMETHICONE (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)		DIMETHICONE	20 mg in 5 mL	
Inactive Ingredients				
Ingredient Name			Strength	
BENZYL ALCOHOL (UNII: LKG8494WBH)				
GLYCERIN (UNII: PDC6A3C0OX)				
POTASSIUM CITRATE (UNII: EE90ONI6FF)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
WATER (UNII: 059QF0KO0R)				
SORBITOL (UNII: 506T60A25R)				
SACCHARIN SODIUM ANHYDROUS (UNII: I4807BK602)				
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)				
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)				
XANTHAN GUM (UNII: TTV12P4NEE)				
Product Characteristics				
Color		Score		
Shape		Size		
Flavor	MINT	Imprint Code		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0432-26	769 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/31/2018	12/29/2023
2	NDC:0363-0432-12	355 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/31/2018	12/29/2023
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
OTC monograph final	part332		03/31/2018	12/29/2024

---

**Labeler** - Walgreens (008965063)

Revised: 1/2023

Walgreens